
Participant Flow:

Participant flow diagram for the pilot implementation follow-up



Baseline Characteristics:

Participant characteristics for the pilot study follow-up									
		Young people			Men who have sex with men				
		Total (N, %)		Recruits (N, %)		Total (N, %)		Recruits (N, %)	
		n = 6,216*		n = 273		n = 5,738*		n = 133	
Gender	Male	1,444	(23%)	67	(25%)				
	Female	4,772	(77%)	206	(75%)				
Age	16-20 years	1,890	(30%)	86	(32%)	149	(3%)	9	(7%)
	21-25 years	4,326	(70%)	69	(68%)	610	(11%)	25	(19%)
	26-35 years					1,639	(29%)	43	(32%)
	36-45 years					1,504	(26%)	29	(22%)
	>45 years					1,836	(32%)	27	(20%)
Ethnicity	White	4,296	(69%)	179	(66%)	4,485	(78%)	102	(77%)
	Mixed	455	(7%)	25	(9%)	228	(4%)	11	(8%)
	Asian	331	(5%)	11	(4%)	350	(6%)	7	(5%)
	Black	568	(9%)	52	(19%)	192	(3%)	6	(5%)
	Other	566	(9%)	6	(2%)	483	(8%)	7	(5%)
Sexual	Heterosexual	5,759	(93%)	245	(90%)				
orientation	Homosexual	25	(0%)	2	(1%)	5,286	(92%)	113	(85%)
	Bisexual	404	(7%)	25	(9%)	452	(8%)	20	(15%)

Outcome Measures:

The primary outcome measures were measured using mixed-methods during a pilot implementation period.

Planned outcome	Status	Result				
Acceptability of the intervention to users a		neouit				
Proportion of eligible service users who attend the clinic that were assigned a score by the triage tool	Collected as planned	1. 19% in young people; 10% in men who have sex with men				
2. Proportion of those who were classified as high risk who are offered the intervention3. Proportion of those who were offered	Partially collected	2. Unable to fully assess3. Unable to fully assess4. Only able to qualitatively assess				
the intervention who took up the intervention		u33C33				
Proportion who took up the intervention who completed the intervention						
5. Reasons for not completing the intervention from the qualitative study of participants	Collected as planned	16 qualitative interviews conducted with service users				
6. Acceptability of the intervention from the qualitative study of the staff	Partially completed	2 focus group discussions and 1 qualitative interview conducted with healthcare providers				
Feasibility of delivering the interventions						
The total time spent by service users within the clinical service compared to normal	Data not collected	N/A				
Total number of service users seen and STIs diagnosed, compared to normal						
Average consultation time compared to normal						
Number of patients seen by health advisors compared to normal						
5. Extra HCP time required for the intervention	Partially collected	Health advisors recorded duration of intervention sessions.				
Feasibility of obtaining follow-up outcome data						
 Proportion of eligible service users who consented to the follow-up Proportion of eligible service users who 	Collected as planned	1. 69% 2. 35% 3. 28%				
were contactable at 6 weeks and complete a questionnaire		3: -5/5				
Proportion who complete follow-up tests						

Adverse Events:

There were no adverse events associated with this trial.