


1. Participant Flow

Figure 1 Participant flow through the trial



2. Baseline Characteristics

Table 1 Characteristics of the study population at baseline by treatment group

	Total (n=1355)		Control (n=714)		Intervention (n=641)		P-value ^a
	n	%	N	%	n	%	
Sex							
Male	623	46.0	322	45.1	301	47.0	0.493
Female	732	54.0	392	54.9	340	53.0	
Age							
13	7	0.5	5	0.7	2	0.3	<0.001
14	472	34.8	255	35.7	217	33.9	
15	702	51.8	397	55.6	305	47.6	
16	170	12.6	57	8.0	113	17.6	
17	4	0.3	0	0.0	4	0.6	
Grade							
8 th	686	50.6	348	48.7	338	52.7	0.142
9 th	669	49.4	366	51.3	303	47.3	
Perceived family affluence							
Very well off	308	22.7	179	25.1	129	20.1	0.192
Quite well off	487	35.9	256	35.9	231	36.0	
Average	524	38.7	263	36.8	261	40.7	
Not well off ^b	36	2.6	16	2.2	20	3.1	
Binge drinking within the last 30 days							
0-3 times	1301	96.1	689	96.5	612	95.6	0.408
4 or more times	53	3.9	25	3.5	28	4.4	
Perception of lifetime binge drinking among peers							
No overestimation	743	55.1	429	60.3	314	49.4	<0.001
Overestimation	605	44.9	283	39.7	322	50.6	
Alcohol-related harms^c							
0-1	395	48.5	170	40.6	225	56.8	0.819
2 or more	145	17.8	64	15.3	81	20.5	
Missing ^d	275	33.7	185	44.1	90	22.7	<0.001

^a Chi²-test for differences between intervention and control group.

^b Response options Not so well off and Not at all well off combined

^c Only measured among pupils with lifetime alcohol use (n=815)

^d Data missing due to an error in the electronic questionnaire in the 2016 data collection

3. Outcome Measures

Table 2 Intervention effects at 3-months follow-up

Outcomes	OR^b	95% CI	P-value	ICC^c	Control n/N	Intervention n/N
Overestimation of peers' lifetime binge drinking ^a	0.52	0.33;0.83	0.006	0.088	323/706	241/633
Binge drinking (4 or more times) within the last 30 days	0.89	0.49;1.61	0.709	0.099	77/713	63/640
Alcohol-related harms (2 or more) (n=540)	0.59	0.37;0.93	0.024	0.004	89/234	95/306

4. Adverse Events

There were no adverse events associated with this trial.