


Simultaneous Endo-Epicardial Mapping of Recurrent Atrial Fibrillation (SEE MORE AF)

Results

Trial registration number: <https://doi.org/10.1186/ISRCTN10615430>

Study Numbers



Demographics

	Number	Percentage
Male	13	65
Female	7	35
70-80 years	8	40
60-70 years	7	35
50-60 yeas	4	20
40-50 years	1	5

Variable	All patients (n=20)
Age, y	65.7 ± 8
Male	13 (65)

Body Mass Index (SD)	31.6 ± 6
Hypertension	12 (60)
Diabetes Mellitus	1 (5)
Coronary Artery Disease	8 (40)
Renal Failure	0
Ischaemic Cardiomyopathy	1 (5)
History of MI	1 (5)
History of Stroke	0
CHA₂DS₂VASc score	2 (IQR 2-3)
Anticoagulation	
Rivaroxaban	13 (65)
Apixaban	5 (25)
Dabigatran	2 (10)
EHRA score pre-procedure	2 (IQR 2-3)
LVEF	
>50%	10 (50)
40-49%	3 (15)
30-39%	5 (25)
<30%	2 (10)
Left atrial dimension (mm) (SD)	48.1 ± 8
Left atrial volume (mls) (SD)	92.4 ± 29
Previous Cryoablation for AF	19 (95)
Previous Radiofrequency Ablation for AF	2 (10)
Long Standing persistent AF	10 (50)

Adverse Events

	Number	Expected	Related	Outcome
Prolonged hospitalisation	3	2	3	Resolved
Hospital admission	2	1	1	Resolved

Outcome Measures

	6 Months Number	6 Months Percentage	12 Months Number	12 Months Percentage
Primary: Endo-epicardial AF free survival	18	90	16	80
Linear lesion acute block/transmural: Roof line	19	95	N/A	N/A
Linear lesion acute block/transmural: Inferior line	19	95	N/A	N/A
Linear lesion acute block/transmural: Anterior mitral line	20	100	N/A	N/A

