


ISRCTN82492510


Basic results summary

Participant flow



Analyses		
<p>Completed self-report survey</p> <ul style="list-style-type: none">• Baseline (7/12, 75%)• Mid-point (7/11, 64%)• End of intervention (5/11, 45%)• 6-month post intervention (3/9, 33%) <p>CRF</p> <ul style="list-style-type: none">• Baseline (100%)• 6 month (100%)• 12 month (100%) <p>Qualitative interview</p> <ul style="list-style-type: none">• Completed (n=5)• Declined (n=4)• Not applicable (n=3)		<p>Completed self-report survey</p> <ul style="list-style-type: none">• Pre intervention (7/12, 58%)• Baseline (5/12, 42%)• Mid-point (5/12, 42%)• End of intervention (5/12, 42%) <p>CRF</p> <ul style="list-style-type: none">• Baseline (100%)• 6 month (100%)• 12 month (100%) <p>Qualitative interview</p> <ul style="list-style-type: none">• Completed (n=7)• Declined (n=5)

Baseline characteristics



Outcome measures

Outcome measure	Result
Consent to screen rate	95%
Screening positivity rate	24%
Recruitment rate, calculated by assessing the proportion of those who screen positive, that complete the consent form	46%
Randomisation acceptability, from participant feedback at randomisation and qualitative interviews	No study withdrawals Not a theme in qualitative interviews
Retention rate	87%
Engagement and adherence with intervention components	See next slide
Completion rates of self-report measures	67% → 52% → 43% → 38%

Adverse events

- 3 deaths, unrelated to trial
 - Withdrawal from treatment
- Suicide risk protocol has not been required to date
 - One patient over PHQ threshold and declined Talkworks; screening scores have subsequently improved