Basic results summary for CASNET2 (ISRCTN15913081)

Participant flow



Baseline characteristics

	All eligible patients	Cancer patients in pre-intervention period	Cancer patients in post-intervention period
Eligible patient population (n)	442,662	6,300	3,503
Female (n, %)	224,527 (50.7%)	2,967 (47.1%)	1,761 (50.3%)
Age in years (mean, SD)	45.9 (18.3)	65.1 (14.1)	64.5 (14.2)
Ethnicity (n, %)			
- White	323,232 (73.0%)	5,511 (87.5%)	2,982 (85.1%)
- Black	20,644 (4.7%)	144 (2.3%)	77 (2.2%)
- Asian	33,808 (7.6%)	112 (1.8%)	80 (2.3%)
- Mixed	7,877 (1.8%)	31 (0.5%)	20 (0.6%)
- Other	5,213 (1.2%)	22 (0.3%)	21 (0.6%)
- Missing	51,888 (11.7%)	480 (7.6%)	323 (9.2%)
IMD (n, %)			
- 1st quintile	75,879 (17.1%)	683 (10.8%)	468 (13.4%)
- 2nd quintile	109,365 (24.7%)	1,041 (16.5%)	713 (20.4%)
- 3rd quintile	106,708 (24.1%)	1,640 (26.0%)	941 (26.9%)
- 4th quintile	88,802 (20.1%)	1,752 (27.8%)	815 (23.3%)
- 5th quintile	61,908 (14.0%)	1,184 (18.8%)	566 (16.2%)
Urban (n,%)	317,747 (71.8%)	3,904 (62.0%)	2,208 (63.0%)

IMD: index of multiple deprivation

Outcome measures

Results of mixed effects modelling for the primary outcome measure (primary care interval)

These results will be updated when analyses are complete.

Adverse events

There were no adverse events associated with this study.