

ISRCTN36033130: Assessing the use of FreeStyle Libre Pro with HbA1c in diabetes management

1. Study Flow




Figure 1 Study flow diagram

2. Baseline characteristics

Variable	Baseline
Mean age \pm SD (years)	62.3 \pm 9.8
Sex, male	28 (63.6%)
Mean HbA1c \pm SD (%)	7.7 \pm 1.6
Mean duration of diabetes \pm SD (years)	8.2 \pm 6.3

Table 1 Baseline characteristics of study population (n=44)

3. Outcome measures

3.1 Primary outcome

		In your opinion, is the patient's current diabetes therapy optimised, based on the HbA1c value and the LibreView reports?		
		Yes	No	Total
In your opinion, is the patient's current diabetes therapy optimised, based on their HbA1c value?	Yes	18	5	23
	No	1	20	21 (48%)
	Total	19	25 (57%)	44

Table 2: HCP questionnaire responses

4. Adverse events

Adverse event	Number of patients affected	Number of events
Serious adverse events	0	0
Other (not including serious) adverse events	3 (6.8%)	5
Device related (anticipated)		
Erythema	3 (6.8%)	3
Pain	1 (2.3%)	1
Bleeding	1 (2.3%)	1

Table 3: Summary of adverse events recorded (n=44)