



Figure 1: Participant flow for the ASPECT trial

Table 2: Baseline characteristics by randomised group for all randomised participants (n=268)

| Characteristic | CBT (n=134) | OST (n=134) | All (n=268) |
|---|-------------------|-------------------|-------------------|
| Child's age | | | |
| N (%) | 134 (100%) | 134 (100%) | 268 (100%) |
| Mean (SD) | 11.8 (2.6) | 11.8 (2.6) | 11.8 (2.6) |
| Median (IQR) | 12.0 (10.0, 14.0) | 12.0 (10.0, 14.0) | 12.0 (10.0, 14.0) |
| Site | | | |
| Humber | 10 (7%) | 9 (7%) | 19 (7%) |
| Tees, Esk and Wear | 39 (29%) | 39 (29%) | 78 (29%) |
| South West Yorkshire | 9 (7%) | 5 (4%) | 14 (5%) |
| Norfolk and Suffolk | 21 (16%) | 35 (26%) | 56 (21%) |
| Northumberland | 14 (10%) | 10 (7%) | 24 (9%) |
| South Tyneside | 17 (13%) | 18 (13%) | 35 (13%) |
| Leeds | 1 (1%) | 0 (0%) | 1 (0%) |
| RDaSH | 10 (7%) | 5 (4%) | 15 (6%) |
| Exeter | 1 (1%) | 1 (1%) | 2 (1%) |
| Dorset | 5 (4%) | 6 (4%) | 11 (4%) |
| Cumbria | 1 (1%) | 1 (1%) | 2 (1%) |
| County Durham | 1 (1%) | 3 (2%) | 4 (1%) |
| Dudley and Walsall | 5 (4%) | 2 (1%) | 7 (3%) |
| Sex | | | |
| Boy | 53 (40%) | 48 (36%) | 101 (38%) |
| Girl | 81 (60%) | 86 (64%) | 167 (62%) |
| Ethnicity | | | |
| English / Welsh / Scottish / Northern Irish / British | 129 (96%) | 127 (95%) | 256 (96%) |
| Irish | 0 (0%) | 1 (1%) | 1 (0%) |

| Characteristic | CBT (n=134) | OST (n=134) | All (n=268) |
|--|----------------|----------------|----------------|
| Any other White background | 0 (0%) | 1 (1%) | 1 (0%) |
| Pakistani | 1 (1%) | 2 (1%) | 3 (1%) |
| White and Black Caribbean | 1 (1%) | 0 (0%) | 1 (0%) |
| White and Black African | 2 (1%) | 0 (0%) | 2 (1%) |
| White and Asian | 0 (0%) | 2 (1%) | 2 (1%) |
| Any other mixed / multiple ethnic background | 0 (0%) | 1 (1%) | 1 (0%) |
| Prefer not to say | 1 (1%) | 0 (0%) | 1 (0%) |
| School attender | | | |
| No | 8 (6%) | 5 (4%) | 13 (5%) |
| Yes | 126 (94%) | 129 (96%) | 255 (95%) |
| Treatment preference | | | |
| CBT | 35 (26%) | 43 (32%) | 78 (29%) |
| OST | 35 (26%) | 30 (22%) | 65 (24%) |
| No preference | 64 (48%) | 61 (46%) | 125 (47%) |
| Taking part in another study | | | |
| No | 132 (99%) | 130 (97%) | 262 (98%) |
| Yes - phobia/anxiety | 2 (1%) | 4 (3%) | 6 (2%) |
| Number of adults in household | | | |
| 1 | 18 (13%) | 25 (19%) | 43 (16%) |
| 2 | 100 (75%) | 101 (75%) | 201 (75%) |
| 3 | 13 (10%) | 6 (4%) | 19 (7%) |
| 4 | 2 (1%) | 2 (1%) | 4 (1%) |
| Number of children in household | | | |
| 1 | 37 (28%) | 36 (27%) | 73 (27%) |

| Characteristic | CBT (n=134) | OST (n=134) | All (n=268) |
|----------------|----------------|----------------|----------------|
| 2 | 65 (49%) | 68 (51%) | 133 (50%) |
| 3 | 28 (21%) | 21 (16%) | 49 (18%) |
| 4 | 2 (1%) | 8 (6%) | 10 (4%) |
| 5 | 2 (1%) | 0 (0%) | 2 (1%) |

Table 3: Baseline characteristics by randomised group for the parents and carers of all randomised participants (n=268)

| Characteristic | CBT (n=134) | OST (n=134) | All (n=268) |
|---|-------------------|-------------------|-------------------|
| Age | | | |
| N (%) | 132 (99%) | 133 (99%) | 265 (99%) |
| Mean (SD) | 41.3 (6.4) | 42.9 (7.2) | 42.1 (6.9) |
| Median (IQR) | 42.0 (37.0, 46.0) | 43.0 (37.0, 48.0) | 42.0 (37.0, 47.0) |
| Sex | | | |
| Male | 15 (11%) | 12 (9%) | 27 (10%) |
| Female | 119 (89%) | 122 (91%) | 241 (90%) |
| Ethnicity | | | |
| English / Welsh / Scottish / Northern Irish / British | 127 (95%) | 126 (94%) | 253 (94%) |
| Irish | 1 (1%) | 0 (0%) | 1 (0%) |
| Any other White background | 4 (3%) | 4 (3%) | 8 (3%) |
| Pakistani | 1 (1%) | 2 (1%) | 3 (1%) |
| White and Asian | 0 (0%) | 1 (1%) | 1 (0%) |
| Any other mixed / multiple ethnic background | 1 (1%) | 0 (0%) | 1 (0%) |
| Prefer not to say | 0 (0%) | 1 (1%) | 1 (0%) |
| Highest qualification | | | |
| Primary or less | 0 (0%) | 2 (1%) | 2 (1%) |
| Secondary | 24 (18%) | 39 (29%) | 63 (24%) |
| Higher (e.g. University, College) | 80 (60%) | 66 (49%) | 146 (54%) |
| Other qualifications | 27 (20%) | 25 (19%) | 52 (19%) |
| Employment status | | | |
| Employed full-time | 39 (29%) | 38 (28%) | 77 (29%) |

| | | | |
|---|-----------|-----------|-----------|
| Employed part-time | 52 (39%) | 50 (37%) | 102 (38%) |
| Self-employed | 17 (13%) | 13 (10%) | 30 (11%) |
| Currently unable to work due to poor health | 6 (4%) | 7 (5%) | 13 (5%) |
| Unemployed | 10 (7%) | 12 (9%) | 22 (8%) |
| Retired | 0 (0%) | 2 (1%) | 2 (1%) |
| Studying/student | 1 (1%) | 6 (4%) | 7 (3%) |
| Other | 9 (7%) | 5 (4%) | 14 (5%) |
| Relationship to child | | | |
| Parent | 129 (96%) | 131 (98%) | 260 (97%) |
| Other relative / legal guardian | 5 (4%) | 3 (2%) | 8 (3%) |
| Marital status | | | |
| Married/Civil partnership - cohabiting | 83 (62%) | 89 (66%) | 172 (64%) |
| Married/civil partnership - separated | 2 (1%) | 4 (3%) | 6 (2%) |
| Single - living in a couple | 21 (16%) | 11 (8%) | 32 (12%) |
| Single - not living in a couple | 14 (10%) | 11 (8%) | 25 (9%) |
| Divorced – cohabiting | 3 (2%) | 5 (4%) | 8 (3%) |
| Divorced – separated | 5 (4%) | 6 (4%) | 11 (4%) |
| Widowed | 1 (1%) | 1 (1%) | 2 (1%) |
| Prefer not to say | 0 (0%) | 1 (1%) | 1 (0%) |
| Partner's employment status | | | |
| Employed full-time | 68 (51%) | 70 (52%) | 138 (51%) |
| Employed part-time | 13 (10%) | 8 (6%) | 21 (8%) |
| Self-employed | 14 (10%) | 18 (13%) | 32 (12%) |
| Currently unable to work due to poor health | 5 (4%) | 1 (1%) | 6 (2%) |
| Unemployed | 4 (3%) | 3 (2%) | 7 (3%) |
| Retired | 2 (1%) | 2 (1%) | 4 (1%) |

| | | | |
|---|----------|----------|-----------|
| Other | 1 (1%) | 3 (2%) | 4 (1%) |
| Parent concerned about CYP psychological or emotional health | | | |
| Yes | 67 (50%) | 90 (67%) | 157 (59%) |
| No | 67 (50%) | 44 (33%) | 111 (41%) |
| Child sought help for psychological or emotional problem in last 6 months | | | |
| Yes | 62 (46%) | 65 (49%) | 127 (47%) |
| No | 72 (54%) | 69 (51%) | 141 (53%) |

Table 4: Baseline assessments and Patient Reported Outcome Measures (PROMs) for all randomised participants (n=268)

| Assessment/PROM | | CBT (n=134) | OST (n=134) | All (n=268) |
|----------------------------|-----------------------------------|----------------|----------------|----------------|
| Principal diagnosis (ADIS) | | | | |
| | Spiders | 5 (4%) | 6 (4%) | 11 (4%) |
| | Dogs | 30 (22%) | 26 (19%) | 56 (21%) |
| | Bees / Insects | 5 (4%) | 6 (4%) | 11 (4%) |
| | Other animals | 1 (1%) | 1 (1%) | 2 (1%) |
| | Heights | 1 (1%) | 2 (1%) | 3 (1%) |
| | Darkness | 2 (1%) | 0 (0%) | 2 (1%) |
| | Getting injections | 23 (17%) | 23 (17%) | 46 (17%) |
| | Having blood tests | 7 (5%) | 8 (6%) | 15 (6%) |
| | Seeing blood from a cut or scrape | 1 (1%) | 5 (4%) | 6 (2%) |
| | Lifts or small enclosed places | 0 (0%) | 2 (1%) | 2 (1%) |
| | Doctors / Dentists | 1 (1%) | 2 (1%) | 3 (1%) |
| | Vomiting | 41 (31%) | 36 (27%) | 77 (29%) |
| | Costumed characters | 5 (4%) | 1 (1%) | 6 (2%) |
| | Choking | 0 (0%) | 4 (3%) | 4 (1%) |
| | Food / Eating | 3 (2%) | 5 (4%) | 8 (3%) |
| | Other | 9 (7%) | 7 (5%) | 16 (6%) |
| ADIS CSR | | | | |
| | N (%) | 134 (100%) | 134 (100%) | 268 (100%) |
| | Mean (SD) | 7.5 (0.9) | 7.6 (0.9) | 7.5 (0.9) |
| | Median (IQR) | 8.0 (7.0, 8.0) | 8.0 (7.0, 8.0) | 8.0 (7.0, 8.0) |
| BAT steps | | | | |
| | N (%) | 134 (100%) | 133 (99%) | 267 (100%) |

| | | | | |
|-----------------------------|--------------|----------------------|----------------------|----------------------|
| | Mean (SD) | 4.1 (4.2) | 4.8 (4.2) | 4.4 (4.2) |
| | Median (IQR) | 2.0 (0.0, 9.0) | 4.0 (0.0, 10.0) | 3.0 (0.0, 10.0) |
| Change in SUDS | | | | |
| | N (%) | 121 (90%) | 120 (90%) | 241 (90%) |
| | Mean (SD) | -0.3 (2.3) | -0.1 (1.9) | -0.2 (2.1) |
| | Median (IQR) | 0.0 (-1.0, 0.0) | 0.0 (-1.0, 1.0) | 0.0 (-1.0, 1.0) |
| CAIS child | | | | |
| | N (%) | 130 (97%) | 133 (99%) | 263 (98%) |
| | Mean (SD) | 17.8 (13.6) | 18.1 (14.1) | 17.9 (13.8) |
| | Median (IQR) | 15.0 (8.0, 24.0) | 14.0 (7.0, 26.0) | 15.0 (7.0, 25.0) |
| CAIS parent | | | | |
| | N (%) | 127 (95%) | 129 (96%) | 256 (96%) |
| | Mean (SD) | 20.2 (16.3) | 21.9 (16.7) | 21.1 (16.5) |
| | Median (IQR) | 16.0 (6.0, 30.0) | 17.0 (8.0, 33.0) | 16.0 (8.0, 32.0) |
| CHU-9D | | | | |
| | N (%) | 133 (99%) | 132 (99%) | 265 (99%) |
| | Mean (SD) | 0.9 (0.1) | 0.8 (0.1) | 0.9 (0.1) |
| | Median (IQR) | 0.9 (0.8, 0.9) | 0.9 (0.8, 0.9) | 0.9 (0.8, 0.9) |
| EQ-5D-Y | | | | |
| | N (%) | 134 (100%) | 133 (99%) | 267 (100%) |
| | Mean (SD) | 0.741 (0.276) | 0.748 (0.258) | 0.745 (0.266) |
| | Median (IQR) | 0.812 (0.656, 1.000) | 0.812 (0.689, 0.883) | 0.812 (0.689, 1.000) |
| EQ-5D VAS | | | | |
| | N (%) | 134 (100%) | 133 (99%) | 267 (100%) |
| | Mean (SD) | 79.1 (19.6) | 77.7 (18.6) | 78.4 (19.1) |
| | Median (IQR) | 85.0 (70.0, 95.0) | 80.0 (65.0, 95.0) | 80.0 (70.0, 95.0) |
| Goal based outcome - goal 1 | | | | |
| | N (%) | 134 (100%) | 134 (100%) | 268 (100%) |

| | | | | |
|--|--------------|-----------------|-----------------|-----------------|
| | Mean (SD) | 1.7 (2.0) | 1.9 (1.9) | 1.8 (2.0) |
| | Median (IQR) | 1.0 (0.0, 3.0) | 1.5 (0.0, 3.0) | 1.0 (0.0, 3.0) |
| Goal based outcome - goal 2 | | | | |
| | N (%) | 127 (95%) | 129 (96%) | 256 (96%) |
| | Mean (SD) | 1.9 (2.1) | 2.1 (1.9) | 2.0 (2.0) |
| | Median (IQR) | 1.0 (0.0, 3.0) | 2.0 (0.0, 4.0) | 2.0 (0.0, 3.5) |
| Goal based outcome - goal 3 | | | | |
| | N (%) | 89 (66%) | 92 (69%) | 181 (68%) |
| | Mean (SD) | 2.0 (2.0) | 1.9 (1.8) | 1.9 (1.9) |
| | Median (IQR) | 1.0 (0.0, 3.0) | 2.0 (0.0, 3.0) | 2.0 (0.0, 3.0) |
| RCADS child social phobia score | | | | |
| | N (%) | 134 (100%) | 133 (99%) | 267 (100%) |
| | Mean (SD) | 8.8 (6.1) | 9.7 (6.5) | 9.2 (6.3) |
| | Median (IQR) | 9.0 (4.0, 12.0) | 9.0 (5.0, 14.0) | 9.0 (4.0, 13.0) |
| RCADS child panic disorder score | | | | |
| | N (%) | 134 (100%) | 133 (99%) | 267 (100%) |
| | Mean (SD) | 6.6 (5.4) | 7.8 (6.7) | 7.2 (6.1) |
| | Median (IQR) | 6.0 (2.0, 10.0) | 6.0 (3.0, 11.0) | 6.0 (2.0, 10.0) |
| RCADS child depression score | | | | |
| | N (%) | 134 (100%) | 133 (99%) | 267 (100%) |
| | Mean (SD) | 7.8 (5.6) | 8.3 (5.7) | 8.0 (5.7) |
| | Median (IQR) | 7.0 (3.0, 11.0) | 7.0 (4.0, 12.0) | 7.0 (4.0, 11.0) |
| RCADS child separation anxiety score | | | | |
| | N (%) | 134 (100%) | 133 (99%) | 267 (100%) |
| | Mean (SD) | 5.8 (4.7) | 6.0 (4.4) | 5.9 (4.5) |
| | Median (IQR) | 5.0 (2.0, 9.0) | 5.0 (2.0, 9.0) | 5.0 (2.0, 9.0) |

| | | | | |
|--|--------------|-------------------|-------------------|-------------------|
| <hr/> | | | | |
| RCADS child generalised anxiety score | | | | |
| | N (%) | 134 (100%) | 133 (99%) | 267 (100%) |
| | Mean (SD) | 6.4 (4.2) | 7.1 (4.4) | 6.7 (4.3) |
| | Median (IQR) | 6.0 (3.0, 9.0) | 7.0 (4.0, 10.0) | 6.0 (3.0, 9.0) |
| RCADS child obsessive compulsive score | | | | |
| | N (%) | 134 (100%) | 133 (99%) | 267 (100%) |
| | Mean (SD) | 4.4 (3.8) | 5.0 (4.2) | 4.7 (4.0) |
| | Median (IQR) | 4.0 (1.0, 7.0) | 4.0 (2.0, 7.0) | 4.0 (2.0, 7.0) |
| RCADS child total anxiety score | | | | |
| | N (%) | 134 (100%) | 133 (99%) | 267 (100%) |
| | Mean (SD) | 32.0 (19.1) | 35.6 (21.8) | 33.8 (20.5) |
| | Median (IQR) | 28.5 (16.0, 47.0) | 31.0 (19.0, 52.0) | 30.0 (17.0, 49.0) |
| RCADS parent social phobia score | | | | |
| | N (%) | 134 (100%) | 134 (100%) | 268 (100%) |
| | Mean (SD) | 9.6 (5.8) | 11.4 (5.8) | 10.5 (5.9) |
| | Median (IQR) | 9.0 (5.0, 13.0) | 11.0 (7.0, 15.0) | 10.0 (6.0, 14.0) |
| RCADS parent panic disorder score | | | | |
| | N (%) | 133 (99%) | 134 (100%) | 267 (100%) |
| | Mean (SD) | 6.1 (5.0) | 7.7 (5.9) | 6.9 (5.5) |
| | Median (IQR) | 5.0 (2.0, 8.0) | 6.0 (3.0, 10.0) | 5.0 (3.0, 10.0) |
| RCADS parent depression score | | | | |
| | N (%) | 134 (100%) | 134 (100%) | 268 (100%) |
| | Mean (SD) | 8.0 (5.5) | 8.8 (5.8) | 8.4 (5.6) |
| | Median (IQR) | 7.0 (4.0, 11.0) | 8.0 (5.0, 13.0) | 7.5 (4.0, 12.0) |
| <hr/> | | | | |

| | | | | |
|---|--------------|-------------------|-------------------|-------------------|
| <hr/> | | | | |
| RCADS parent separation anxiety score | | | | |
| | N (%) | 134 (100%) | 134 (100%) | 268 (100%) |
| | Mean (SD) | 6.4 (4.9) | 7.3 (5.2) | 6.9 (5.1) |
| | Median (IQR) | 5.0 (2.0, 10.0) | 6.0 (3.0, 12.0) | 6.0 (3.0, 10.0) |
| RCADS parent generalised anxiety score | | | | |
| | N (%) | 133 (99%) | 134 (100%) | 267 (100%) |
| | Mean (SD) | 6.8 (4.3) | 7.7 (4.5) | 7.2 (4.4) |
| | Median (IQR) | 6.0 (3.0, 10.0) | 7.0 (4.0, 11.0) | 7.0 (4.0, 10.0) |
| RCADS parent obsessive compulsive score | | | | |
| | N (%) | 132 (99%) | 134 (100%) | 266 (99%) |
| | Mean (SD) | 3.6 (3.8) | 3.9 (3.1) | 3.7 (3.5) |
| | Median (IQR) | 2.0 (0.5, 6.0) | 3.0 (2.0, 6.0) | 3.0 (1.0, 6.0) |
| RCADS parent total anxiety score | | | | |
| | N (%) | 132 (99%) | 134 (100%) | 266 (99%) |
| | Mean (SD) | 32.4 (19.1) | 37.9 (20.3) | 35.2 (19.9) |
| | Median (IQR) | 29.0 (17.5, 45.0) | 35.0 (23.0, 51.0) | 32.0 (19.0, 50.0) |

ADIS CSR is measured on a 0-8 scale, higher score means the phobia is more disturbing

BAT steps is measured on 0 (no steps completed) to 10 (all steps completed)

SUDS is measured on a 0-8 scale, higher scores mean more anxiety/fear

CAIS measures impact of anxiety symptoms on psychological function on the 0-81 scale, higher scores represent greater impact

CHU-9D is measured on a 0.33-1 scale higher scores mean greater health related quality of life

EQ-5D-Y is measured on a scale of -0.594 to one (full health)

Goal based outcome is measured from 0 (goal not at all met) to 10 (goal reached)

RCADS measures anxiety and depression symptoms, higher scores mean more anxiety/depression symptoms RCADS total anxiety is measured on 0-111 scale

Table 5: Comparison of mean 6-month assessment and quality of life outcomes by treatment group (n=197)

| Outcome | Treatment group | | | |
|----------------------------------|-----------------|-------------|-----|-------------|
| | CBT | | OST | |
| | N | Mean (SD) | n | Mean (SD) |
| BAT steps | 76 | 7.1 (3.9) | 73 | 7.4 (3.6) |
| SUDS before | 74 | 3.8 (3.0) | 72 | 3.7 (2.7) |
| SUDS change | 74 | -0.8 (2.1) | 70 | -0.5 (2.0) |
| ADIS CSR | 97 | 5.1 (2.6) | 100 | 4.9 (2.7) |
| CAIS child | 93 | 12.0 (11.2) | 96 | 14.5 (14.3) |
| CAIS parent | 92 | 14.4 (14.5) | 92 | 15.0 (15.8) |
| CHU-9D | 95 | 0.89 (0.10) | 96 | 0.87 (0.10) |
| EQ-5D-Y | 96 | 0.81 (0.24) | 98 | 0.82 (0.23) |
| EQ-5D-Y VAS | 96 | 84.2 (16.3) | 98 | 82.6 (17.1) |
| Goal based outcome - goal 1 | 94 | 6.4 (3.2) | 95 | 5.8 (3.4) |
| Goal based outcome - goal 2 | 91 | 6.2 (3.5) | 94 | 5.9 (3.2) |
| Goal based outcome - goal 3 | 66 | 6.5 (3.7) | 69 | 6.1 (3.2) |
| RCADS child social phobia score | 96 | 7.3 (5.8) | 97 | 8.6 (6.0) |
| RCADS child panic disorder score | 96 | 5.0 (4.7) | 97 | 5.6 (5.2) |

BAT is measured on a 0 (no steps) to 10 (all steps completed) scale, a positive difference means the CBT group completed more steps at six-months than the OST group

SUDS is measured on a 0-8 scale, higher scores mean more anxiety/fear

ADIS CSR is measured on a 0-8 scale higher score means phobia is more disturbing

CAIS measures impact of anxiety symptoms on psychological function on the 0-81 scale, higher scores represent greater impact

CHU-9D is measured on a 0.33-1 scale higher scores mean greater health related quality of life

EQ-5D-Y is measured on a scale of -0.594 to one (full health)

Goal based outcome is measured from 0 (goal not at all met) to 10 (goal reached)

RCADS measures anxiety and depression symptoms, higher scores mean more anxiety/depression symptoms RCADS total anxiety is measured on 0-111 scale

Table 6: Adverse events and serious adverse events by treatment group (n=268)

| Adverse Events | CBT | OST | All |
|---|----------------|----------------|----------------|
| | (n=134) | (n=134) | (n=268) |
| Number (%) of participants who experienced ≥ 1 AE | 4(3.0%) | 5(3.7%) | 9(3.4%) |
| Number of all AEs (including repeated events) | 5 | 5 | 10 |
| Number (%) of participants who experienced ≥ 1 SAE | 2(1.5%) | 1(0.7%) | 3(1.1%) |
| Number of all SAEs (including repeated events) | 3 | 1 | 4 |
| Occurred | | | |
| Before randomisation | 1 (20.0%) | 0 (0.0%) | 1 (10.0%) |
| After randomisation | 4 (80.0%) | 5 (100.0%) | 9 (90.0%) |
| AE category | | | |
| fainting/light headedness | 1 (20.0%) | 1 (20.0%) | 2 (20.0%) |
| Needle injury/scratch | 1 (20.0%) | 1 (20.0%) | 2 (20.0%) |
| Inpatient hospitalisation | 3 (60.0%) | 1 (20.0%) | 4 (40.0%) |
| Other | 0 (0.0%) | 2 (40.0%) | 2 (20.0%) |
| Seriousness | | | |
| Inpatient hospitalisation | 3 (60.0%) | 1 (20.0%) | 4 (40.0%) |
| Intensity | | | |
| Moderate | 1 (20.0%) | 1 (20.0%) | 2 (20.0%) |
| Severe | 2 (40.0%) | 0 (0.0%) | 2 (20.0%) |
| Relationship to phobia treatment | | | |
| Unrelated | 3 (60.0%) | 1 (20.0%) | 4 (40.0%) |