



Examining the benefit of graduated
compression stockings in the Prevention
of vEnous Thromboembolism in low-risk
Surgical patients: a multicentre cluster
randomised controlled trial (PETS Trial)

V7.0, 07.05.2024

SPONSOR: Imperial College London

FUNDER: National Institute for Health Research (NIHR) – Health Technology Assessment (HTA)

STUDY COORDINATION CENTRE: Imperial College London

IRAS Project ID: 312752

REC reference: 22/LO/0390

Revision History

Protocol Version	Date	Amendments
V7.0	07.05.2024	<ul style="list-style-type: none"> The addition of a mixed-methods sub-study exploring the barriers, facilitators and attitudes towards VTE Clinical Practice Guidelines
V6.0	11/04/2024	<ul style="list-style-type: none"> Update to confirm that non-NHS sites can be included
V5.0	23/02/2024	<ul style="list-style-type: none"> An update to the eligibility criteria to confirm that patients who require extended thromboprophylaxis beyond discharge are not excluded.
V4.0	28/09/2023	<ul style="list-style-type: none"> Confirmation that patients who undergo varicose vein related procedures should not be enrolled in the study The option to collect follow-up data via post
V3.0	29/11/2022	<ul style="list-style-type: none"> Addition of “pharmacological prophylaxis” into the exclusion criteria Clarity on the eligibility of clusters, method of randomisation for clusters and blinding to treatment allocation Updated study flow chart to include the eligibility and randomisation of clusters Additional information on the follow-ups at 7 and 90 days Clarity around the sample size calculation and statistical analyses section Clarification surrounding the withdrawal of participants from the study

		<ul style="list-style-type: none"> • Additional information regarding the internal pilot • Addition of data management section • Correction of typographical errors
V 2.0	18/07/2022	<ul style="list-style-type: none"> • Addition of the EQ-5D pre-procedure • Addition of the REC reference/name of the REC
V 1.0	29/03/2022	Protocol submitted the REC

Protocol authorised by:

Name & Role

Professor Alun Davies
Chief investigator

Date

28/09/2023

Signature

Professor John Norrie
Lead Statistician and Co-Applicant

28/09/2023

Ms Becky Ward
Sponsor Representative

28/09/2023

Study Management Group

Chief Investigator:

Professor Alun Davies
Imperial College London, Academic Section of Vascular Surgery

Room 3, 4th Floor East Wing, Charing Cross Hospital
Fulham Palace Road, London, W6 8RF
Email: a.h.davies@imperial.ac.uk

Senior Statistician:

Professor John Norrie
Chair of Medical Statistics and Trial Methodology
Director of Edinburgh Clinical Trials Unit
Usher Institute- University of Edinburgh
9 Little France Road
Nine Edinburgh BioQuarter
Edinburgh, EH16 4UX
Email: j.norrie@ed.ac.uk

Co-investigators:

Mr Joseph Shalhoub, Consultant Vascular Surgeon/Honorary Clinical Senior Lecturer

Professor John Norrie, Director of Edinburgh Clinical Trials Unit

Mr Matthew Machin, Academic Clinical Research Fellow

Miss Sasha Smith, Clinical Trial Manager

Professor Beverley Hunt, Professor of Thrombosis and Haemostasis

Dr Tamara Everington, Consultant in Haemostasis and Thrombosis

Dr David Epstein, Associate Professor

Mr Manjit Gohel, Consultant Vascular Surgeon

Professor David Warwick, Consultant Orthopaedic Surgeon

Dr Christopher Baker, Consultant Cardiologist

Mr Zaed Hamady, Consultant General Surgeon

Ms Layla Bolton, Clinical Nurse Specialist

Ms Annya Stephens-Boal, PPI representative

Ms Beverley Gray, PPI representative

Study Coordination Centre

For general queries, supply of study documentation, and collection of data, please contact:

Study Coordinator: Mrs Sarrah Peerbux

Address:

Imperial College London
Section of Vascular Surgery
Room 3E, 4th Floor East Wing
Charing Cross Hospital
Fulham Palace Road
London W6 8RF

Tel: +44 (0)203 311 7371

E-mail: s.peerbux@imperial.ac.uk/petstrial@imperial.ac.uk

Clinical Queries

Clinical queries should be directed to the Trial Manager at petstrial@imperial.ac.uk who will direct the query to the appropriate person

Sponsor

Imperial College London is the main research Sponsor for this study. For further information regarding the sponsorship conditions, please contact the Head of Regulatory Compliance at:

Research Governance and Integrity Team
Imperial College London and Imperial College Healthcare NHS Trust
Room 215, Level 2, Medical School Building
Norfolk Place
London, W2 1PG

Tel: 0207 594 1862

[Imperial College - Research Governance and Integrity Team \(RGIT\) Website](#)

Funder

This study is funded by the National Institute for Health Research (NIHR) Health Technology Assessment programme (NIHR133776). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

This protocol describes the PETS study and provides information about procedures for entering participants. Every care was taken in its drafting, but corrections or amendments may be necessary. These will be circulated to investigators in the study. Problems relating to this study should be referred, in the first instance, to the Chief Investigator.

This study will adhere to the principles outlined in the UK Policy Frame Work for Health and Social Care Research. It will be conducted in compliance with the protocol, the Data Protection Act and other regulatory requirements as appropriate.

Table of Contents**Page No**

1. INTRODUCTION	- 8 -
1.1. BACKGROUND	- 9 -
1.2. RATIONALE FOR CURRENT STUDY	- 10 -
2. STUDY OBJECTIVES	- 10 -

2.1 PRIMARY OUTCOME	- 11 -
2.2 SECONDARY OUTCOMES	- 11 -
3. STUDY DESIGN	- 11 -
3.1. STUDY OUTCOME MEASURES	Error! Bookmark not defined.
4. PARTICIPANT ENTRY	- 15 -
4.1. INCLUSION CRITERIA	- 15 -
4.2. EXCLUSION CRITERIA	- 15 -
4.3. WITHDRAWAL CRITERIA	- 16 -
5. ADVERSE EVENTS	Error! Bookmark not defined.
5.1. DEFINITIONS	Error! Bookmark not defined.
5.2. REPORTING PROCEDURES	Error! Bookmark not defined.
6. ASSESSMENT AND FOLLOW-UP	- 16 -
6.1 ELIGIBILITY AND ASSESSMENT	- 16 -
6.2 INFORMED CONSENT	- 17 -
6.3 FOLLOW-UP	- 17 -
7. STATISTICS AND DATA ANALYSIS	- 20 -
7.1 Sample size calculation	- 20 -
7.2 Statistical analyses	- 21 -
7.3 Internal pilot of feasibility and progression criteria	Error! Bookmark not defined.
7.4 Measurement of cost-effectiveness	- 22 -
8. REGULATORY ISSUES	- 24 -
8.1. ETHICS APPROVAL	- 24 -
8.2. CONSENT	- 25 -
8.3. CONFIDENTIALITY	- 25 -
8.4. INDEMNITY	- 25 -
8.5. SPONSOR	- 25 -
8.6. FUNDING	- 25 -
8.7. AUDITS	- 25 -
9. STUDY MANAGEMENT	- 26 -
10. PUBLICATION POLICY	- 26 -
11. REFERENCES	- 26 -

GLOSSARY OF ABBREVIATIONS

CHEERS	Consolidated Health Economic Evaluation Reporting Standards
CI	Confidence Interval
CTU	Clinical trials unit
DHRA tool	Department of health risk assessment tool
DVT	Deep venous thrombosis
EQ-5D	Euro-QoL 5D instrument for measuring generic health status
GAPS	Graduated compression stockings as adjuvant to pharmacothromboprophylaxis in elective surgical patients' study
GCS	Graduated compression stocking
HAT	Hospital-acquired thrombosis
ICC	Intraclass correlation
iDMC	Independent data monitoring committee

LMWH	Low-molecular weight heparin
NHS	National Health Service
NICE	National Institute for Health and Care Excellence
NNT	Number needed to treat
PE	Pulmonary embolism
PPI	Patient and public involvement
PTS	Post-thrombotic syndrome
QALY	Quality-adjusted life year
QoL	Quality of Life
RCT	Randomised Controlled Trial
RGIT	Research Governance and Integrity Team
TEDS	Thromboembolic deterrent stockings
SAP	Statistical Analysis Plan
SMS	Short Messaging Service
SOP	Standard operating procedure
TSC	Trial Steering Committee
VTE	Venous thromboembolism
WHO	World Health Organisation

KEYWORDS

Venous Thromboembolism, graduated compression stockings, deep venous thrombosis, pulmonary embolism, cluster design

STUDY SUMMARY

TITLE Examining the benefit of graduated compression stockings in the Prevention of vEnous Thromboembolism in low-risk Surgical patients: a multicentre cluster randomised controlled trial (PETS Trial)

TRIAL DESIGN Assessor blind, multicentre, parallel-group cluster, randomised controlled trial

AIMS To evaluate the potential benefit of Graduated Compression Stockings (GCS) in the prevention of Venous Thromboembolism (VTE) in patients undergoing short-stay surgical procedures, assessed as being at low-risk for VTE.

OUTCOME MEASURES Primary Outcome:
The occurrence of symptomatic VTE within 90 days from date of surgery .

Secondary outcomes:

- Mortality at 7 and 90 days post-surgery

- Incremental Cost-Effectiveness Ratio (ICER) up to 2 years
- Quality of life – EQ-5D at 7 and 90 days post-surgery
- Adverse events with GCS (assessed at 7-days post-surgery, in those enrolled in the intervention clusters only).

POPULATION Patients undergoing short-stay surgical procedures, assessed as being at low-risk of developing VTE by Department of Health Risk Assessment (DHRA) tool (equivalent to scoring 0, i.e. absence of all assessed thrombosis risk factors).

SAMPLE SIZE We plan to recruit a total of 21,472 participants from 25 intervention and 25 control sites.

PATIENT ELIGIBILITY Inclusion criteria

- Adults (18-59 years of age) scheduled to undergo a surgical procedure with a hospital stay <48 hours
- Individuals assessed as being at low-risk of developing VTE as per the DHRA Tool (i.e. no assessed thrombosis risk factors / scoring 0)

Exclusion criteria

- Individuals with a contraindication to GCS
- Individuals assessed as being at moderate or high-risk of VTE as per the DHRA tool
- Individuals requiring therapeutic anticoagulation
- Individuals requiring pharmacological prophylaxis
- Individuals with thrombophilia/ thrombogenic disorder
- Individuals with a previous history of VTE
- Individuals requiring intermittent pneumatic compression therapy beyond theatre and recovery
- Female patients of childbearing age who have a positive pregnancy test
- Individuals with lower limb immobilisation
- Inability to provide informed consent

CLUSTER ELIGIBILITY In order to be deemed suitable to participate, Trusts must conduct either day case or short stay surgical procedures.

DURATION 45-months

1. INTRODUCTION

1.1. BACKGROUND

Hospital-acquired thrombosis (HAT) is defined as any venous thromboembolism (VTE) related event within 90 days of hospital admission (1), and is a term that encompasses both deep vein thrombosis (DVT) and pulmonary embolism (PE). HAT accounts for significant morbidity and mortality with 57.1 VTE-related deaths per 100,000 hospital admissions reported in the year 2018-2019 within the NHS (1). The probability of untreated moderate-risk surgical inpatients developing HAT was previously estimated to be as high as 15%, this reducing to 4.1% with pharmacological and mechanical prophylaxis (2).

The UK annual VTE-related mortality is estimated to be 32,000 fatalities, with associated costs of £640 million per annum (3). After suffering from a DVT, around half of patients develop post-thrombotic syndrome (PTS) (4), characterised by leg pain, oedema, skin changes and leg ulceration (5). Surgery is an established risk factor for VTE (6). It is estimated that the untreated risk of VTE for patients undergoing most general surgical, urological or open gynaecological procedures is 10-40% (7). Furthermore, for those undergoing hip or knee arthroplasty this risk increases to 40-80% (7). Specific surgical factors such as abdominal and pelvic procedures, operations for cancer, and procedures with a greater duration are associated with a greater risk of VTE (8,9). Conversely, operations with a short anaesthetic and procedural time, that can be performed within a <48-hour hospital stay, and that permit early ambulation are associated with a low risk of VTE. An example of this is groin hernia repair - in a large consecutive cohort of 5649 patients, half of whom received low-molecular weight heparin (LMWH), the rate of DVT was just 0.12% (10).

Mechanical thromboprophylaxis is often utilised in the form of graduated compression stockings (GCS), also known by the brand name ThromboEmbolic Deterrent (TED) (7,11,12). GCS apply a level of graduated pressure at rest, i.e. not whilst ambulant - they are designed for immobile patients at-risk of VTE. However, it has recently been shown that GCS provide no added benefit in the reduction of VTE for moderate-high VTE risk surgical inpatients receiving LMWH (13). Additionally, the CLOTS 1 trial, which randomised acute stroke patients to either LMWH alone or the combination of GCS and LMWH, reported no difference in the rate of DVT (14). The findings of these studies have cast doubt on the use of GCS in prevention of VTE. Importantly, the patients in the aforementioned trials were at a higher risk from this distinct low VTE risk group to which this application refers.

Limited evidence is available on the rate of hospital-acquired VTE in low-risk surgical patients. NICE guidelines for the prevention of VTE published in 2007 previously recommended that all surgical patients should receive GCS to reduce the risk of VTE, irrespective of the absence of thrombosis risk factors (15). The subsequent updated NICE guidelines did not include this blanket recommendation, but instead provided specific recommendations for differing procedure types. The most contemporary NICE guidelines, published in 2018, recommend that all patients undergoing abdominal, thoracic, spinal, bariatric, head and neck, and elective joint surgery should be treated with GCS and to consider treatment with GCS for all those undergoing cardiac, vascular and ear, nose and throat surgery (16). The interpretation of these recommendations has meant that patients undergoing short-stay procedures, who are able to ambulate early and have no other thrombosis risk factors, are still treated with GCS. An example of this interpretation is the provision of GCS for inguinal hernia repair procedures - an operation that fits into the recommendation of provision of GCS for an abdominal procedure, however, patients ambulate immediately and are discharged from hospital on the same day. This blanket provision of GCS is evidenced by our detailed feasibility analysis that demonstrated many NHS Trusts within the UK offer GCS to all surgical patients, even those at low-risk of VTE. Hence, it is now common practice for patients undergoing day case

procedures, even those that have no other assessed risk factors, to be prescribed GCS in the absence of contraindications.

1.2. RATIONALE FOR CURRENT STUDY

The evidence to support GCS for short-stay ambulant patients is poor. A systematic review (17) examining the use of GCS in comparison to no prophylaxis in low VTE risk short stay surgical patients confirmed that there have been no RCTs to support this practice. Furthermore, a meta-analysis in the 2018 Cochrane Review for patients at moderate and high-risk of VTE revealed a significantly lower odds ratio of DVT in those treated with GCS in comparison to those that were not (18). However, this analysis is not relevant to the current research question which comprises of the use of GCS alone in low VTE risk surgical patients, i.e. not receiving low molecular weight heparin (LMWH) and undergoing short stay procedures. None of the included 18 surgical RCTs identified in the Cochrane Review consisted of low risk procedures, i.e. ambulatory day case procedures. The case mix consisted of the following types of procedures: neurosurgery, orthopaedic trauma, major lower limb arthroplasty and major abdominal surgery.

Moreover, the NIHR HTA-funded RCT (GAPS trial) assessing the use of GCS in addition to LMWH for the prevention of VTE in moderate and high-risk, i.e. not low risk, elective surgical patients demonstrating that LMWH alone was non-inferior to dual thromboprophylaxis with LMWH and GCS, has further drawn the role of GCS in the prevention of VTE in surgery into question (13). This is on the background of other RCTs, such as the CLOTS 1 trial which randomised acute stroke patients to either LMWH alone or the combination of GCS and LMWH and reported no difference in the rate of DVT (14). In the GAPS trial there was no adjuvant benefit of GCS in patients receiving LMWH, while the clinical community are adopting these findings into clinical practice, they cannot safely be extrapolated to this large cohort of low VTE risk patients.

Due to this lack of evidence to support the use of GCS in the prevention of VTE for low VTE risk surgical patients, an adequately powered trial is required to provide level-1A evidence in relation to this practice.

If this practice is demonstrated to be without therapeutic benefit, then foregoing this will advance the environmental agenda of the NHS as well as save money. These valuable resources can also be redistributed elsewhere, and the potential adverse effects of GCS avoided.

To summarise, this trial is necessary because:

- a) There is no evidence to support clinicians in the prescription of GCS to prevent VTE in low VTE risk surgical patients
- b) There is no information to counsel patients when informing them regarding the use of GCS
- c) There is potential to save a significant financial resource
- d) Evidence is required to guide future clinical practice policy

2. STUDY OBJECTIVES

The aim of this study is to evaluate the potential benefit of GCS in the prevention of VTE in patients undergoing short-stay surgical procedures, assessed as being at low-risk for VTE.

2.1 PRIMARY OUTCOME

The primary outcome is the occurrence of symptomatic VTE (DVT or PE) within 90 days, of date of surgery, for surgical patients undergoing short-stay procedures assessed as being at low-risk of VTE for those treated with GCS in comparison to those not given thromboprophylaxis.

2.2 SECONDARY OUTCOMES

- Mortality at 7 and 90 days post-surgery
- Incremental Cost-Effectiveness Ratio (ICER) at 90 days post-surgery and, if relevant up to 2 years
- Quality of life – EQ-5D (19) at 7 and 90 days post-surgery
- Adverse events with GCS (assessed at 7-days post-surgery, in those enrolled in the intervention cluster only).


3. STUDY DESIGN

3.1 TYPE OF STUDY

This is an assessor-blind multicentre, parallel group sequential cluster randomised controlled trial with internal pilot and one interim analysis at 50% of recruitment with mature data on the primary outcome. A total of 50 sites (21,472 participants) will be randomised (1:1) to either GCS or no GCS. Inclusion criteria stipulates these participants are adult, undergoing short stay procedures, and assessed low-risk for VTE as per the Department of Health risk assessment tool (20).

The duration of the study will be 45 months including 6-months set up, 30 months of recruitment, 3 months follow-up and 6 months for analysis and dissemination. **Figure 1** is a flow diagram summarising the study design.

Figure 1: Study flow diagram



*A formal interim analysis with the possibility of stopping early for futility (no prospect of a clinically meaningful treatment effect), at the point of 50% mature primary outcome data.

3.2 BLINDING

The primary outcome will be assessed blindly by blinded assessors at Imperial College London during the 7 and 90 day follow ups.

3.3 FOLLOW UP

Participants will be contacted at 7 and 90 days to obtain follow up data.

Day 7 post procedure (data collected via telephone or online survey [link to survey sent via email or SMS])

- VTE outcome (participants will be asked to report on whether or not they have been diagnosed with a DVT or PE within the past 7-days)
- EQ-5D (assessing generic quality of life)
- Adverse events associated with GCS will be captured (for those enrolled in the Intervention cluster site)

Day 90 post procedure (data collected via telephone or online survey [link to survey sent via email or SMS])

- VTE outcome (participants will be asked to report on whether or not they have been diagnosed with a DVT or PE within the past 90-days)
- EQ-5D (assessing generic quality of life) • Resource use data will be collected (capturing information on any consultations with a primary/secondary care healthcare professional over the past 90-days)

Mortality within the 90-day follow-up period will also be captured. Follow-up data will be assessed blindly.

3.4 Internal Pilot

There will be an internal pilot to assess feasibility of recruitment over the first 6 months of recruitment to the trial, in which we will start recruiting the (minimum) 50 sites at 4 sites per month. The participant consent rate to meet the target sample size is 20 participants per centre month (accounting for the staggered site set-up). This equates to a target of 1440 participants at the end of the 6-month internal pilot.

For the internal pilot we will use a stop-go criteria based on a Green-Amber-Red statistical approach, as follows:

Figure 2: Progression criteria

Progression criteria	Red	Amber	Green
% of total consented participants	<5%	5 – 6.9%	≥7%
Participant consent rate per site per month	<16	16 - 19	≥20
Number of sites opened	<20	20 -23	≥24
Total number of participants consented	<1000	1000 – 1439	≥1440

4. PARTICIPANT ENTRY

4.1. INCLUSION CRITERIA

- Adults (18-59 years of age) scheduled to undergo a surgical procedure with a hospital stay <48 hours.
- Individuals assessed as being at low-risk of developing VTE as per the DHRA tool (20) i.e. no assessed thrombosis risk factors / scoring 0.

Examples of procedures from which patients are at low-risk of VTE include (but not limited to): open general surgery (e.g. inguinal hernia repair), laparoscopic general surgery (e.g. laparoscopic cholecystectomy), ear, nose and throat (e.g. myringotomy), urology (e.g. cystoscopy), gynaecology (e.g. diathermy for endometriosis), and orthopaedics (e.g. joint arthroscopy).

4.2. EXCLUSION CRITERIA

- Individuals with a contraindication to GCS
- Individuals assessed as being at moderate or high-risk of VTE as per the DHRA tool
- Individuals requiring therapeutic anticoagulation
- Individuals with thrombophilia/ thrombogenic disorder
- Individuals with a previous history of VTE
- Individuals requiring intermittent pneumatic compression therapy beyond theatre and recovery
- Individuals requiring pharmacological prophylaxis
- Female patients of childbearing age who have a positive pregnancy test
- Individuals with lower limb immobilisation
- Inability to provide informed consent

Procedures undertaken to treat varicose veins (e.g. endovenous radiofrequency ablation of the great saphenous vein or avulsion of varicose veins) are ineligible. This is due to the fact

that such varicose vein procedures pose a higher risk of VTE than standard short-stay surgical procedures. Patients undergoing these procedures are being recruited to the [THRIVE](https://fundingawards.nihr.ac.uk/award/NIHR152877) trial (<https://fundingawards.nihr.ac.uk/award/NIHR152877>) .

4.3. WITHDRAWAL CRITERIA

Participants may discontinue study intervention for the following reasons:

- At the request of the participant.
- Adverse event/ Serious Adverse Event
- If the investigator considers that a participant's health will be compromised due to adverse events or concomitant illness that develop after entering the study.

The reason for participant withdrawal should be documented in the Study Completion CRF in REDCap. Participants that are withdrawn or lost to follow-up do not need to be replaced.

5. ASSESSMENT AND FOLLOW-UP

5.1 ELIGIBILITY AND ASSESSMENT

This is a “cluster study” i.e randomisation will be conducted at recruitment centre level rather than on an individual participant level. Recruiting centres will be randomised/allocated to either the intervention arm (where all participants who undergo short-stay surgery will receive GCS to wear during their short-stay procedure) or the control arm (where no GCS will be provided to those undergoing short-stay surgery) once confirmation of capacity and capability has been issued by the local R&D team. This treatment allocation will apply to **all** patients (cohorts in whom equipoise has been agreed) who undergo short-stay surgery within the Trust, i.e. the Trust will adopt this allocation as their standard of care.

Patients are only required to consent to be contacted for follow-up (7 and 90-days post procedure).

This trial will take place in at least 50 NHS Trusts and non-NHS sites offering day case and short stay surgical services. Patients from a variety of surgical specialties will be included in this pragmatic trial. Adults who are scheduled to undergo short-stay surgery will be pre-screened by a member of the direct care team and invited to speak to a member of the research team. Permission from the patient will be granted by the direct care team before any information is passed or approach made by the research team.

The reasons for non-inclusion will be logged anonymously along with a minimum data set of age, gender and reason for exclusion. The anonymised screening logs will be transferred to the Trial Coordinating Centre for the purposes of monitoring recruitment.

5.1.1. RANDOMISATION

Randomisation schedules were generated using permuted blocks and stored by ECTU. As sites are set up, the site is allocated to either intervention or control as per the random allocation sequence and the site administrator is informed of the allocation.

5.1.2. INTERVENTION

Centres randomised to the intervention arm, which is the current standard of care, will consist of participants receiving GCS. Clinical staff (e.g. theatre support workers) will issue stockings to all patients who are scheduled to undergo short-stay surgery. Participants will be instructed to put on their stockings just before undergoing the surgical procedure and to wear for the duration advised by the clinician.

5.1.3. CONTROL

In those centres randomised to the control arm, participants will not receive GCS.

5.2 INFORMED CONSENT

Participants will be consented to be contacted for follow-up (7 and 90-days post procedure).

Eligible patients will be provided with further information about the study. Formal informed consent to be contacted for follow-up (7 and 90-days) will be obtained prior to the day of procedure (e.g. at the preassessment visit) or at any point before the 7-day follow-up is due (including on the day of the surgical procedure). Informed consent can be collected in written, verbal or electronic form.

Prior to the surgical procedure, all participants will be provided with a leaflet which explains the signs and symptoms of developing a blood clot. Although VTE outcome will be assessed at 7 and 90-days post-procedure, participants will be advised to visit the emergency department if they suspect they have developed a blood clot (i.e. and not to wait for the study team to make contact).

Prior to the procedure, the following information will be collected from the patient and medical records:

- Baseline demographic information
- Name of surgical procedure
- Previous medical history and current medication
- EQ-5D (issued immediately after the participant provides consent either in-person, over the telephone or electronically).

5.3 FOLLOW-UP

Participants will be followed-up at 7 and 90-days post operation. This follow-up can be conducted via telephone or online questionnaire (the link to the questionnaire will be sent via email or SMS). Data can also be collected via post.

Data collection is summarised below:

Day 7 post procedure (assessed blindly)

- VTE outcome
- Mortality
- Quality of life questions – EQ-5D
- Adverse events with GCS (for participants enrolled in the intervention cluster only)

Day 90 post procedure (assessed blindly)

- VTE outcome

- Mortality
- Quality of life questions – EQ-5D
- Resource use questionnaire

The collection of follow-up data will be performed by blinded assessors based at the coordinating centre (Imperial College London).

5.4 END OF STUDY

The end of the study is defined as the last patient last visit.

5.5 MIXED METHODS SUB-STUDY

A mixed-methods sub-study (comprising qualitative and quantitative data collection) will be conducted to complement the main PETS trial and will recruit Healthcare Professionals (HCPs) responsible for conducting the VTE risk assessment. The mixed-methods sub-study will explore the perceived barriers and facilitators to clinician adherence to VTE clinical practice guidelines.

The quantitative study will involve HCPs completing the Clinical Guideline Determinants Questionnaire (CGDQ). The qualitative study will involve one-to-one semi-structured interviews with HCPs involved in conducting VTE risk assessment.

Inclusion criteria

- Healthcare professional (any background and grade)
- involved in conducting VTE risk assessments in surgical patients or responsible for prescribing thromboprophylaxis to surgical patients.

Exclusion criteria

- Not involved in conducting VTE risk assessments in surgical patients or not responsible for prescribing thromboprophylaxis to surgical patients,

5.5.1 Quantitative study

The CGDQ is a validated questionnaire comprising four sections to assess; 1) clinician demographic information (including two determinants [attitudes towards and experience with the guidelines]); 2) closed-ended items reflecting clinician- and guideline-specific determinants (ranked on a 7-point Likert scale); 3) four open-ended items reflecting barriers and facilitators perceived to be the most important; and 4) three items on learning style (preferred sources of guideline information).

The adapted CGDQ will be used to assess the determinants of clinician adherence to VTE Clinical Practice Guidelines (including the perceived barriers and facilitators to adhering to these guidelines).

HCPs will be contacted through a range of recruitment methods, including contacting HCPs involved in recruiting to PETS and utilising contacts from professional member networks.

HCPs responsible for conducting the VTE risk assessment will be invited to complete the CGDQ via the Qualtrics online survey software. Recruitment is targeted at a minimum of n=100 HCPs as a convenience sample to complete the CGDQ.

The link to the survey will be shared via social media posts or in the ethically approved communication with potential participants. A link to the PIS will be provided on the landing page of the survey. The web-based Qualtrics Survey Platform (<https://www.qualtrics.com>) enables opt-in consent by completion.

The survey will be available for completion for at least four months. Potential participants can complete the survey at any point whilst it is live.

Participants will provide the data anonymously. However, they will have the opportunity to share personally identifiable information if they choose (e.g. if they would like to find out more information about the qualitative study).

5.5.2 Quantitative data analysis

Responses to the CGDQ will provide quantitative data from both nominal and ordinal scales as well as qualitative data from open ended questions. Descriptive statistics will be presented where appropriate. Responses to the Likert scale will be collapsed where appropriate, such that a three-point scale will be established to reflect 'disagree', 'agree' and 'neutral'. Items coded as disagree will be classified as a barrier, whilst those coded as agree will be classified as a facilitator. The data will be used to assess the most frequently reported barriers and facilitators.

Responses will be stored within the secure, password protected Qualtrics account. Any relevant exports to facilitate the analysis of data will be stored in a secure location.

5.5.3 Qualitative study

Healthcare professionals responsible for conducting the VTE risk assessment will be invited to participate in an in-depth, one-to-one, semi-structured interview (remotely [e.g. telephone or Microsoft Teams] or face-to-face) with a member of the research team based at the co-ordinating centre (Imperial College London). Potential participants will be recruited via social media (Facebook, Twitter or Instagram) or via clinician networks. Potential participants will be invited to contact the study team if they are interested in taking part. They will be provided with the information sheet to further consider participation. Participants will be given as long as they deem necessary to consider participation. If the participant is happy to participate, they will be sent a consent form to complete. Following completion of the consent form, an interview will be arranged.

To allow data to approach saturation, up to 20 HCPs will be interviewed.

There may be some overlap between participants who participate in the quantitative study (survey) and those who participate in the qualitative study - e.g. some people may participate if they see a social media advert for the survey and the qualitative interviews and want to participate in both.

Participants will be sampled purposively across a range of professional backgrounds (surgical specialities and clinical grades). HCPs will be interviewed to explore attitudes towards the VTE Clinical Practice Guidelines and the perceived barriers and facilitators to adhering to VTE Clinical Practice Guidelines. There will be two parts to the interview. The first part is semi-structured, where participants will discuss their attitudes towards VTE clinical practice guidelines. This includes describing any barriers or facilitators to conducting the VTE risk assessment or prescribing the appropriate thromboprophylaxis. The second part of the interview is a structured questionnaire where participants will be asked about their practices around conducting the VTE risk assessment and prescribing thromboprophylaxis.

Topic guides will be used to ensure that similar topics are covered in each interview but applied in a flexible manner, enabling issues of importance to the participants to emerge. Interviews will be semi-structured and conducted in a conversational manner with probes, facilitation and clarification of responses as required by the interviewer. Respondent demographic characteristics will also be recorded.

5.5.4 Qualitative data analysis

All the interviews will be audio-recorded (with consent) and transcribed (by the study team or via Microsoft Teams transcription feature) verbatim. Two members of the study team will participate in data analysis to ensure transparency and credibility of the themes that are identified and their interpretation. Data will be analysed using thematic analysis. Researchers will familiarise themselves with the data through reading and re-reading the transcripts. The qualitative study will be conducted in line with and written up in accordance with the COREQ guidelines.

5.5.5 Data storage

Study data gathered by the coordinating centre will be kept in a secure location and in accordance with the terms of the Data Protection Act 2018. Participant confidentiality will be maintained at all times and all study records will be kept in pseudonymised form identifying participants by a pseudonym rather than name. At the point of consent, participants will be assigned a participant ID on the consent form which will allow for pseudonymisation.

Any paper interview transcripts or notes made by the coordinating centre during the interview will be kept in locked filing cabinets in locked rooms, only accessible by authorised personnel and destroyed at the end of the study. The key to the participant code list will be kept separately to these documents and destroyed at the end of the study. The audio recording will be destroyed after transcription.

6. STATISTICS AND DATA ANALYSIS

6.1 Sample size calculation

Good quality data for the incidence of VTE in low VTE risk surgical patients not receiving VTE prophylaxis is lacking. The symptomatic VTE occurrence at 90 days for day-case cholecystectomy and inguinal hernia repair is approximately 0.3-0.5% (21). These patients are

given GCS as standard practice. We are unable to determine how many of these patients received additional LMWH which would not be in line with NICE guidelines. Data from the recently published GAPS RCT on moderate and high risk patients reported 29/1858 VTE events at 90 days (1.6%) (13).

The occurrence of VTE for short stay procedures in individuals formally assessed as being at low risk of VTE would be expected to be lower than this, however, as these short stay patients are not receiving LMWH we have conservatively assumed an overall control (no stockings nor drugs) of 1.0%.

The sample size calculation is for a superiority comparison based on symptomatic VTE (DVT or PE) at 90 days(Y/N) at 90% power with a significance level of 5%, assuming 1.0% in the no stocking (control) and 0.5% in the stocking (active) groups. That is, we expect that the stockings would need to achieve an absolute 0.5% reduction (50% relative reduction from 1%) to be clinically and cost-effective.

For an individually randomised trial, 6256 participants per group are required (equalling a total of 12,512). We will instead use a parallel groups cluster design in at least 50 hospitals. If we assume an ICC on the VTE at 90 days outcome of 0.001 (consistent with the control outcome rate varying uniformly between 0.5% and 1.5%) (22) and a coefficient of variation of site size of 0.25 (consistent with sites varying according to a Normal distribution mean=350 and SD=88 i.e. 95% of the sites would be between 174 and 526 participants), with 25 sites randomised to active and 25 sites to control, we would need to recruit 17,500 patients.

The estimate was further increased to 18,251 participants to allow for one interim analysis at 50% mature day on the primary outcome under a two-sided, asymmetric, group sequential design implementing the non-binding Hwang-Shih-DeCani spending function [lower bounds (futility) with $\gamma=-2$; upper bound (efficacy) with $\gamma=-4$].

Finally, to allow for 15% missing data on the primary outcome the sample size was inflated to 21,472 participants.

6.2 Statistical analyses

Data will be reported according to the CONSORT statement for cluster trials (Campbell et al 2012). Analyses will primarily be intention-to-treat where all consented participants will be included in the analysis retained in the group to which they were allocated (i.e. "as-randomised") and for whom outcome data are available. All results will be presented as point estimates, confidence intervals at the appropriate level with associated p-values. Absolute measures of effects will be presented alongside relative measures. A sensitivity analysis using imputation of missing values will be considered only if the proportion of cases with missing values is sufficiently large. All statistical analyses will be governed by a comprehensive Statistical Analysis Plan (SAP), authored by the study statistician(s) and agreed by the independent Trial Steering Committee (TSC).

The primary analysis will compare the occurrence of VTE at 90 days post surgery, measured at an individual level, using a hierarchical (multilevel) logistic regression, adjusting for any other pre-specified strongly prognostic individual baseline covariates and site level baseline covariates, with site itself as a random effect (to account for cluster of participants within sites). The intervention effect will be presented by the adjusted odds ratio (and corresponding

confidence interval and p-value) for GCS vs. no GCS as well as the ICC for study centre. Unadjusted results will also be presented to support the findings of the primary analysis.

Secondary outcomes will be analysed in a similar fashion with generalized linear models appropriate to the distribution of the outcome. Safety data will be summarised descriptively

6.3 Subgroup analyses

We will perform the following sub-group analyses of the primary outcome, and test for sub-group interactions if appropriate:

Predefined sub-group analyses (primary outcome) include:

- Sex (male, female)
- Age
- BMI
- Smoking status (ex-smoker, current smoker, never smoker)
- Type of anaesthesia (local, general)

The study is not formally powered for these subgroups analyses, and will be exploratory.

6.4 Interim analysis

We have scheduled 1 interim analysis which will consider the primary outcome of VTE at 90 days post-surgery at 50% of recruitment. Full details of the stopping boundaries and analysis will be included in the SAP, which the iDMC will approve prior to seeing unblinded data. This interim analysis will be presented by the unblinded statistician at ECTU to the iDMC, who will include an independent statistician. The unblinded ECTU statistician would have no other role in the study while it was ongoing. The stopping rules are statistically non-binding. The iDMC may recommend early stopping of the study if the boundaries are crossed. They would make a recommendation to the independent Trial Steering Committee (TSC) who may or may not endorse that recommendation. Note also that the trial can stop at any time for safety, if there is an excess of events in the intervention group that is considered to generate avoidable harm – and this would be a decision not based on any statistical criterion and taken by the iDMC and endorsed by the TSC.

6.5 Measurement of cost-effectiveness

Two health economic analyses will be conducted. The main analyses will be performed from the perspective of the NHS and Personal Social Services, with secondary analyses from a societal perspective.

- 1) A within-trial analysis will compare GCS to no thromboprophylaxis over the 90 days of the study. Resource use items associated with treatments in hospital and community care will be collected using case notes and self-completed patient resource use diaries over the 90-day follow-up, and costed using manufacturers' list prices, previous literature and national reference costs. Days off work and normal activities and other patient-related costs will be collected for a secondary analysis. EQ-5D will be collected at baseline and follow up, analysed using the NICE approved tariff. Appropriate methods will be used to handle missing data and any relevant subgroups in line with the SAP.

- 2) If there are clinically relevant and measurable differences in VTE or quality of life between the study arms at 90 days, a Markov (state–transition) decision model will be constructed to compare the ICER up to 2 years for GCS versus no thromboprophylaxis. The time horizon of the model will be 2 years allowing extrapolation of sequelae of VTE events (such as PTS) over the longer term to quantify the impact of VTE on patient health (quality adjusted life years) and resource use. A preliminary model has been constructed based on published literature to identify the key variables that would need to be collected during the clinical study, and to estimate the number needed to treat (NNT) to avoid 1 VTE, above which GCS would not be considered cost-effective at NICE thresholds. The 2-year time point was chosen as we know from previous research that the incidence of PTS after acute DVT levels out after the first year.

This preliminary model assumes 30% of patients with VTE develop PTS, with 3% of those patients having severe PTS. The cost of purchasing and applying GCS stockings are approximately £22.46 (23). The model assumes the cost of treatment of VTE, non-severe and severe PTS, are £451, £872 and £1547 respectively, and estimates of the utility decrement associated with symptomatic VTE and PTS which are 0.8628, 0.7745 and 0.6752 respectively (24). Using a 2-year time horizon, the ICER of GCS versus no prophylaxis would be £20,603 per quality-adjusted life year (QALY) if the NNT were 200 participants. Hence, for GCS to be cost-effective at a NICE willingness to pay threshold, the NNT would need to be below 200 (see sample size).

The health economic analyses will be conducted and reported according to NICE reference case and CHEERS guidelines (25,26), including sensitivity analyses and probabilistic sensitivity analyses. The results will be presented as estimates of mean incremental costs, effects, and incremental cost per QALY.

7. ADVERSE EVENTS

7.1. DEFINITIONS

Adverse Event (AE): any untoward medical occurrence in a patient or clinical study subject.

Serious Adverse Event (SAE): any untoward medical occurrence or effect that:

- **Results in death**
- **Is life-threatening** – *refers to an event in which the subject was at risk of death at the time of the event; it does not refer to an event which hypothetically might have caused death if it were more severe*
- **Requires hospitalisation, or prolongation of existing inpatients' hospitalisation**
- **Results in persistent or significant disability or incapacity**
- **Is a congenital anomaly or birth defect**

Medical judgement should be exercised in deciding whether an AE is serious in other situations. Important AEs that are not immediately life-threatening or do not result in death or hospitalisation but may jeopardise the subject or may require intervention to prevent one of the other outcomes listed in the definition above, should also be considered serious.

7.2. REPORTING PROCEDURES

For the purposes of the study, only AEs related to the study intervention (graduated compression stockings) will be recorded. This information will be collected at 7-days only. Participants enrolled in the center which has been randomized to the stockings allocation are

only required to wear the stockings for the duration of the short-stay surgery (donning the stockings just prior to undergoing the surgery and removing when ambulant), hence AEs associated with the stockings are only expected in the short-term and thus collected at 7-days. Participants who report a treatment related AE which requires further investigation/follow-up will be advised to consult with their GP/relevant clinical team where necessary. Depending on the nature of the event the reporting procedures below should be followed. Any questions concerning adverse event reporting should be directed to the Chief Investigator in the first instance.

7.2.1 Non serious AEs

Only AEs related to the study intervention (graduated compression stockings) will be recorded.

7.2.2 Serious AEs

An SAE form should be completed and emailed to the Chief Investigator within 24 hours. However, hospitalisations for elective treatment of a pre-existing condition do not need reporting as SAEs.

All SAEs should be reported to the London - Camden & Kings Cross Research Ethics Committee where in the opinion of the Chief Investigator, the event was:

- 'related', ie resulted from the administration of any of the research procedures; and
- 'unexpected', ie an event that is not listed in the protocol as an expected occurrence

Reports of related and unexpected SAEs should be submitted within 15 days of the Chief Investigator becoming aware of the event, using the NRES SAE form for non-IMP studies. The Chief Investigator must also notify the Sponsor of all related and unexpected SAEs.

Local investigators should report any SAEs as required by their Local Research Ethics Committee, Sponsor and/or Research & Development Office.

Contact details for reporting SAEs

RGIT@imperial.ac.uk

Please send SAE forms to: petstrial@imperial.ac.uk

Tel: +44 (0)203 311 7371 (Mon to Fri 09.00 – 17.00)

8. REGULATORY ISSUES

8.1. ETHICS APPROVAL

The Study Coordination Centre has obtained approval from the London - Camden & Kings Cross Research Ethics Committee (REC) and Health Research Authority (HRA). The study must also receive confirmation of capacity and capability from each participating NHS Trust before accepting participants into the study or any research activity is carried out. The study will be conducted in accordance with the recommendations for physicians involved in research

on human subjects adopted by the 18th World Medical Assembly, Helsinki 1964 and later revisions.

8.2. CONSENT

Consent to enter the study will be sought from each participant only after a full explanation has been given, an information leaflet offered and time allowed for consideration. Consent will be obtained in written, verbal or electronic form. The right of the participant to refuse to participate without providing any reason will be respected. After the participant has entered the study, the clinician remains free to give alternative treatment to that specified in the protocol at any stage if he/she feels it is in the participant's best interest, but the reasons for doing so should be recorded. In these cases the participants remain within the study for the purposes of follow-up and data analysis. All participants are free to withdraw at any time from the protocol treatment without giving reasons and without prejudicing further treatment.

8.3. CONFIDENTIALITY

The Chief Investigator will preserve the confidentiality of participants taking part in the study and is registered under the Data Protection Act.

As follow-ups will be performed by the coordinating centre (Imperial College London), patient identifiable data (name, address, email address and contact telephone number[s]) will be stored on the REDCap database. This identifiable data will only be accessible by researchers at the local site (who will enter the data onto REDCap in the first place) and by the blinded assessors based at the coordinating centre who are responsible for conducting the follow-ups.

The trial manager will only have access to pseudonymised data on REDCap.

The investigator shall permit direct access to subjects' records and source document for the purposes of monitoring, auditing, or inspection by the Sponsor, authorised representatives of the Sponsor and RECs.

Pseudonymised data will be transferred to the University of Granada for the purposes of conducting the cost-effectiveness analysis.

8.4. INDEMNITY

Imperial College London holds negligent harm and non-negligent harm insurance policies which apply to this study.

8.5. SPONSOR

Imperial College London will act as the main Sponsor for this study. Delegated responsibilities will be assigned to the NHS trusts or non-NHS site taking part in this study.

8.6. FUNDING

The National Institute for Health Research (NIHR) are funding this study.

8.7. AUDITS

The study may be subject to audit by Imperial College London under their remit as sponsor and other regulatory bodies to ensure adherence to GCP and the UK Policy Frame Work for Health and Social Care Research.

9. DATA MANAGEMENT

9.1. DATA COLLECTION

Details of procedures for CRF/eCRF completion will be provided in a study manual.

9.2. ARCHIVING

Data and all appropriate documentation will be stored for a minimum of 10 years after the completion of the study, including the follow-up period.

10. STUDY MANAGEMENT

The day-to-day management of the study will be co-ordinated through Imperial College London.

11. PUBLICATION POLICY

The following outputs are anticipated to arise from the PETS trial:

- Publications in peer reviewed journals (including the protocol paper, main trial analysis and cost-effectiveness analysis)
- The NICE guidelines aiming to prevent VTE “Venous thromboembolism in over 16s: reducing the risk of hospital-acquired deep vein thrombosis or pulmonary embolism [NG89]” will be subsequently updated
- Cost-effectiveness information to guide clinical commissioning groups and NICE
- Updated systematic review of literature and meta-analysis
- An understanding of the safety of GCS and subsequent quality of life
- Presentation at international academic conferences including European and American vascular, venous, general surgery and haematology societies
- Dissemination of results to the wider public through social media streams
- If GCS are found to be ineffective, this could prompt a wider effect on the design and application of graduated compression stocking devices within the healthcare setting

12. REFERENCES

1. 5.1 Deaths from venous thromboembolism (VTE) related events within 90 days post discharge from hospital [Internet]. NHS Digital. [cited 2022 Mar 29]. Available from: <https://digital.nhs.uk/data-and-information/publications/statistical/nhs-outcomes-framework/march-2022/domain-5---treating-and-caring-for-people-in-a-safe-environment-and-protecting-them-from-avoidable-harm-nof/5.1-deaths-from-venous-thromboembolism-vte-related-events-within-90-days-post-discharge-from-hospital>
2. Cohen AT, Agnelli G, Anderson FA, Arcelus JI, Bergqvist D, Brecht JG, et al. Venous thromboembolism (VTE) in Europe. The number of VTE events and associated morbidity and mortality. *Thromb Haemost.* 2007 Oct;98(4):756–64.
3. The Prevention of Venous Thromboembolism in Hospitalised Patients Second Report of Session 2004-05.

4. Kahn SR, Shapiro S, Wells PS, Rodger MA, Kovacs MJ, Anderson DR, et al. Compression stockings to prevent post-thrombotic syndrome: a randomised placebo-controlled trial. *Lancet Lond Engl*. 2014 Mar 8;383(9920):880–8.
5. Eklof B, Perrin M, Delis KT, Rutherford RB, Gloviczki P, American Venous Forum, et al. Updated terminology of chronic venous disorders: the VEIN-TERM transatlantic interdisciplinary consensus document. *J Vasc Surg*. 2009 Feb;49(2):498–501.
6. Anderson FA, Spencer FA. Risk factors for venous thromboembolism. *Circulation*. 2003 Jun 17;107(23 Suppl 1):I9-16.
7. Geerts WH, Bergqvist D, Pineo GF, Heit JA, Samama CM, Lassen MR, et al. Prevention of venous thromboembolism: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest*. 2008 Jun;133(6 Suppl):381S-453S.
8. Gould MK, Garcia DA, Wren SM, Karanicolas PJ, Arcelus JL, Heit JA, et al. Prevention of VTE in nonorthopedic surgical patients: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012 Feb;141(2 Suppl):e227S-e277S.
9. Sweetland S, Green J, Liu B, Berrington de González A, Canonico M, Reeves G, et al. Duration and magnitude of the postoperative risk of venous thromboembolism in middle aged women: prospective cohort study. *BMJ*. 2009 Dec 3;339:b4583.
10. Hair A, Duffy K, McLean J, Taylor S, Smith H, Walker A, et al. Groin hernia repair in Scotland. *Br J Surg*. 2000 Dec;87(12):1722–6.
11. Overview | Venous thromboembolic diseases: diagnosis, management and thrombophilia testing | Guidance | NICE [Internet]. NICE; [cited 2022 Mar 29]. Available from: <https://www.nice.org.uk/guidance/ng158>
12. Compression stockings | Health topics A to Z | CKS | NICE [Internet]. [cited 2022 Mar 29]. Available from: <https://cks.nice.org.uk/topics/compression-stockings/>
13. Shalhoub J, Lawton R, Hudson J, Baker C, Bradbury A, Dhillon K, et al. Graduated compression stockings as adjuvant to pharmaco-thromboprophylaxis in elective surgical patients (GAPS study): randomised controlled trial. *BMJ [Internet]*. 2020 May 13 [cited 2020 Sep 25];369. Available from: <https://www.bmj.com/content/369/bmj.m1309>
14. CLOTS Trials Collaboration, Dennis M, Sandercock P a. G, Reid J, Graham C, Murray G, et al. Effectiveness of thigh-length graduated compression stockings to reduce the risk of deep vein thrombosis after stroke (CLOTS trial 1): a multicentre, randomised controlled trial. *Lancet Lond Engl*. 2009 Jun 6;373(9679):1958–65.
15. Hill J, Treasure T. Reducing the risk of venous thromboembolism (deep vein thrombosis and pulmonary embolism) in inpatients having surgery: summary of NICE guidance. *BMJ*. 2007 May 19;334(7602):1053–4.

16. Overview | Venous thromboembolism in over 16s: reducing the risk of hospital-acquired deep vein thrombosis or pulmonary embolism | Guidance | NICE [Internet]. NICE; [cited 2022 Mar 29]. Available from: <https://www.nice.org.uk/guidance/ng89>
17. Machin M, Younan HC, Smith S, Salim S, Davies AH, Shalhoub J. Systematic review on the benefit of graduated compression stockings in the prevention of venous thromboembolism in low-risk surgical patients. *Phlebology*. 2021 Apr;36(3):184–93.
18. Graduated compression stockings for prevention of deep vein thrombosis - Sachdeva, A - 2018 | Cochrane Library [Internet]. [cited 2022 Mar 29]. Available from: <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD001484.pub4/full>
19. EuroQol Group. EuroQol--a new facility for the measurement of health-related quality of life. *Health Policy Amst Neth*. 1990 Dec;16(3):199–208.
20. Department of Health. Risk assessment for Venous Thromboembolism (VTE) [Internet]. 2010 [cited 2021 May 11]. Available from: <http://www.nice.org.uk/guidance/CG92>
21. Bouras G, Burns EM, Howell A-M, Bottle A, Athanasiou T, Darzi A. Risk of Post-Discharge Venous Thromboembolism and Associated Mortality in General Surgery: A Population-Based Cohort Study Using Linked Hospital and Primary Care Data in England. *PloS One*. 2015;10(12):e0145759.
22. Measures of between-cluster variability in cluster randomized trials with binary outcomes - Thomson - 2009 - Statistics in Medicine - Wiley Online Library [Internet]. [cited 2022 Mar 29]. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/sim.3582>
23. Unit Costs of Health and Social Care 2019 | PSSRU [Internet]. [cited 2022 Mar 29]. Available from: <https://www.pssru.ac.uk/project-pages/unit-costs/unit-costs-2019/>
24. Enden T, Resch S, White C, Wik HS, Kløw NE, Sandset PM. Cost-effectiveness of additional catheter-directed thrombolysis for deep vein thrombosis. *J Thromb Haemost JTH*. 2013 Jun;11(6):1032–42.
25. Guide to the methods of technology appraisal 2013. 2013;94.
26. Husereau D, Drummond M, Petrou S, Carswell C, Moher D, Greenberg D, et al. Consolidated Health Economic Evaluation Reporting Standards (CHEERS) statement. *BMJ* [Internet]. 2013 Mar 25 [cited 2020 May 15];346. Available from: <https://www.bmj.com/content/346/bmj.f1049>

Appendix 1. Summary of investigations, treatment and assessments

Assessment	Week of assessment			
	Pre-surgical procedure	0 (Day of procedure)	1	12
Inclusion/Exclusion	X	X		

Informed consent¹	X			
Screening assessments	X			
Demographics¹		X		
Medical history (including concurrent medications)¹		X	X	X
Provision of leaflet explaining the signs and symptoms of developing a blood clot	X			
Provision of stockings*		X		
EQ-5D		X	X	X
VTE outcome (self-reported)			X	X
Adverse Events Assessment*			X	
Resource use information				X

*Sites randomised to the intervention arm only

¹Can be collected at any point up to the 7-day follow-up, including prior to the day of procedure (i.e. at the preassessment stage) or on the day of procedure