


Figure 1: Flowchart showing the number of patients in each stage of the study - CONSORT



**Table 1. Background characteristics of patients by intervention groups**

Characteristics	Intervention groups			p
	Total (n=360)	Estradiol oral (n=182)	Estradiol gel (n=178)	
	n (%)	n (%)	n (%)	
<b>Age (Years)<sup>Y</sup></b>	32,0 (4,3)	32,3 (4,5)	31,7 (4,1)	0,141
<b>Age groups (years)</b>				
<25	9 (2,5)	4 (2,2)	5 (2,8)	0,273 <sup>c</sup>
25-34	259 (71,9)	127 (69,8)	132 (74,2)	
35-40	82 (22,8)	43 (23,6)	39 (21,9)	
>40	10 (2,8)	8 (4,4)	2 (1,1)	
<b>Age groups (years)</b>				
>30	217 (60,3)	115 (63,2)	102 (57,3)	0,254
≤30	143 (39,7)	67 (36,8)	76 (42,7)	
<b>Type of infertility</b>				
Primary	215 (59,7)	106 (58,2)	109 (61,2)	0,563
Secondary	145 (40,3)	76 (41,8)	69 (38,8)	
<b>IVF indications</b>				
Ovulation disorder	73 (20,3)	37 (20,3)	36 (20,2)	0,362 <sup>c</sup>
Tubal	95 (26,4)	48 (26,4)	47 (26,4)	
Male factor infertility	102 (28,3)	46 (25,3)	56 (31,5)	
Unexplained	30 (8,3)	15 (8,2)	15 (8,4)	
Others	58 (16,1)	36 (19,8)	22 (12,4)	
<b>Duration infertility (months)<sup>Y</sup></b>	53,3 (34,1)	53,4 (34,0)	53,3 (34,4)	0,965
<b>Duration infertility (years)<sup>Y</sup></b>	4,4 (2,8)	4,5 (2,8)	4,4 (2,9)	0,965
<b>Duration infertility (years)</b>				
≤1	32 (8,9)	13 (7,1)	19 (10,7)	0,395
>1-<5	192 (53,3)	102 (56,0)	90 (50,6)	
≥5	136 (37,8)	67 (36,8)	69 (38,8)	
<b>BMI<sup>Y</sup></b>	22,1 (3,1)	22,2 (3,2)	22,1 (3,1)	0,794
<b>BMI group</b>				
<18,5	24 (6,7)	14 (7,7)	10 (5,6)	0,713
18,5-<23	222 (61,7)	110 (60,4)	112 (62,9)	
>23	114 (31,7)	58 (31,9)	56 (31,5)	

<sup>Y</sup>Mean &SD<sup>c</sup> Fisher 's exact test

**Table 2. Cycle characteristics of patients by intervention groups**

Cycle characteristics	Intervention groups			p
	Total (n=360) n (%)	Estradiol oral (n=182) n (%)	Estradiol gel (n=178) n (%)	
<b>Total amount of drugs use estradiol<sup>y</sup></b>	19,4 (2,3)	19,4 (2,2)	19,5 (2,4)	0,821
<b>Total amount of drugs use estradiol<sup>y</sup></b>	63,5 (20,8)	63,0 (20,5)	64,0 (21,2)	0,670
<b>Serum Estradiol on progesterone day (pg/ml)<sup>§</sup></b>	232,5 (155,5 - 315,0)	270,5 (197,0 - 394,0)	186,5 (126,0 - 258,0)	<0,001
<b>Endometrium thickness<sup>y</sup></b>	10,5 (1,2)	10,5 (1,1)	10,6 (1,3)	0,850
<b>Endometrium thickness on 7<sup>th</sup> day of cycle<sup>y</sup></b>	7,3 (6,8 - 8,0)	8,0 (7,2 - 8,1)	7,0 (6,3 - 7,5)	<0,001
<b>Type of embryos</b>				
Cleavage embryo	34 (9,4)	21 (11,5)	13 (7,3)	0,170
Blastocyst	326 (90,6)	161 (88,5)	165 (92,7)	
<b>Number of embryos transfer</b>				
1	309 (85,8)	156 (85,7)	153 (86,0)	0,999 <sup>c</sup>
≥2	50 (13,9)	25 (13,7)	25 (14,0)	
<b>Number of excellent embryos transfer</b>				
0	92 (25,6)	41 (22,5)	51 (28,7)	0,418 <sup>c</sup>
1	265 (73,6)	139 (76,4)	126 (70,8)	
2	3 (0,8)	2 (1,1)	1 (0,6)	
<b>Number of good embryos transfer</b>				
0	266 (73,9)	139 (76,4)	127 (71,3)	0,388 <sup>c</sup>
1	90 (25,0)	42 (23,1)	48 (27,0)	
2	4 (1,1)	1 (0,5)	3 (1,7)	
<b>Number of average embryos transfer</b>				
0	316 (87,8)	161 (88,5)	155 (87,1)	0,853 <sup>c</sup>
1	41 (11,4)	20 (11,0)	21 (11,8)	
2	3 (0,8)	1 (0,5)	2 (1,1)	

<sup>y</sup>Mean & SD<sup>§</sup>Median and interquartile range<sup>c</sup> Fisher's exact test

**Table 3. Outcome measures of cycles of patients by intervention groups**

Outcome	Intervention groups			p
	Total (n=360) n (%)	Estradiol oral (n=182) n (%)	Estradiol gel (n=178) n (%)	
<b>Pregnancy rate (positive HCG)</b>				
positive	204 (56,7)	99 (54,4)	105 (59,0)	0,379
negative	156 (43,3)	83 (45,6)	73 (41,0)	
<b>Clinical pregnancy rate (n = 208)</b>				
positive	195 (93,8)	95 (95,0)	100 (92,6)	0,474
negative	13 (6,3)	5 (5,0)	8 (7,4)	
<b>Ectopic pregnancy (n = 208)</b>				
positive	4 (1,9)	2 (2,0)	2 (1,9)	0,999 <sup>c</sup>
negative	204 (98,1)	98 (98,0)	106 (98,1)	
<b>Ongoing pregnancy rate (n = 208)</b>				
positive	178 (85,6)	86 (86,0)	92 (85,2)	0,867
negative	30 (14,4)	14 (14,0)	16 (14,8)	
<b>Miscarriage rate (n = 208)</b>				
positive	33 (15,9)	18 (18,0)	15 (13,9)	0,417
negative	175 (84,1)	82 (82,0)	93 (86,1)	

<sup>c</sup> Fisher's exact test

**Table 4: Side effects of patients by intervention groups**

	Intervention groups			p
	Total (n=360)	Estradiol oral (n=182)	Estradiol gel (n=178)	
	n (%)	n (%)	n (%)	
<b>Side effects</b>				
Yes	55 (15,3)	37 (20,3)	18 (10,1)	0,007
No	305 (84,7)	145 (79,7)	160 (89,9)	
<b>Side effects</b>				
No	305 (84,7)	145 (79,7)	160 (89,9)	<0,001 <sup>c</sup>
Headache	11 (3,1)	10 (5,5)	1 (0,6)	
Nausea	6 (1,7)	5 (2,7)	1 (0,6)	
Epigastric pain	22 (6,1)	18 (9,9)	4 (2,2)	
Itchy skin	12 (3,3)	0 (0)	12 (6,7)	
Hot flashes	4 (1,1)	4 (2,2)	0 (0)	

<sup>c</sup> Fisher's exact test