



Circulating tumour DNA guided Adaptive BRAF and MEK Inhibitor therapy (DyNAMIC)

DyNAMIC Protocol

V8.0, 14/01/2026

CTA Reference Number: 21266/0255/001-0001

ISRCTN: ISRCTN14643179

ClinicalTrials.gov: NCT06470880

Research Ethics Ref: 22/LO/0453

Sponsor Ref: CFTSp175

Funder Ref: 151

IRAS Number: 1004759

CPMS Ref: 55327

Study Sponsor:

The Christie NHS Foundation Trust

550 Wilmslow Road

Manchester

M20 4BX

Chief Investigator:

Professor Paul Lorigan

Consultant Medical Oncology

The Christie NHS Foundation Trust

The funder of this research trial had no role in the design of this study and will not have any role during its execution, analyses, interpretation of the data, or decision to submit results.

Protocol Approval

I, the undersigned, hereby approve this clinical study protocol:

Authorised by Chief Investigator:

Signature:  **Date:** 14 Jan 2026

box SIGN 462Z92J4-4LX3YZ9Q

Professor Paul Lorigan

Consultant in Medical Oncology

The Christie NHS Foundation Trust

Authorised on behalf of The Sponsor:

Signature:  **Date:** 22 Jan 2026

box SIGN 1J68K7L1-4LX3YZ9Q


Dr Clare Griffin

Senior Research Governance Manager (Sponsorship and Regulatory Affairs)

Sponsor Representative

The Christie NHS Foundation Trust

Authorised on behalf of University of Liverpool:

Signature:  **Date:** 14 Jan 2026

box SIGN 42YWJYX1-4LX3YZ9Q

Dr Ashley Jones

Lead Statistician

University of Liverpool

General Information

For the purposes of clarity, use of the words 'Study' and 'Trial' are interchangeable synonyms, and their usage will differ depending on the context in which they are used.

Similarly, as numerous institutions and systems are involved in the study, interchangeable terminology, e.g. Patient Trial ID, Participant Study Number etc. are also used in this and other associated documents as synonyms of the pseudonymised reference number allocated to participants.

This document describes the DyNAMIC trial including detailed information about procedures and recruitment. The protocol should not be used as an aide-memoir or guide for the treatment of other patients. Every care was taken in its drafting, but corrections or amendments may be necessary. Any amendments will be circulated to the investigators participating in the trial, but sites entering participants for the first time are advised to contact Liverpool Clinical Trials Centre (LCTC), the coordinating clinical trials unit, to confirm they have the most up to date version. Clinical problems relating to this trial should be referred to the relevant Chief Investigator, Professor Paul Lorigan, via the LCTC.

This protocol defines the participant characteristics required for study entry and the schedule of treatment and follow-up. Participant recruitment will be undertaken in compliance with this document and applicable regulatory and governance requirements. Waivers to authorise non-compliance are not permitted.

Incidence of protocol non-compliance whether reported prospectively (e.g. where a treatment cannot be administered on a scheduled date as a result of public holidays) or retrospectively noted (e.g. as a result of central monitoring) are recorded as protocol deviations. These are monitored and reported to trial oversight committees.

The template content structure is consistent with the SPIRIT (Standard Protocol Item: Recommendations for Interventional Trials 2013) and has regard for the Health Research Authority guidance. Regulatory and ethical compliance information is located in Section [1313](#).

The LCTC has achieved full registration by the UK Clinical Research Collaboration (www.ukcrc.org) as their standards and systems were assessed by an international review panel as reaching the highest quality. The LCTC has a diverse trial portfolio underpinned by methodological rigour, a GCP compliant data management system, and quality management system.

Contact Details: Institutions		
Sponsor:	Clinical Trials Unit:	Pharmacy:
The Christie NHS Foundation Trust, Wilmslow Road, Manchester, M20 4BX ☎ 0161 918 7432 ✉ the-christie.sponsoredresearch@nhs.net	Liverpool Clinical Trials Centre, University of Liverpool, Block C, Waterhouse Building, 3 Brownlow Street, Liverpool, L69 3GL ☎ 0151 795 1400 ✉ dynamic_study@liverpool.ac.uk	The Christie NHS Foundation Trust, Oak Road Pharmacy, Wilmslow Road, Manchester, M20 4BX ☎ 0161 956 1065 ✉ Lydia.sutherland1@nhs.net
Central Laboratory:		
Cancer Research UK National Biomarker Centre, The University of Manchester, Wilmslow Road, Manchester, M20 4BX ☎ 0161 200 8925 ✉ ceptrials@cruk.manchester.ac.uk		

Contact Details: Individuals		
Individual Authorised to Sign the Protocol and Protocol Amendments on behalf of the Sponsor:	Chief Investigator (CI):	
Clare Griffin, The Christie NHS Foundation Trust, Wilmslow Road, Manchester, M20 4BX ☎ 0161 918 7432 ✉ clare.griffin1@nhs.net	Prof Paul Lorigan MB MD FRCP, The University of Manchester and The Christie NHS Foundation Trust, Wilmslow Road, M20 4BX ☎ 0161 446 8002 ✉ Paul.Lorigan@nhs.net	
Medical Expert who will Advise on Protocol Related Clinical Queries:	Alternative Medical Expert who will Advise on Protocol Related Clinical Queries:	Alternative Medical Expert who will Advise on Protocol Related Clinical Queries:
Chief Investigator Details as above	Dr Rebecca Lee MbChb PhD, The University of Manchester and The Christie NHS Foundation Trust, Wilmslow Road, M20 4BX ☎ 07764801907 ✉ Rebecca.lee-3@manchester.ac.uk	Dr Avinash Gupta, The Christie NHS Foundation Trust, Wilmslow Road, Manchester, M20 4BX ☎ 0161 446 3000 (switchboard) ✉ Avinash.gupta@nhs.net

Contact Details: Individuals		
Medical Expert who will Evaluate SAE Reports:	Alternative Medical Expert who will Evaluate SAE Reports:	Alternative Medical Expert who will Evaluate SAE Reports:
Chief Investigator Details as above	Dr Rebecca Lee MbChb PhD, The University of Manchester and The Christie NHS Foundation Trust, Wilmslow Road, M20 4BX ☎ 07764801907 ✉ Rebecca.lee-3@manchester.ac.uk	Dr Avinash Gupta, The Christie NHS Foundation Trust, Wilmslow Road, Manchester, M20 4BX ☎ 0161 446 3000 (switchboard) ✉ Avinash.gupta@nhs.net
	Dr Antonio Rullan, Royal Free London NHS Foundation Trust, Royal Free Hospital, Pond Street, London, NW3 2QG ☎ 020 3758 2000 (ext. 35839) ✉ Antonio.rullan@nhs.net	Dr Kate Young, Renal/Melanoma Unit, The Royal Marsden NHS Foundation Trust, Fulham Road, London, SW3 6JJ ☎ 020 7811 8151 ✉ kate.young@rmh.nhs.uk

Additional Contacts:

The contact details for the trial oversight committee members and participating sites are detailed in documents supplementary to the protocol and stored in the Trial Master File.

Table of Contents

1.	GLOSSARY	10
2.	PROTOCOL OVERVIEW	12
3.	ROLES AND RESPONSIBILITIES	17
3.1	Sponsor	17
3.2	Funder	17
3.3	The Study Team	17
3.4	Oversight Committees	17
3.5	Protocol Contributors	18
4	INTRODUCTION	19
4.1	Background	19
4.2	Rationale	26
4.3	Risk and Benefits	29
4.4	Trial Hypothesis, Objectives and Outcomes	31
5	STUDY DESIGN	35
5.1	Molecular tumour board ongoing assessment of TAB levels.....	36
5.2	Blinding	38
5.3	Study Setting.....	39
6	STATISTICAL CONSIDERATIONS	40
6.1	Introduction	40
6.2	Sample Size and Recruitment	40
6.3	Method of Randomisation.....	41
6.4	Interim Analyses	41
6.5	Analysis Plan.....	42
7	ELIGIBILITY CRITERIA.....	43
7.1	Inclusion Criteria at Screening.....	43
7.2	Inclusion Criteria prior to randomisation.....	43
7.3	Exclusion Criteria	44
7.4	Co-enrolment Guidelines	44
7.5	Lifestyle Restrictions	45
8	TRIAL TREATMENT / INTERVENTIONS	47
8.1	Introduction	47
8.2	Treatment Name / Description.....	47
8.3	Manufacturing and Distribution	50
8.4	Preparation and Administration	53
8.5	Treatment Modifications	53
8.6	Accountability Procedures	53
8.7	Assessment of Compliance	53
8.8	Product Recall.....	53
8.9	Concomitant Medications/Treatments and Specific Restrictions.....	54
8.10	Overdose.....	55
9	PARTICIPANT TIMELINES AND ASSESSMENTS.....	57
9.1	Identification and Screening	57
9.2	Informed Consent	58
9.3	DyNAMIC Clinical Study Database	59
9.4	Assessments and Confirmation of Eligibility (Screening)	60
9.5	Baseline 1 Assessments.....	61
9.6	Commencement of Initial Targeted 28 day E+B Cycle	63
9.7	Randomisation	63
9.8	Intervention	63
9.9	Schedule for Assessments and Follow-up - Arm A.....	63
9.10	Schedule for Assessments and Follow-up – Arm B.....	72
9.11	Definitions of Assessments:.....	82
9.12	Disease progression definition and procedures	86
9.13	Sampling	88
9.14	Intervention Discontinuation and Participant Discontinuation/Withdrawal.....	90
9.15	End of Trial.....	92
10	Management of COVID-19	93

11	SAFETY REPORTING	94
11.1	Terms and Definitions	94
11.2	Assessment of Seriousness	95
11.3	Severity of Adverse Events	95
11.4	Assessment of "Causality" - Relationship to Trial Treatment/Intervention	96
11.5	Assessment of "Expectedness"	97
11.6	Time Period for Active Monitoring of Safety Events	98
11.7	Notes on Safety Event Recording	98
11.8	Reporting of Pregnancy	99
11.9	Notification of Deaths	100
11.10	Reporting Procedures	100
11.11	Investigator Reporting Responsibilities	102
11.12	LCTC Responsibilities	104
11.13	Contact Details and Out-of-hours Medical Cover	106
12	DATA MANAGEMENT AND TRIAL MONITORING	107
12.1	Source Documents	107
12.2	Data Collection Methods	107
12.3	Monitoring	107
12.4	Risk Assessment	108
12.5	Confidentiality	108
12.6	Quality Assurance and Control	109
12.7	Records Retention	109
13	REGULATORY AND ETHICAL CONSIDERATIONS	111
13.1	Statement of Compliance	111
13.2	Ethical Considerations	111
13.3	Approvals	111
13.4	Non-Compliances	111
14	INDEMNITY	113
15	PUBLICATION AND DISSEMINATION	114
15.1	Publication Policy	114
15.2	Dissemination to Key Stakeholders	114
15.3	Data Sharing	114
16	CHRONOLOGY OF PROTOCOL AMENDMENTS	115
16.1	Version 1.0 (05/05/2022)	115
16.2	Version 2.0	115
16.3	Version 3.0 (28/11/2022)	115
16.4	Version 4.0 (27/03/2023)	115
16.5	Version 5.0 (21/05/2024)	116
16.6	Version 6.0 (05/03/2025)	117
16.7	Version 7.0 (23/06/2025)	118
16.8	Version 8.0 (14/01/2026)	119
17	REFERENCES	122
18	DOCUMENTS SUPPLEMENTARY TO THE PROTOCOL	130
19	APPENDICES	131

1. GLOSSARY

AE	Adverse Event
AR	Adverse Reaction
cfDNA	Cell free DNA
CI	Chief Investigator
CRUK	Cancer Research UK
CRUK NBC	Cancer Research UK National Biomarker Centre
CTA	Clinical Trial Authorisation
CTCAE	Common Terminology Criteria for Adverse Events
ctDNA	Circulating Tumour DNA
CTIMP	Clinical Trial of an Investigational Medicinal Product
ddPCR	Droplet Digital PCR
DNA	Deoxyribonucleic acid
ECG	Electrocardiograph
ECOG	Eastern Cooperative Oncology Group
eCRF	Electronic Case Report Form
EU	European Union
EUCTD	European Clinical Trials Directive
FFPE	Formalin Fixed Paraffin Embedded
GCP	Good Clinical Practice
GP	General Practitioner
HCP	Health Care Professional
HRA	Health Research Authority
IB	Investigator's Brochure
ICH	International Conference on Harmonisation
ICO	Information Commissioner's Office
IDSMC	Independent Data and Safety and Monitoring Committee
IMP	Investigational Medicinal Product
ISF	Investigator Site File
ISRCTN	International Standard Randomised Controlled Trials Number
LCTC	Liverpool Clinical Trials Centre
LDH	Lactate Dehydrogenase
MA	Marketing Authorisation
MHRA	Medicines and Health Care Products Regulatory Agency
MTA	Material Transfer Agreement

NBC	National Biomarker Centre
NHS	National Health Service
NIHR CRN	National Institute for Health Research Clinical Research Network
NRES	National Research Ethics Service
OS	Overall Survival
PD	Progressive Disease
PFS	Progression Free Survival
PI	Principal Investigator
PISC	Participant Information Sheet and Consent Form
PR	Partial Response
PSF	Pharmacy Site File
QA	Quality Assurance
QC	Quality Control
R&D	Research & Development
RCT	Randomised Controlled Trial
REC	Research Ethics Committee
RECIST	Response Evaluation Criteria In Solid Tumours
RSI	Reference Safety Information
RSO	Research Support Office
SAE	Serious Adverse Event
SAR	Serious Adverse Reaction
SD	Stable Disease
SDV	Source Data Verification
SmPC	Summary of Product Characteristics
SOP	Standard Operating Procedure
SSAR	Suspected Serious Adverse Reaction
SUSAR	Suspected Unexpected Serious Adverse Reaction
TAB	Tumour activity and burden level
TMF	Trial Master File
TMG	Trial Management Group
ToR	Terms of Reference
TSC	Trial Steering Committee
TTP	Time to Progression
UAR	Unexpected Adverse Reaction
VAF	Variant Allele Frequency
WOCBP	Women of child bearing potential

2. PROTOCOL OVERVIEW

Full Title:	Circulating tumour DNA guided Adaptive BRAF and MEK Inhibitor therapy
Acronym:	DyNAMIC
Phase:	II
Target Population:	Patients in the United Kingdom, aged 18 years and over, with stage III unresectable, or IV, cutaneous melanoma with <i>BRAF</i> mutation and mutant ctDNA levels of BRAF VAF ≥ 7 copies/ml of plasma.
Sample size:	40 participants.
Trial Arms	ARM A (Standard of Care): Continuous dosing of encorafenib 450mg once daily plus binimetinib 45mg twice daily. ARM B (Adaptive Therapy): 4 weeks of encorafenib 450mg once daily plus binimetinib 45mg twice daily, followed by adaptive cycles based on ctDNA TAB level.
Inclusion Criteria:	At Screening 1. Written and informed consent obtained from participant and agreement of participant to comply with the requirements of the study 2. Histological confirmation of cutaneous melanoma including acral 3. ≥ 18 years of age 4. Stage III un-resectable/ IV disease 5. Measurable disease on CT (thorax, abdomen and pelvis, $\pm$ neck if indicated) and/or PET-CT, or clinically accurately measurable (RECIST v1.1) 6. <i>BRAF</i> p.V600E/K/R/D mutation confirmed (exact point^a mutation must be known) 7. ECOG performance status 0/1/2 8. Prior radiotherapy or radiosurgery must have been completed at least 2 weeks prior to the first dose of study drugs 9. Adequate organ function enabling safe administration of the study treatment

	10. Women of childbearing potential participating in the study (WOCBP see Appendix B for definition) must have a negative serum or urine pregnancy test (minimum sensitivity 25 IU/L or equivalent units of HCG) within 72 hours prior to the start of study drug. 11. WOCBP must agree to follow instructions for method(s) of contraception for the duration of treatment with study drugs plus at least 28 days following last dose of drug (either encorafenib or binimetinib), (see Appendix B). 12. Males who are sexually active with WOCBP must agree to follow instructions for method(s) of contraception for the duration of treatment plus 90 days (duration of sperm turnover) from last dose of drug (either encorafenib or binimetinib), (see Appendix B). Prior to randomisation: 13. BRAF ctDNA TAB level of ≥ 7 copies/ml of plasma. (If there is a discrepancy between the screening and baseline 1 ctDNA TAB levels this should be discussed via the DyNAMIC trial team at the LCTC.) 14. Left Ventricular Ejection fraction (LVEF) $\geq 50\%$ or $\geq \text{LLN}^b$ by ECHO a. The 3-letter abbreviation is also acceptable, (see Appendix F) b. LLN = lower limit of normal
Exclusion Criteria:	1. Prior systemic targeted BRAF/MEKi therapy for stage IV (metastatic) melanoma (treatment for stage III allowed as long as RFS ≥ 26 weeks following discontinuation of drugs) 2. BRAF wild-type malignant melanoma 3. Current evidence of active metastasis to the brain or leptomeninges (the designation of “active metastasis” should be determined by the local clinical team in discussion with the DyNAMIC study team at LCTC if needed and informed by recent imaging.) 4. Any contraindication to treatment with encorafenib or binimetinib as per the local Summary of Product Characteristics 5. Hypersensitivity to the active substance or to any of the excipients of encorafenib or binimetinib 6. Current use of a prohibited medication as described in Section 8.98.9 7. History of another malignancy. Exception: Patients who have been disease-free for 3 years, (i.e. patients with second malignancies that are indolent or definitively treated at least 3 years ago), curatively treated in situ cancer of the cervix, ductal carcinoma in situ (DCIS);

	stage 1, grade I endometrial carcinoma, or patients with a history of completely resected non-melanoma skin cancer. No additional therapy should be required whilst the patient is on study. Discussion via the DyNAMIC trial team at the LCTC as to whether they can be included in the study is allowed and the outcome of the discussion will determine entry. 8. Any serious or unstable pre-existing medical conditions (aside from malignancy exceptions specified above), psychiatric disorders, or other conditions that could interfere with the patient's safety, obtaining informed consent, or compliance with study procedures 9. Known Human Immunodeficiency Virus (HIV), Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV) infection 10. Females who are pregnant or breast-feeding and are not able to stop breast-feeding prior to first dose of study drugs (see section 7.57.5) 11. Prisoners or patients who are involuntarily incarcerated 12. Patients who are compulsorily detained for treatment of either a psychiatric or physical (e.g., infectious disease) illness	
Study Centres and Distribution:	11-15 Secondary / Tertiary NHS Hospital sites across the UK	
Participant Study Duration:	Duration of treatment: All participants will be treated until progressive disease or clinical deterioration following disease progression, or unacceptable toxicity or withdrawal of patient consent, at which point their care will be managed as per standard of care. Duration of follow-up: A minimum follow-up of 26 weeks for all participants.	
Study Duration:	30 months recruitment followed by an additional 6 month continued treatment/follow-up.	
	encorafenib	
	Form:	Capsule
	Dose:	450mg once daily

Investigational Medicinal Products (all arms)	Route:	Oral
	binimetinib	
	Form:	Tablet
	Dose:	45 mg twice daily
	Route:	Oral
	Objectives:	
Primary:	1. To assess whether tumours respond to re-introduction of encorafenib plus binimetinib following the first “drug off” period, assessed by % decrease of tumour activity and burden level (TAB; mutant copies of ctDNA/ml of plasma).	
Secondary:	1. To optimise the thresholds of percentage reduction in TAB level in ctDNA as a measure of response to stop drugs and the percentage increase in TAB as a decision to restart drugs (see section 5.15.4 for further details) 2. To assess maximal response (complete response (CR)/partial response (PR)/stable disease (SD)/progressive disease (PD)) to therapy in Arm A vs. Arm B 3. To assess progression free survival (PFS) in Arm A vs. Arm B 4. To assess time to clinical deterioration in Arm A vs. Arm B. This can either be at the point of progression (defined as per RECIST v1.1) or in participants treated beyond RECIST v1.1 progression, at the time of clinically determined deterioration defined at the discretion of the treating clinician, specifically due to the underlying cancer (see Section 8.28.2 for more information) 5. To assess overall survival 6. To assess melanoma specific overall survival in Arm A vs. Arm B 7. To assess number of adaptive therapy cycles completed by participants on Arm B 8. To assess median duration of adaptive therapy cycles completed by participants on Arm B	

	9. To assess whether ctDNA result can be provided within 5 working days from sample receipt into NBC 10. To assess quality of life in Arm A vs. Arm B 11. To assess toxicity (all grade adverse events) in Arm A vs. Arm B 12. To assess the number and location of sites of disease progression in Arm A vs. Arm B
Exploratory/ Translational:	<u>Exploratory objectives</u> 1. To describe maximal reduction in TAB level for each adaptive cycle 2. To describe rise in TAB levels during drug off period for each patient 3. To examine clonal evolution during continuous vs. adaptive drug dosing 4. To explore the relationship between observing a rise in ctDNA levels of mutant <i>BRAF</i> copies/ml and progressive disease observed scheduled scan results in Arm A <u>Translational work</u> Samples will be collected for research work including tumour biopsies and blood. Any translational studies using DyNAMIC samples must be authorised by the Sponsor. Due to the hypothesis-generating design of translational studies, these will not be performed to GCP standard however, they will be conducted with scientific rigour in accordance with principles of respect for persons, beneficence and justice. Translational work aims will include but are not limited to: • Understanding of tumour evolution on targeted therapy which will include in depth genetic and epigenetic analysis of ctDNA • Understanding of response and resistance to therapy which can include analysis of DNA, RNA metabolites and proteins • Characterisation of stage IV/stage III unresectable melanoma which can include analysis of DNA, RNA metabolites and proteins Any leftover samples will be stored for up to 25 years in sponsor approved laboratories and will be available for research work authorised by the Sponsor.

3. ROLES AND RESPONSIBILITIES

3.1 Sponsor

The Christie NHS Foundation Trust is the sponsoring organisation and is legally responsible for the study. They will centrally manage the budget and allocate funding to collaborating organisations and formally delegate specific sponsoring roles to the Chief Investigator (CI) and Liverpool Clinical Trials Centre (LCTC).

3.2 Funder

This study is funded by The Jon Moulton Charity Trust, a registered charitable organisation, via a £1.25 million grant to sponsor; the study will be run as a non-commercial investigator led clinical trial.

The funder has reviewed the proposal in line with their funding policies and will receive updates on progress as required by the terms and conditions of funding.

3.3 The Study Team

Chief Investigator (CI):

Professor Paul Lorigan is the CI and is an employee of The Christie NHS Foundation Trust and is responsible for overall design and conduct of the trial in collaboration with other members of the study team.

Principal Investigators (PIs):

In each participating centre, a Principal Investigator (PI) will be identified to be responsible for identification, recruitment, data collection and completion of electronic case report forms (eCRFs), along with follow-up of study participants and adherence to study protocol at site. They will also be responsible for safety reporting and processing any applicable safety information.

Clinical Trials Unit:

LCTC at the University of Liverpool (UoL), in collaboration with the CI, will have overall management responsibility and will be responsible for trial management activities including (but not limited to) study planning, Trial Master File management, safety reporting, data management, randomisations, statistical analysis, coordination of participating sites. LCTC will manage the budget allocated to UoL.

3.4 Oversight Committees

The DyNAMIC trial is subject to oversight from the following committees:

Trial Management Group (TMG):

A Trial Management Group (TMG) will be formed comprising the CI, other lead Investigators (clinical and non-clinical), members of the LCTC and representation from the Sponsor, as documented in the TMG Terms of Reference (ToR). The TMG will operate under the terms of their specific ToR and is responsible for monitoring all aspects of the progress and conduct of the trial and will be responsible for the day-to-day running and management of the trial. The TMG will meet at least monthly at setup stage and then reduce at

a reduced frequency throughout the trial unless more frequent meetings are required. See section 3.4 for further information.

Trial Steering Committee (TSC):

The Trial Steering Committee (TSC) will consist of an independent chairperson, at least 2 independent experts, one must be in the field of melanoma and the other in ctDNA, an independent biostatistician and will include the CI, selected PIs and observers as documented in the TSC ToR. The TSC will also include at least one lay patient representative member. The role of the TSC is to provide overall supervision for the trial and provide advice through its independent Chairman. The decision for the continuation of the trial lies with the TSC.

The TSC will operate under the terms of their specific ToR and members must disclose any conflicts of interest prior to any contribution. The TSC will meet at least annually, but as agreed in the TSC ToR. See section 3.4 for further information.

Independent Data and Safety Monitoring Committee (IDSMC):

The Independent Data and Safety Monitoring Committee (IDSMC) will consist of an independent chairperson, plus an independent member with clinical expertise in melanoma and ctDNA, and an independent biostatistician.

The IDSMC will be responsible for reviewing and assessing recruitment, interim monitoring of safety and effectiveness, trial conduct and external data. The IDSMC will first convene before the first participant is recruited and will then define frequency of subsequent meetings (at least annually). Details of the interim analysis and monitoring are provided in sections 6.4 and [12.312.3](#) respectively.

An IDSMC Charter, membership and member declarations will be drafted and agreed by the IDSMC in the first meeting. The IDSMC will provide a recommendation to the TSC concerning the continuation of the study. See section 3.4 for further information.

Molecular Tumour Board (MTB):

The Molecular Tumour Board (MTB) will consist of (but is not limited to) the CI, one other clinician, a statistician, a representative from CRUK NBC and a mathematician.

The MTB will discuss each threshold in addition to a general overview of the data and kinetics of ctDNA levels. Features of any participants who have progressed clinically on scan without a corresponding rise in TAB level will also be discussed.

An MTB Charter, membership and member declarations will be drafted and agreed by the MTB in the first meeting. The MTB will immediately adjust the threshold levels as required and update the TSC retrospectively.

3.5 Protocol Contributors

Appendix C lists in alphabetical order (by surname) those who have substantively contributed clinically and scientifically to the design of the protocol.

4 INTRODUCTION

4.1 Background

Melanoma:

Melanoma has been increasing in incidence over the last two decades due to a number of environmental and social factors as well as increased surveillance (1). It is the 4th most common cancer in the United Kingdom with 13,348 new cases diagnosed in 2011 (2). The World Health Organisation (WHO) reported in 2012 approximately 130,000 new diagnoses worldwide, with 37,000 deaths (3). More recent data from Cancer Research UK (CRUK) report 2333 melanoma-related deaths between 2016-2018, accounting for 1% of total cancer deaths (4).

The majority of patients are diagnosed with stage I/II melanoma, however 9% will be diagnosed with stage III/IV disease (5). Whilst only a very small percentage of patients with stage I disease will relapse, a further 10-25% of patients with stage II disease will progress to stage IV disease (6). Although survival has significantly improved with the paradigm-shift of treatment type from chemotherapy to immune and targeted therapies, approximately 40-50% of patients with stage III unresectable or stage IV disease who receive treatment will not survive to 5 years (7). New strategies are therefore required to improve both progression free and overall survival for this group of patients.

Current standard treatment of stage IV melanoma:

With the growing understanding of genetic alterations in melanoma and immune modulation by checkpoint molecules, two new therapeutic approaches have become standard therapy in late-stage melanoma, with both strategies improving overall survival (OS) in randomised trials.

Immune therapy:

One of the major mechanisms cancer cells use to avoid T cell mediated death is by up-regulation of immune checkpoints. In late stage melanoma, ipilimumab, a monoclonal antibody targeting CTLA-4, was the first checkpoint inhibitor to achieve improved OS in two randomised trials (8,9). The identification of the PD-L1/PD-1 axis as important for T cell responses to cancer cells (10), led to two therapies gaining FDA and NICE approval in melanoma: pembrolizumab and nivolumab. The Phase III Checkmate 066 trial randomised 418 treatment naïve, BRAF wild-type patients to receive either nivolumab (3mg/kg) or dacarbazine (11). It showed an objective response rate (ORR) of 40.0% (95% CI, 33.3 to 47.0) in the nivolumab group versus 13.9% (95% CI, 9.5 to 19.4) in the dacarbazine group (11). Median progression free survival (PFS) was 5.1 months in the nivolumab group versus 2.2 months in the dacarbazine group (HR=0.43; 95% CI, 0.34 to 0.56; P<0.001) (11). The 1 year OS rate was 72.9% (95% CI, 65.5 to 78.9) in the nivolumab group, as compared with 42.1% (95% CI, 33.0 to 50.9) in the dacarbazine group (HR 0.42; 99.79% CI, 0.25 to 0.73; P<0.001) (11). Similar results were obtained in a Phase III trial of pembrolizumab with a response rate of 32.9% vs. 11.9% for ipilimumab and 1 year survival rate of 68.4% for pembrolizumab vs. 58.2% for ipilimumab (HR 0.69; 95% CI 0.52 to 0.90) (12).

Combination therapy results in increased depth of response compared to single agents. The Checkmate 067 study randomised 945 previously untreated patients to combination nivolumab and ipilimumab (N+I) vs. ipilimumab alone or nivolumab vs. ipilimumab (13). The study was not powered to detect a difference between N+I and nivolumab however descriptive analyses have been subsequently performed and showed N+I to have better efficacy especially in those patients with poor prognostic features (>3 sites disease, high LDH, M1c/d disease) at baseline (14). The overall response rate (ORR) was 45% (95% CI, 39-50) in the nivolumab group, 58% (95% CI, 53-64) in the N+I group, and 19% (95% CI, 15-24) in the ipilimumab group with time to objective response 2.8 months for all groups (13,15). At 5 years, median PFS was 11.5 months (95% CI, 8.7-19.3) with N+I, compared to 2.9 months (95% CI, 2.8 to 3.2) with ipilimumab (HR for death or disease progression, 0.42; 99.5% CI, 0.35 to 0.51; $P<0.001$), and 6.9 months (95% CI, 4.3 to 9.5) with nivolumab (HR compared to ipilimumab, 0.53; 99.5% CI, 0.44-0.64; $P<0.001$) (15). Overall survival at 5 years was 52% in the nivolumab-plus-ipilimumab group and 44% in the nivolumab group, as compared with 26% in the ipilimumab group (15). The median OS was more than 60.0 months (median not reached; 95% CI, 38.2 to not reached) in the N+I group, 36.9 months (95% CI, 28.2 to 58.7) in the nivolumab group, and 19.9 months (95% CI, 16.8 to 24.6) in the ipilimumab group (15). The HR of N+I vs. ipilimumab was 0.54 (0.44-0.67) and HR of nivolumab vs. ipilimumab was 0.65 (0.53-0.79) (15). A recent update with median follow-up of 6.5 years confirmed the ongoing benefits of both combination ipilimumab + nivolumab, with no new safety signals (16). The best outcomes were seen in patients with small volume disease, normal LDH and good performance status.

More recently, the RELATIVITY Phase III trial randomised 714 previously untreated patients with stage III unresectable or stage IV melanoma to nivolumab or nivolumab plus relatlimab; a LAG-3 blocking antibody (17). At a median follow-up of 13.2 months, the median PFS in the nivolumab plus relatlimab group was 10.1 months vs. 4.6 months for nivolumab (HR 0.75; 95% CI 0.6–0.9, $P=0.0055$). PFS rates at 12 months were 47.7% (95% CI, 41.8–53.2) for nivolumab plus relatlimab vs. 36.0% (95% CI, 30.5–41.6) for nivolumab. Patients benefited from the combination irrespective of LAG-3 expression and therefore it is unlikely that it will be a clinically useful predictive biomarker. Incidence of grade 3/4 side effects was 18.9% for N+R vs. 9.7% for nivolumab with treatment discontinuation rates 14.6% vs. 6.7% respectively. Thus, in indirect comparisons with N+I, this combination is considerably better tolerated and is likely to become standard of care first line immune therapy at least in patients with lower risk disease.

Targeted therapy:

Activation of the mitogen-activated protein kinase (MAPK) pathway is crucial to the progression of melanoma (18). Approximately 35-45% of patients diagnosed with melanoma harbour a mutation in *BRAF*, which results in increased kinase activity and downstream phosphorylation of MEK and ERK (19,20). Due to this reliance of *BRAF* mutant melanomas on the MAPK pathway for their continued survival and proliferation, a number of treatments targeting elements crucial to its signalling have been developed (21–23). Initially, single agent *BRAF* inhibitors were reported to improve OS, resulting in a paradigm shift in melanoma management (21).

More recently, Phase III results combining *BRAF* and MEK inhibition have been reported (24–26). In the COMBI-D Phase III trial, ORR was 67% in the dabrafenib plus trametinib (D+T) group vs. 51% in the

dabrafenib/placebo group in 423 randomised patients (24). After 301 events, median PFS was 11 months (95% CI 8.0-13.9) in the D+T arm vs. 8.8 months (95% CI 5.9–9.3) in the dabrafenib only group (HR 0.67, 95% CI 0.53-0.84; $P=0.0004$) (27). The COLUMBUS Phase III trial randomised patients 1:1:1 to encorafenib plus binimetinib (E+B), encorafenib or vemurafenib. ORR by central review was 63% (56-70) in the E+B group vs. 51% (43-58) in the encorafenib group vs. 77 (40%; 33-48) in the vemurafenib group (26). At a median follow-up of 16.6 months (95% CI 14.8-16.9), median PFS was 14.9 months (95% CI 11.0-18.5) in the encorafenib plus binimetinib group, and 7.3 months (5.6-8.2) in the vemurafenib group (HR 0.54, 95% CI 0.41-0.71; $P<0.0001$) (26). Updated results at 48.8 months revealed median OS was 33.6 months (95% CI, 24.4-39.2) for E+B, 23.5 months (95% CI, 19.6-33.6) for encorafenib and 16.9 months for vemurafenib (95% CI, 14.0-24.5) (28). E+B vs. vemurafenib decreased the relative risk of death by 39% (HR, 0.61; 95% CI, 0.48–0.79) (28). Further follow-up at a median of 70.4 months presented at ASCO 2021, reported the median OS (95% CI) and 5-year OS rate with E+B were 33.6 (95% CI 24.4–39.2) months and 34.7% (95% CI 28.0–41.5), respectively (29). The 5-year PFS rate was 22.9% for E+B vs. 10.2% for vemurafenib; ORR (95% CI) was 64.1% (56.8–70.8) vs. 40.8% (33.8–48.2); and the median duration of response (DOR) was 18.6 in the E+B group vs. 12.3 months in the vemurafenib group (29). No new safety concerns were reported.

The Food and Drug Administration (FDA) approved vemurafenib/cobimetinib (V+C), D+T and encorafenib/binimetinib (E+B) for use in patients with metastatic *BRAF* V600E/K/R/D mutated melanoma on the basis of these phase III trials, however the National Institute for Clinical Excellence has not authorised V+C for use in the UK (21,23,25,26,30,31).

Newer agents such as next generation paradox-breaking *BRAF* inhibitors (e.g. PLX8394 (32)), which prevent *RAF* dimerization and *ERK* inhibitors (33), which target further down the *MAPK* pathway could potentially overcome some of the resistance mechanisms to *BRAF* and *MEK* inhibition, but have yet to be tested beyond Phase I/II trials.

Targeted vs. immune therapy and scheduling of therapy lines:

Whilst there have been no direct comparisons of targeted therapy and immunotherapy, a number of indirect comparisons have been carried out (7,34), and these show a clear benefit in terms of long term survival for immunotherapy over targeted therapy in the metastatic setting. The results of the DREAMseq (Doublet, Randomized Evaluation in Advanced Melanoma Sequencing) trial examining dabrafenib and trametinib followed by ipilimumab and nivolumab or ipilimumab and nivolumab followed by dabrafenib and trametinib in patients with stage III-IV *BRAF* V600 melanoma (NCT02224781) are awaited. This trial switches patients on treatment progression rather than at response to either therapy.

An alternative approach is to switch patients when they are responding to targeted therapy as this can result in changes to the tumour and the microenvironment that facilitates immune response. Pre-clinical data has revealed that *BRAF* inhibition results in an environment that can enhance immune responses (35,36). Tumours responding to *BRAF* inhibitors have been shown to have increased T cell infiltration, improved T cell recognition of melanoma associated antigens and reduced production of immunosuppressive cytokines (36–39). Microphthalmia-associated transcription factor (MITF) is up-regulated following *BRAF* inhibition,

resulting in increases in the expression of lineage antigens, such as gp100, melan-A, and tyrosinase-related proteins 1 and 2 (39,40). PD-L1 expression has also been shown to increase following 10-14 days of BRAF inhibitor therapy, which could enhance the effect of PD-1 inhibitors (40,41).

The SEquential COMBo Immuno and Targeted therapy (SECOMBIT; NCT02631447) study and the Immunotherapy With Ipilimumab and Nivolumab Preceded or Not by a Targeted Therapy With Encorafenib and Binimetinib (EBIN; NCT03235245) studies are both assessing whether a pre-specified period of induction targeted therapy (12 weeks for EBIN and 8 weeks for SECOMBIT) results in improved survival (42,43). Early results from SECOMBIT showed promising results for the induction arm of the trial (42). However, these strategies do not personalise treatment to the individual response of a patient and are at risk (particularly with 12 weeks of therapy) of patients already developing resistant clones which are more immune suppressive due to the length of the induction period. The CACTUS (Circulating Tumour DNA gUided therapy Switch; NCT03808441) trial in *BRAF* mutant patients, uses circulating tumour DNA (ctDNA) to determine when patients are responding to targeted therapy in order to personalise the timing of the switch. All of these trials allow restart of targeted therapy upon progression on immune therapy, therefore even if these strategies are shown to be effective, optimising targeted therapy scheduling remains critical for these patients.


Combining targeted and immune therapy

Studies examining combination targeted and immune therapy have been largely negative with no significant PFS difference seen for combination of spartalizumab, dabrafenib and trametinib (D+T) vs. D+T in a phase III trial (44). Only a small benefit was seen for PFS as per investigator assessment from 10.6 months (95% CI 9.3–12.7) in the control group (vemurafenib plus cobimetinib) to 15.1 months (11.4–18.4) in the triplet group (atezolizumab, vemurafenib plus cobimetinib; HR 0.78; 95% CI 0.63–0.97; log-rank $p=0.025$ which was not confirmed by independent assessment (45). Neither trial compared triplet therapy with anti-PD-1 single agent or combination N+I which would be more in line with standard of care in this group of patients. Sub-group analysis of these trials is ongoing and may provide a signal as to which patients could benefit from this strategy; however, this would need confirming in further randomised trials.

Adaptive therapy:

Adaptive therapy relies on the competition between drug-sensitive and drug-resistant subclones to control overall tumour growth (Figure 1) (46,47). It aims to stabilise tumour burden by allowing a significant population of treatment sensitive cells to survive, which suppress proliferation of the less fit, resistant populations (46).

Figure 1. Conventional sequential therapy of two treatments selects for a clone resistant to treatment one (red) upon tumour progression and subsequently a clone resistant to treatment two (green). B. Adaptive therapy maintains a stable tumour volume by introducing treatment holidays. Drug sensitive clones (blue) suppress the growth of less fit resistant clones (red). Adapted from West et al 2020.



Research has shown a reduction in drug resistant clones when targeted therapies are withdrawn. Mathematical analyses of tumour evolution have shown that continuous dosing selects for drug resistant clones, whilst an adaptive approach can control these (48,49). A number of models have been developed which have considered the proportion of resistant cells at start of therapy, the spatial constraints of tumours, the phenotypic switch of resistant cells with a tendency to result in slower growth rates, cell turnover rate, and velocity of evolution (47,49–55). A framework for considering how cancer evolves has been proposed based on 4 major components; the diversity of a cancer and how this changes over time (Evolutionary index) and the hazards impacting on clonal survival and resources available to neoplastic cells (Ecological index) (56). Adaptive therapy disrupts both the evolutionary and ecological indices in order to slow tumour growth.

To support these models in patients, a study with colorectal cancer revealed that resistant *KRAS* mutant clones identified in circulating tumour DNA (ctDNA) declined upon withdrawal of EGFR-specific antibodies (57). In addition, an adaptive therapy strategy has recently been tested in a pilot study treating patients with abiraterone in prostate cancer until a > 50% decline in their baseline levels of PSA pre-abiraterone was observed (46). Upon achieving this decline, abiraterone therapy was suspended and only restarted when the PSA increased to the baseline level (46). Preliminary data showed that all of the patients achieved PSA control when abiraterone was restarted and there was a signal that progression free survival (PFS) was improved compared to historical controls (46).

In melanoma, a number of models have shown that adaptive therapy could delay progression compared to continuous dosing. One study examined the differences in transcriptional states of melanoma cells *in vitro* using single cell analysis at baseline, when treated with vemurafenib and when drug was removed (58). They showed there was heterogeneity of these states, which mainly associated with levels of MITF, AXL and c-JUN, which have previously been associated with melanoma invasive/proliferative states (58–61). Different phenotypic states were associated with different growth kinetics and therefore they developed a two-compartment mathematical (ODE: ordinary differential equation) model to describe the competition between the sensitive and resistant states. The model was used to develop a personalised adaptive schedule, which was then tested in mice. This revealed that the adaptive therapy schedule delayed tumour progression in comparison with continuous or fixed intermittent schedules (58). The same group has subsequently examined the optimal timing of therapy and identified potential predictive factors as to which patients would likely benefit the most from an adaptive strategy (54). They compared two models; the first assumed genetically fixed drug-sensitive and resistant populations that competed for limited resource and the second considered phenotypic switching between drug-sensitive and resistant cells. They found that in both models adaptive therapy delayed time to progression by 6–25 months compared to continuous therapy. Factors predicting response to adaptive therapy included number of initial sensitive cells, competitive effect, switching rate from resistant to sensitive cells, and sensitive cell growth rate (54).

Further human-based data in melanoma supporting the potential benefit and safety of an adaptive strategy include a number of studies that have also shown that further responses can be seen to BRAF/MEK targeted therapy upon rechallenge following a drug holiday after progression or toxicity (62–65). An open-label phase II clinical trial showed that patients who had previously progressed on targeted therapy and had at least 12 weeks off treatment responded to rechallenge with 8/25 (32%) patients having a partial response and 10/25 (40%) having stable disease (63). Taken together, these studies suggest that melanoma resistance is plastic and can change following periods of treatment withdrawal.

Circulating tumour DNA (ctDNA):

An adaptive strategy requires precise and dynamic monitoring of changes in tumour activity and growth in order to be viable for use in the clinic. We have developed a number of assays based on circulating tumour DNA and have shown that it has potential to meet these criteria. Cell-free DNA (cfDNA) is DNA within the blood circulation, which is released from cells. Circulating tumour DNA (ctDNA) is DNA present in the bloodstream arising from cancer cells, which can be detected through identifying somatic mutations within it (66). Mandel and Métais first identified circulating nucleic acids in the blood stream in 1948, however it was not until 1994 that their potential utility as biomarkers for cancer detection and monitoring was realised (67). At that time, Sorenson *et al* showed the presence of mutated *KRAS* ctDNA sequences in the blood of patients with pancreatic cancer whose tumours also possessed mutated *KRAS* (68). The mechanisms behind the release of ctDNA are not completely understood, however it is thought it is produced by cell necrosis, apoptosis and secretion from macrophages that have phagocytosed cells (67,69). In addition circulating tumour cells (CTCs) can also release nucleic acids into the blood, which can be detected through a variety of methods (70).

Many studies have examined the role of ctDNA as a “liquid biopsy” which can predict disease progression, monitor treatment response and reflect tumour resistance. CtDNA has been shown to reflect disease status and tumour burden whilst on treatment for a variety of cancers (71–73). CtDNA levels correlated with treatment response when compared to computerised tomography (CT) imaging in patients with metastatic breast cancer on treatment (74). Levels increased on average 5 months (range 2-9) before progressive disease was seen on imaging (74). In melanoma, baseline pre-treatment level of mutant *BRAF* in cfDNA has been shown to be a predictive biomarker of duration of therapy in patients treated with the BRAF/MEK inhibitors. In a pooled analysis from four clinical trials enrolling patients with tissue confirmed *BRAF* V600E/K mutant melanoma, copies of mutant *BRAF* were identified in 76% (V600E) and 81% (V600K) of 732 baseline plasma samples using BEAMing (beads, emulsion, amplification). Its presence was associated with lower response rates compared to those patients where mutant *BRAF* was undetectable (75). Furthermore, patients who had detectable mutant *BRAF* had a significantly shorter PFS and OS compared to those with undetectable levels (76). This remained significant in multivariable analysis comparing baseline factors including LDH and performance status in 3 out of 4 of the studies (76). More recently, a further validation study of patients treated in the COMBI-d and COMBI-MB trials of dabrafenib plus trametinib supported baseline detectable ctDNA as a predictor of PFS and OS on targeted therapy (72). Using a cut-off of 64 copies of ctDNA per millilitre of plasma (determined using droplet digital PCR [ddPCR]) they stratified patients as high and low risk. Patients with low vs. high risk disease had a significantly longer PFS 12.7 vs. 6.5 months (HR 1.74; 95% CI 1.37–2.21, $p < 0.0001$) and 35.1 vs. 13.4 months (HR 2.23; 95% CI 1.73–2.87, $p < 0.0001$) (72).

There is also increasing evidence that ctDNA could be used to monitor patients on therapy, and that ctDNA changes may be detectable ahead of radiological or biochemical (LDH) change, giving an early indicator as to treatment efficacy or an early herald of disease progression. One study showed that progression in ctDNA was identified in 2 patients 86 and 120 days prior to radiological progression in patients treated with targeted therapy (77). A further study took longitudinal samples from 36 patients on targeted therapy, with 12 of these having detectable ctDNA levels prior to commencement of treatment (78). There was a significant decrease in ctDNA in all of the patients ($p < 0.01$), with median time to becoming undetectable ($n = 7$) or $< 1\%$ ($n = 5$) of 13 days (range 6–40 days) (78). An increase in the *BRAF* V600 mutant ctDNA fraction was detected prior to the clinical diagnosis of PD in 12 out of 27 (44 %) patients and simultaneously with PD in 7 out of 27 (26 %) patients (78). These data were subsequently confirmed in another study, which showed ctDNA levels increased before radiological progression by a mean of 110 days (79).

Although one study did not support the use of ctDNA as an on-treatment biomarker of response to targeted therapy (78), larger studies have shown that it could be a useful tool (72,80). One study investigated whether ctDNA levels of 10 copies per ml of plasma or higher in the first plasma sample since treatment initiation were predictive of PFS. They found patients with undetectable ctDNA had a median PFS of 9 vs. 4 months in patients with detectable ctDNA (HR, 4.05; 95% CI, 1.56 to 10.53). A large analysis of 224 patients from the COMBI-d trial, compared ctDNA levels in paired samples pre-treatment and at 4 weeks on dabrafenib plus trametinib (72). From the patients with paired samples, 201 (90%) had detectable ctDNA at baseline, and

121 (60%) still had detectable ctDNA following 4 weeks of treatment (72). All patients with conversion to undetectable levels had disease control (complete response, partial response, or stable disease) as their best overall response, compared with 103 (87%) of 118 patients with detectable ctDNA at week 4 (proportional odds likelihood ratio test for association $P=0.0002$) (72). In patients with high LDH only (above upper limit of normal), undetectable ctDNA at week 4 was significantly associated with PFS (HR 1.99; 95% CI 1.08–3.64, $p=0.027$) and OS (HR 2.38; 95% CI 1.24–4.54, $p=0.0089$) (72). Conversion to undetectable ctDNA at week 4 was independently associated with both PFS and OS in a Cox regression model that included clinical prognostic factors such as performance status and 3 or more organ sites with metastases (72). Taken together, these studies suggest that decreasing ctDNA levels on treatment are predictive of response to targeted therapy.

In addition, ctDNA can be used to detect mechanisms of resistance to therapy. In their seminal study, Murtaza *et al* demonstrated the proof of principle that circulating-free DNA (cfDNA) extracted from the plasma of patients over a course of treatment could be used to detect new mutations conferring resistance to therapy (71). Using whole exome sequencing (WES) of ctDNA, they demonstrated the emergence of an activating mutation in *PIK3CA* following treatment with paclitaxel in a patient with breast cancer and an *EGFR* T790M mutation on disease progression in a patient with lung cancer on gefitinib (71). In melanoma, studies have shown the development of mutations associated with resistance to targeted therapy, including *NRAS* and *PIK3CA* mutations (81,82). Thus, ctDNA can be used to not only assess tumour activity, but also clonal evolution during therapy, which can inform further models and future treatment strategies.

4.2 Rationale

Limitations of current treatments:

Despite the advances in melanoma management, patients with stage IV melanoma still die from their disease, with a 5 year survival of 39-70% depending on prognostic features such as number of disease sites, brain/liver/bone metastasis and raised lactate dehydrogenase (6). Combination BRAF and MEK inhibitor therapy has resulted in improved survival for patients (26). However resistance develops after a median of 12-15 months for the majority (24,26). Recently, combination anti-PD-1, BRAF and MEK inhibitors have been shown to be disappointing with COMBI-I not meeting its primary PFS endpoint and only a marginal improvement against targeted therapy (rather than immune therapy typically given as standard of care in the first line) seen for combination atezolizumab, vemurafenib and cobimetinib, which was not significant by independent committee review (45). New strategies are therefore required to prevent resistance occurring.


Adaptive therapy vs. intermittent therapy:

Preclinical studies in melanoma have shown a cell intrinsic fitness disadvantage when drug is withdrawn from targeted therapy resistant cells (83–85). The mechanism underlying this dependency of melanoma cells on the same therapeutic drugs to which they have acquired resistance or “drug addiction” phenotype is due to

ERK2 hyperactivation, transcriptional re-programming, DNA damage and parthanatos-related cell death (84–86).

Based on these data, a number of clinical trials have examined alternative dosing strategies for targeted therapy in melanoma. Firstly, the INTERIM study (a randomised phase II feasibility study of INTERmittent versus continuous dosing of oral targeted combination therapy in patients with *BRAF* V600 mutant stage 3 unresectable or metastatic Melanoma NCT03352947) uses a fixed intermittent dosing compared to standard dosing of dabrafenib plus trametinib. Patients are treated with dabrafenib (150mg twice daily) on days 1 – 21 of a 28 day cycle and trametinib (2mg once daily), on days 1 – 14 of a 28 day cycle. The SWOG S1320 randomized phase II trial of intermittent versus continuous dosing of dabrafenib and trametinib in BRAFV600E/K mutant melanoma, compared intermittent dosing 5 weeks on, 3 weeks off schedule with continuous dosing after an initial 8-week continuous run-in of therapy. Continuous dosing was shown to have a statistically significant progression free survival compared with intermittent dosing (median 9.0 months versus 5.5 months, $P = 0.064$, pre-specified two-sided $\alpha = 0.2$) (87). Although results are awaited for INTERIM and the scheduling is different, SWOG S1320 does not suggest that fixed intermittent dosing is a strategy to pursue further into Phase III studies. One of the possible reasons for this lack of efficacy may be due to the pharmacokinetics of trametinib in particular, in humans vs. mice/*in vitro*. As the half-life of trametinib in humans is approximately 10 days (86), the potential pharmacodynamic impact of sudden drug removal seen *in vitro* and in mice where the half-life is much shorter, cannot be reproduced in humans. In addition, a fixed schedule does not account for intra-patient tumour heterogeneity, with individual patients experiencing different tumour kinetics. Therefore, a personalised approach may be a better strategy.

Figure 2. A. In vivo experiment treating WM164 cells with continuous vs. fixed intermittent dosing (2 weeks on 1 week off) vs adaptive PLX4720 treatment. **B.** Tumour volume at experiment end shows significant improvement with adaptive therapy vs continuous or fixed intermittent.



As described above, adaptive therapy can personalise therapy to the kinetics of tumour growth and development of resistant clones vs. sensitive clones within each individual patient. When compared against continuous and fixed intermittent dosing *in vivo* a study showed a significant improvement in tumour control with an adaptive strategy (Figure 2) (58). This was better than a fixed intermittent/metronomic schedule modelling SWOG S1320 or INTERIM, which was not significantly different from continuous therapy.

Taken together, these data show that an adaptive therapy strategy can be used to overcome resistance to encorafenib plus binimetinib and in preclinical models of melanoma is superior to continuous or fixed intermittent dosing.

ctDNA as a measure of tumour activity and burden:

A dynamic marker of response and increasing tumour activity is required for an effective adaptive strategy. As discussed above, we have developed techniques using ctDNA in early and late stage melanoma, which are able to accurately monitor tumour burden and activity over time (88–91). By following tumour-specific mutations in the blood, we were able to show responses to treatment and detect disease progression prior to radiological relapse (91). We argue that ctDNA is not only a measure of tumour burden, but also tumour activity. This is supported by a study which analysed ctDNA and compared levels to tumour metabolic activity measured using FDG-PET in 52 patients who received systemic therapy for metastatic melanoma (92). Levels of ctDNA correlated with metabolic tumour activity seen on FDG-PET and early changes in ctDNA and metabolic disease burden were important indicators of treatment response (92). Thus, in DyNAMIC we refer to the tumour activity and burden level (TAB) level, which is equal to the number of mutant *BRAF* copies per millilitre (ml) of plasma.

ctDNA as an accurate biomarker of response dynamics vs. other biomarkers:

The “Adaptive BRAF-MEK inhibitor therapy for advanced BRAF mutant melanoma” trial (NCT03543969) is using participant's LDH levels, along with scans to decide when to hold or resume vemurafenib plus cobimetinib. However, we and others have shown that ctDNA is a better biomarker of tumour activity/burden than LDH (Figure 3) (81). Supporting our data, a small study of 6 patients with advanced melanoma treated with both targeted and immune therapies showed that longitudinal monitoring using ddPCR to detect BRAF/NRAS mutations in plasma was more accurate than LDH (93). A decrease in ctDNA levels corresponded with disease response (as early as one week into treatment) and a subsequent increase in levels corresponded with radiological progression (93). Similarly, another study compared ctDNA vs. LDH in identifying disease progression within 15 days of confirmed radiological




Figure 3. Progression in ctDNA is seen prior to progression on scan or rise in LDH

progression in 26 patients. In 82% of patients ctDNA progression was detected vs. only 40% having an LDH rise, with a median difference in sensitivity of 42% (95% confidence interval, 27%–58%; $p < 0.001$ (77). Basing decisions to treat on ctDNA rather than LDH also enables us to include a greater number of patients

as patients with normal LDH can still have detectable ctDNA. We therefore believe that using ctDNA enables more accurate monitoring of the disease, which increases the safety of an adaptive strategy.

Use of encorafenib plus binimetinib vs. other targeted therapy combinations:

There are currently two combination targeted therapy options available within the United Kingdom, dabrafenib plus trametinib and encorafenib plus binimetinib. Encorafenib has been shown to have increased potency and dissociation half-life, which translated into a superior overall response rate (51% vs. 40%) compared to vemurafenib another BRAF inhibitor (26,94). However, there have been no head-to-head randomised comparisons of D+T vs E+B. In their respective Phase III registration studies, median PFS was 11 months (95% CI 8.0-13.9) for D+T whilst for E+B it was 14.9 months (95% CI 11.0-18.5), but this could just be reflective of a different patient population. Encorafenib is an ATP-competitive BRAF inhibitor that suppresses the MAPK pathway in *BRAF* mutant tumour cells (V600E/V600D/V600K mutations), with a more than 10-times longer dissociation half-life (>30 hours) than either dabrafenib or vemurafenib, which enables sustained inhibition of the kinase (26). These preclinical data suggest that the encorafenib binimetinib combination will therefore enable more rapid control of sensitive clones when drug is resumed, making it a better therapy for an adaptive approach.

In addition, E+B and D+T have different toxicity profiles with D+T associated with pyrexia, skin rash and diarrhoea, whereas E+B is associated with nausea, vomiting and diarrhoea (28,95). One of the issues of the temperatures associated with dabrafenib (due to its metabolite) is that it can lead to multiple interruptions of therapy, which although has not been shown to impact outcomes in terms of survival, may cause issues with the logistics of an adaptive strategy, therefore we believe E+B may be a better option for this approach.

4.3 Risk and Benefits

Potential Risks:

This trial is categorised as Type B (somewhat higher than the risk of standard medical care) as per the risk-adapted approach to clinical trials adopted by the Medicines and Healthcare Products Regulatory Agency (MHRA).

Disadvantages for participants include the extra visits and blood tests that will need to be made, side effects of the medication, CT scans and radioactive tracer used in PET-CT scans. They will be fully informed of these in the participant information sheet. The main risk however is that E+B does not control the cancer upon restarting drugs or that the ctDNA does not identify the cancer becoming active. Our extensive preclinical data and experience in the CACTUS trial does not suggest that identification of the cancer becoming more active will be a problem as ctDNA is very accurate in identifying disease progression. The only scenario where ctDNA has been shown to be less effective in identifying disease progression is in the context of brain metastases, especially in isolated brain progression (77,96,97). This is likely due to the blood brain barrier reducing the variant allele frequency (VAF) of ctDNA in the blood. We have defined inclusion/exclusion criteria which only permits patients without current evidence brain metastases to enroll. We will monitor for

development of brain metastasis throughout the study with regular scans. In addition, we plan to reassess outcomes and response to restart of drug has occurred following the first drug holiday for every 6 participants on Arm B of the study. This will enable us to assess the trajectory of the ctDNA copies/ml of plasma of each individual participant and the cohort to determine whether drug off or drug on thresholds should change or whether there are any safety signals regarding tumour control.

While all exposure to radiation carries a very low risk that it may itself cause cancer in the future, the majority of scans on this study would also be performed as part of standard treatment.

As with all medications, E+B both carry risks of side effects. The participant should be encouraged to familiarize themselves with the patient information leaflet that will be provided for each drug.

More information regarding management of risks associated with this trial will be detailed and assessed in a separate Risk Assessment, which will be reviewed and agreed by the CI and Sponsor as part of the overall trial set-up process.

Potential Benefits:

There are two main potential benefits for participants on the study. The first is that based on the preclinical findings discussed above, they may have a longer time before they would require their next treatment line with delayed clinical progression (whilst taking drug) using an adaptive strategy. This may also result in improved overall survival. In addition, an adaptive strategy may result in improved quality of life and reduction in side effects from treatment, especially during time off drug.

4.4 Trial Hypothesis, Objectives and Outcomes

Research hypothesis:

- Participants will have ctDNA response upon re-introduction of encorafenib plus binimetinib following first drug holiday.
- Encorafenib plus binimetinib therapy using adaptive scheduling based on ctDNA level will improve PFS, reduce toxicity and enhance quality of life compared to continuous dosing in participants with stage III unresectable or stage IV melanoma.

Objectives and Outcomes

Objectives	Outcome Measures	Timepoint(s) of evaluation
Efficacy:		
Primary: To assess whether tumours respond to re-introduction of encorafenib plus binimetinib following the first “drug off” period, assessed by % decrease of tumour activity and burden level (TAB; mutant copies of ctDNA/ml of plasma).	Maximal reduction in percentage ctDNA mutant <i>BRAF</i> copies/ml of plasma from baseline 2 TAB level upon restart of E+B following first drug off period.	Longitudinally throughout the study
Secondary: To optimise the thresholds of percentage reduction in TAB level in ctDNA as a measure of response to stop drugs and the percentage and/or minimum increase in TAB as a decision to restart drugs (see section 5.1 for further details).	ctDNA mutant <i>BRAF</i> copies/ml of plasma	Longitudinally throughout the study
Secondary: To assess maximal response (complete response (CR)/partial response (PR)/stable disease (SD)/progressive disease (PD)) to therapy in Arm A vs. Arm B	Maximal response (complete response (CR)/partial response (PR)/stable disease (SD)/progressive disease (PD)) using RECIST v1.1 criteria	Baseline 1 until RECIST progression/clinical deterioration

Objectives	Outcome Measures	Timepoint(s) of evaluation
Secondary: To assess progression free survival (PFS) in Arm A vs. Arm B	Percentage of participants who have experienced disease progression or death. PFS defined as time from randomisation to progression (RECIST v1.1 and in Arm B defined as progression whilst on targeted therapy unless stopped due to toxicity/choice).	Continuously throughout the study
Secondary: To assess time to clinical deterioration in Arm A vs Arm B	Time to clinical deterioration. Defined as either the point of progression (defined as per RECIST v1.1) or in participants treated beyond progression, at the time of clinically determined deterioration defined at the discretion of the treating clinician, specifically due to progression of the underlying cancer (see section 8.2 for more information)	Continuously throughout the study
Secondary: To assess overall survival in Arm A vs. Arm B	Overall survival (OS) defined as time from randomisation until death from any cause	Continuously throughout the study
To assess melanoma specific overall survival in Arm A vs. Arm B	Overall survival (OS) defined as time from randomisation until death from melanoma	Continuously throughout the study
Secondary: To assess number of adaptive therapy cycles completed by participants on Arm B	Number of adaptive therapy cycles completed	Longitudinally throughout the study
Secondary: To assess median duration of adaptive therapy cycles on Arm B	Median duration of adaptive therapy cycles	Longitudinally throughout the study

Objectives	Outcome Measures	Timepoint(s) of evaluation
Secondary: To assess whether ctDNA result can be provided within 5 working days from sample receipt into NBC	Percentage of ctDNA results reported within 5 days from sample receipt into NBC	Longitudinally throughout the study
Secondary: To assess quality of life in Arm A vs. Arm B	EORTC QLQ-C30 and PRO-CTCAE	Longitudinally throughout the study
Secondary: To assess toxicity (all grade adverse events) in Arm A vs. Arm B	Adverse Events and Serious Adverse Events defined by CTCAE version 5	Continuously throughout the study
Secondary: To assess the number and location of sites of disease progression in Arm A vs. Arm B	Number and location of sites of progression	At disease progression on imaging
Exploratory:		
To describe maximal reduction in TAB level for each adaptive cycle.	Outcome measure is undefined and is subject to change due to the exploratory nature of the objective.	Longitudinally throughout the study
To describe rise in TAB levels during drug off period for each participant.	ctDNA mutant <i>BRAF</i> copies/ml of plasma	Longitudinally throughout the study
To examine clonal evolution during continuous vs. adaptive drug dosing	Outcome measure is undefined and is subject to change due to the exploratory nature of the objective.	Longitudinally throughout the study
To explore the relationship between observing a rise in ctDNA levels of mutant <i>BRAF</i> copies/ml and progressive disease observed on scheduled scan in Arm A	Outcome measure is undefined and is subject to change due to the exploratory nature of the objective.	Longitudinally throughout the study

Translational research:

Samples will be collected for research work including tumour biopsies and blood. Any translational studies using DyNAM1c samples must be authorised by the Sponsor. Due to the hypothesis-generating design of translational studies these will not be performed to GCP standard, however, will be conducted with scientific rigour in accordance with principles of respect for persons, beneficence and justice.

Translational work aims will include but are not limited to:

- Understanding of tumour evolution on targeted therapy which will include in depth genetic and epigenetic analysis of ctDNA
- Understanding of response and resistance to therapy which can include analysis of DNA, RNA metabolites and proteins
- Characterisation of stage IV/stage III unresectable melanoma which can include analysis of DNA, RNA, metabolites and proteins

Any leftover samples will be stored for up to 25 years in sponsor approved laboratories and will be available for research work authorised by the Sponsor.


5 STUDY DESIGN

DyNAMIC is designed as a randomised, parallel arm proof-of-concept study to:

1. Test our hypothesis that tumours respond (measured by ctDNA mutant *BRAF* TAB level) to re-introduction of targeted therapy (following drug holiday)
2. Test the thresholds we have simulated based on our pre-clinical data are correct or whether they need refining based on response in participants examined prospectively.

Participants with stage III un-resectable or stage IV *BRAF* mutant metastatic melanoma eligible for the study will be randomised 1:3 (Arm A [Standard of Care] vs Arm B [Adaptive Therapy]). All participants will commence encorafenib (E) 450mg once daily plus binimetinib (B) 45mg twice daily continuous dosing for 1 cycle of 28 days. A baseline 1 blood sample will be taken within 1 day prior to commencing E+B to test for the level of ctDNA (Baseline 1 TAB). At week 2 and week 4, further blood for ctDNA will be taken. All participants will have been randomised to Arm A or Arm B by Cycle 1 Day 14 of the initial E+B cycle. Participants will be stratified by ctDNA level at Baseline 1. Participants will be treated until disease progression (RECIST v1.1) or beyond RECIST v1.1 to clinical deterioration whilst taking E+B or stop due to participant choice/toxicity (See Section [8.28-2](#)).

Figure 4 Trial Schema:



The study will recruit 10 participants into Arm A and 30 into Arm B. As DyNAMlc is a pilot study, a balance is required to ensure the aims of the study are addressed in a timely way to move on to further studies powered for efficacy endpoints whilst providing sufficient data to ensure that there is certainty regarding the optimal thresholds and that a signal is seen to support the concept. The aims of the study are to demonstrate a reduction in tumour burden through assessing a drop in TAB upon restart of E+B. This will be demonstrated both within Arm B and by comparison against the Arm A.

Sample size justifications are based on demonstrating that the adaptive approach can continue to induce a reduction in tumour burden when drugs are re-started. The aim is to show that this occurs both solely within Arm B and by comparing the reduction against what is shown in Arm A (which at that timepoint should be stable due to ongoing response or increasing if resistance emerging).

For comparisons within Arm B only, the aim is to show that a reduction in TAB level given by a log fold change of 2 is observed. Analysis is based on estimation as opposed to demonstrating any pre-specified hypothesis. We expect the log-fold change in TAB level to have a standard deviation itself of 2 units. With 30 participants this will translate into 95% Confidence interval half-length of 0.72 [e.g. if a log fold change of 2 is observed it will have an approximate 95% CI of (1.28 – 2.72)] which is sufficient precision to demonstrate a meaningful TAB reduction. Attrition bias will be minimised by analysing data based on the principle of intention-to-treat (ITT).

5.1 Molecular tumour board ongoing assessment of TAB levels

The DyNAMlc trial has 2 key decisions repeated for each treatment cycle and break for Arm B:

- i. The TAB level to stop E+B
- ii. The TAB level to restart E+B

The criteria for the stopping and restarting of treatment for participants on Arm B are subject to change. The decision as to whether the participant should start or stop drug will be taken centrally using current thresholds, and the site will be informed immediately by email. The initial threshold criteria are provided below **and are for central use only**.

- During the first drug withdrawal, the TAB level will be compared against the **Baseline 1 TAB** level (see Figure 5). If the ctDNA shows a rise in TAB level of $\geq 10\%$ of **Baseline 1 TAB** or for participants with a **Baseline 1 TAB** of < 200 copies/ml, a rise by the equivalent of 15 copies/10ml (depending on the volume of plasma used and subject to the limit of quantitation) or there is disease progression (RECIST v1.1), the participant will restart E+B.
- Following the first period of drug withdrawal, upon restart of E+B, the TAB level will be compared to the **Baseline 2 TAB** level, which is the TAB level taken within 48 hours prior to E+B restart. When the TAB level falls to $\leq 50\%$ of **Baseline 2 TAB**, E+B are withdrawn.
- During the next period of drug withdrawal, E+B are restarted when the TAB level increases by $\geq 10\%$ of **Baseline 2 TAB**, or for participants with a **Baseline 1 TAB** of < 200 copies/ml, a rise by the equivalent of 15 copies/10ml (depending on the volume of plasma used and subject to the limit of

quantitation), or there is disease progression (RECIST v1.1). For each “on drug” targeted therapy cycle, when the TAB level falls to $\leq 50\%$ of **Baseline 2 TAB**, E+B are withdrawn.

- For each “drug off” period part of the adaptive therapy cycle E+B are restarted when the TAB level increases by $\geq 10\%$ of **Baseline 2 TAB**, or for participants with a **Baseline 1 TAB** of < 200 copies/ml, a rise by the equivalent of 15 copies/10ml (depending on the volume of plasma used and subject to the limit of quantitation), or there is disease progression (RECIST v1.1).
- In addition, if the participant is “drug off” period but there is disease progression (RECIST v1.1) drug will be restarted.

In all cases where the participant’s scan shows disease progression the site PI must notify the CI and the CI’s representative by email within 48 hours.

We have determined these levels through our preclinical data tracking ctDNA levels in participants treated with targeted therapy and through analysis of baseline values of participants with stage IV *BRAF* mutant melanoma screened for the CACTUS trial.

Throughout DyNAMIC, there will be ongoing monitoring of these thresholds by the MTB. In addition to ad hoc meetings that may be required, The MTB will meet after every 6th occurrence of recommenced therapy following the first period of drug withdrawal on Arm B and will analyse and discuss the TAB levels for each individual on Arm B and for the cohort as a whole. Any changes to the thresholds must be agreed by the MTB and will be documented in “Circulating Tumour DNA Threshold Determination and Ongoing Evaluation”. For safety reasons, changes to ctDNA thresholds for the stopping and restarting of treatment will be implemented immediately.

The MTB will discuss 5 main aspects including an overview of the radiological and clinical response data and kinetics of ctDNA levels. Features of any participants who have progressed on scan without a corresponding rise in TAB level will also be discussed.

- To assess what value the TAB rise during first drug withdrawal vs. Baseline 1 TAB (first point above) should be, given the data from participants on study.
- To assess what value the TAB rise vs. Baseline 2 TAB level (second point above) should be, given the data from participants on study.
- To assess what value of TAB decrease compared to Baseline 2 TAB should trigger a decision to withdraw drugs (third point above), given the data from participants on study.
- Whether these levels should be further adapted as participants complete more cycles of adaptive therapy or for different kinetics of TAB level. Rules may be further created based on mathematical modelling of participant data from DyNAMIC.
- To assess whether the frequency of ctDNA testing should be changed on a per participant basis


Figure 5: Adaptive treatment schedule on Arm B:

Figure 5: Adaptive treatment schedule on Arm B. A baseline TAB level (Baseline 1 TAB) will be taken within 24 hours of taking first dose of study drugs. Participants will initially receive E+B continuously for 4 weeks before E+B are withdrawn. Whilst off E+B, TAB level will be compared to Baseline 1 TAB and when criteria for drug restart are fulfilled, E+B will be restarted. Within 48 hours of the first drug restart, a further TAB level will be taken, termed Baseline 2 TAB level. Further adaptive cycles will compare the TAB level to the Baseline 2 TAB. When on E+B, if the criteria for drug withdrawal are fulfilled, E+B will be stopped. When off E+B the TAB level will be compared to Baseline 2 TAB and if the criteria for drug restart are fulfilled, E+B will be restarted.

To achieve one of the main objectives of the trial, which is to establish the optimal thresholds to stop and start drugs based on TAB level, DyNAMIC will include assessment of these levels at regular intervals for each individual participant. This will also provide additional safety monitoring. Thus, after every 6th occurrence of recommenced therapy following the first period of drug withdrawal on Arm B, the MTB will convene and will analyse and discuss the TAB levels for everyone on Arm B and for the cohort as a whole. Additional discussion regarding individual participants may occur outside of the meeting and any decisions e.g. to visit frequency will be documented. Any revisions to the thresholds will be documented in a new version of “Circulating Tumour DNA Threshold Determination and Ongoing Evaluation”.

5.2 Blinding

This is an open label study with no blinding requirements. All investigators and participants know which treatment is being administered.

5.3 Study Setting

Participants will be identified and recruited from approximately 11-15 hospitals in the UK selected for their expertise in the treatment of melanoma and a track record in conducting clinical trials. Each of the centres must have the required research nurses, pharmacy support and laboratory equipment to undertake the study.

Selection of Participating Sites:

Criteria for the selection of centres will be determined by the Trial Management Group and will be described in a separate document 'DyNAMIC Site Feasibility Questionnaire' maintained in the TMF.

Sites fulfilling the trial-specific criteria will be selected to be recruitment centres for the DyNAMIC trial and will be opened to recruitment upon successful completion of all global (e.g. REC and MHRA) and trial-specific conditions (e.g. site personnel training requirements) and once all necessary documents have been returned to the LCTC. Initiation of sites will be undertaken in compliance with LCTC internal processes. Conditions and documentation required will be detailed on an LCTC Green Light Checklist maintained in the TMF and must be fully completed prior to opening sites to recruitment.

Selection of Principal Investigators:

PIs will be required demonstrate equipoise, relevant experience, and commitment during early-stage feasibility assessment. All investigators will have the particular medical expertise necessary to conduct the study in accordance with this protocol and all regulatory and ethical requirements. Written agreement to conduct research as such will be obtained prior to site initiation. A suitable co-investigator should be identified at each site to deputise in case of PI absence.

6 STATISTICAL CONSIDERATIONS

6.1 Introduction

This section details the statistical principles relating to the design and conduct of the DyNAMIC trial. Please note that whilst an overview of the analytical techniques to be used are included here, a separate Statistical Analysis Plan (SAP) will be developed for evaluation of the primary and secondary outcomes. A separate SAP for exploratory work will also be produced.

6.2 Sample Size and Recruitment

Sample Size Calculation:

The primary endpoint is the maximal reduction in percentage mutant copies/ml from Baseline 2 TAB level upon restart of E+B following first drug off period. Whilst DyNAMIC contains a randomised component, success of the trial will depend on being able to demonstrate that at least 75% of participants randomised to Arm B can achieve a 50% reduction. From 30 participants, this success will be determined by ≥ 23 participants satisfying the endpoint criteria. Assuming a 'true' response rate of 85%, based on exact binomial probabilities, there is a 93% probability (Power) of observing this outcome. The probability of observing this result under varying assumptions of the 'true' success rates are provided in the table below.


'True' rate	60%	65%	70%	75%	80%	85%	90%
Probability (≥ 23 successes)	4%	12%	28%	51%	76%	93%	99%

Precision of QoL Estimates as a key secondary endpoint:

Given the sample size of 40 participants and the large number of quality of life domains that can be observed from the EORTC QLQ-C30 and PRO-CTCAE then power calculations based on pre-set hypotheses are not provided. It is anticipated that overall scores such as the global health status will be of greatest interest. Here, based on previous studies, a standard deviation of 12 units on the change in score between baseline 1 and follow-up is reasonable. From a sample size of 40 participants, it is reasonable to expect data suitable for analysis on 32 participants. Based on these assumptions then a standard error of 2.12 will be observed. This will result in a 95% confidence interval half -length of 4.16 being observed.

Feasibility of Sample Size:

Study recruitment is planned from approximately 11-15 sites which will recruit at an average recruitment rate of 1-3 participants/year/site. Assuming all sites are opened within 52 weeks, then a total of 2.5 years are required to obtain 40 participants.

Figure 7: Predicated recruitment curve:

6.3 Method of Randomisation

Allocation Sequence Generation:

Randomisation will be performed using a 1:3 allocation ratio. A pre-generated list produced on the principle of randomly permuted blocks will be used. Level of ctDNA will be included as the only stratification factor. It will have 3 levels: 7-50, 51-200 and >200 copies/ml of plasma. These are based on unpublished data from the CACTUS trial.

Concealment and Implementation of Allocation Sequence:

Participant allocations will be irrevocably generated upon completion of the web-based randomisation by a delegated member of the DyNAMIC Research Team at site. Allocation concealment will be ensured as the service will not release the randomisation code until the participant has been recruited into the trial; this takes place after confirmation of ctDNA levels.

6.4 Interim Analyses

Interim analyses to assess the ctDNA response to restart of E+B will occur after every 6th occurrence of recommenced therapy following the first period of drug withdrawal. The results will be discussed by the MTB, which has been specifically established to assess whether the levels of starting/stopping drugs determined by our preclinical data fit with our predictions. No formal stopping rules are included within the study design. Analyses of the accumulating data will also be performed at regular intervals (at least annually) for review by an IDSMC. These analyses will be performed at the LCTC. The IDSMC will be asked to give advice on whether the accumulated data from the trial, together with results from other relevant trials, justifies continuing recruitment of further participants or further follow-up. A decision to discontinue recruitment, in all participants

or in selected subgroups will be made only if the result is likely to convince a broad range of clinicians, including PIs, and the general clinical community.

6.5 Analysis Plan

Data Summaries:

Continuous data will be summarised as median (IQR) and categorical data will be presented as frequencies of counts with associated percentages. Summaries of the participant population will be reported by randomised group with Wilcoxon tests and Fisher tests (as appropriate) used to assess any differences.

Participant Groups for Analysis:

Analyses will be performed on an intention-to-treat principle, retaining all participants in the planned analyses irrespective of any protocol violations.

Levels of Significance:

As assessment of the primary outcome is based on the observed count of successes, no pre-defined level of significance is required to demonstrate a successful study. All results will be presented alongside nominal 95% two-sided confidence intervals and a p-value used to determine statistical significance.

Analysis of the Primary Endpoint:

The primary endpoint is the number of participants to achieve a 50% maximal reduction in percentage mutant copies/ml of plasma from Baseline 2 TAB level upon restart of E+B following first “drug off” period. This is measured as a binary endpoint and will be reported as a rate with associated 95% confidence intervals.

Further analyses will be performed on the TAB level over the course of the study. Here longitudinal regression techniques will be performed including random effects for the participant identifier. Details on the analytical approaches for each secondary endpoint are included in the statistical analysis plan (SAP).

7 ELIGIBILITY CRITERIA

The DyNAMlc trial aims to recruit 40 participants based on sample size calculations described in Section [6.26-2](#). All participants must provide written, informed consent before any study procedures occur (see Section [9.29-2](#)) and must meet all eligibility criteria as described below before being randomised.

7.1 Inclusion Criteria at Screening

Patients eligible for the DyNAMlc trial must comply with all of the following inclusion criteria at Screening:

1. Written and informed consent obtained from participant and agreement of participant to comply with the requirements of the study
2. Histological confirmation of cutaneous melanoma including acral.
3. ≥ 18 years of age
4. Stage III un-resectable/ IV disease
5. Measurable disease on CT (thorax, abdomen and pelvis, $\pm$ neck if indicated) and/or PET-CT, or clinically accurately measurable (RECIST v1.1)
6. *BRAF* p.V600E/K/R/D mutation confirmed (exact point^a mutation must be known)
7. ECOG performance status 0/1/2
8. Prior radiotherapy or radiosurgery must have been completed at least 2 weeks prior to the first dose of study drugs.
9. Adequate organ function enabling safe administration of the study treatments
10. Women of childbearing potential participating in the study (WOCBP see Appendix B for definition) must have a negative serum or urine pregnancy test (minimum sensitivity 25 IU/L or equivalent units of HCG) within 72 hours prior to the start of study drug.
11. WOCBP must agree to follow instructions for method(s) of contraception for the duration of treatment with study drugs plus at least 28 days following last dose of drug (either encorafenib or binimetinib), (see Appendix B).
12. Males who are sexually active with WOCBP must agree to follow instructions for method(s) of contraception for the duration of treatment plus 90 days (duration of sperm turnover) from last dose of drug (either encorafenib or binimetinib), (see Appendix B).

7.2 Inclusion Criteria prior to randomisation

To be eligible for the DyNAMlc trial, patients must also comply with the following inclusion criteria:

13. *BRAF* ctDNA TAB level of ≥ 7 copies/ml of plasma. (If there is a discrepancy between the screening and baseline 1 ctDNA TAB levels this should be discussed via the DyNAMlc trial team at the LCTC.)
14. Left Ventricular Ejection fraction (LVEF) $\geq 50\%$ or $\geq$ LLN^b by ECHO

- a) *The 3-letter abbreviation is also acceptable, (see Appendix F)*
- b) *LLN = lower limit of normal*

7.3 Exclusion Criteria

Any patient meeting any of the criteria listed below at Screening or Baseline 1 will be excluded from study participation:

1. Prior systemic targeted BRAF/MEKi therapy for stage IV (metastatic) melanoma (treatment for stage III allowed as long as RFS $\geq$ 26 weeks following discontinuation of drugs)
2. BRAF wild-type malignant melanoma
3. Current evidence of active metastasis to the brain or leptomeninges (the designation of “active metastasis” should be determined by the local clinical team in discussion with the DyNAMIC study team at LCTC if needed and informed by recent imaging.)
4. Any contraindication to treatment with encorafenib or binimetinib as per the local Summary of Product Characteristics
5. Hypersensitivity to the active substance or to any of the excipients of encorafenib or binimetinib
6. Current use of a prohibited medication as described in Section [8.98-9](#)
7. History of another malignancy. **Exception:** Patients who have been disease-free for 3 years, (i.e. patients with second malignancies that are indolent or definitively treated at least 3 years ago), curatively treated in situ cancer of the cervix, ductal carcinoma in situ (DCIS); stage 1, grade I endometrial carcinoma, or patients with a history of completely resected non-melanoma skin cancer. No additional therapy should be required whilst the patient is on study. Discussion via the DyNAMIC trial team at the LCTC as to whether they can be included in the study is allowed and the outcome of the discussion will determine entry.
8. Any serious or unstable pre-existing medical conditions (aside from malignancy exceptions specified above), psychiatric disorders, or other conditions that could interfere with the patient’s safety, obtaining informed consent, or compliance with study procedures
9. Known Human Immunodeficiency Virus (HIV), Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV) infection
10. Females who are pregnant or breast-feeding and are not able to stop breast-feeding prior to first dose of study drugs (see Section [7.57-5](#))
11. Prisoners or patients who are involuntarily incarcerated
12. Patients who are compulsorily detained for treatment of either a psychiatric or physical (e.g., infectious disease) illness

7.4 Co-enrolment Guidelines

Individuals who have participated in a CTIMP and received any IMP within 28 days or 5 half-lives (whichever is shorter) preceding screening will be ineligible for the DyNAMIC trial.

Those participating in DyNAMIC are allowed to consent and undergo screening phases within other CTIMPs, however they must have experienced disease progression before they are randomised and/or enter any treatment phase.

Co-enrolment to non-CTIMP observational studies is permitted.

Should participants wish to withdraw from DyNAMIC to take part in another CTIMP trial, provisions for follow-up should be discussed with the CI and LCTC.

7.5 Lifestyle Restrictions

Meals and Dietary Restrictions:

Participants are not allowed to consume St John's Wort as part of herbal remedies as it could affect the efficacy of trial drugs. Grapefruit juice is contraindicated. No other meal/dietary restrictions apply.

Effects on ability to drive and use machines:

Encorafenib and binimetinib have a minor influence on the ability to drive or use machines. Visual disturbances have been reported in some patients treated with encorafenib during clinical studies. Participants should be advised not to drive or use machines if they experience visual disturbances or any other adverse reactions that may affect their ability to drive and use machines.

Contraception:

Women of childbearing potential must use effective contraception during treatment and for at least 28 days following the last dose of trial drug (either encorafenib or binimetinib) (see Appendix B). Combination treatment may decrease the efficacy of hormonal contraceptives. Therefore, female participants using hormonal contraception must use an additional or alternative method such as a barrier method (e.g. condom) during treatment and for at least 28 days following the last dose of trial drug.

During treatment and for at least 90 days (duration of sperm turnover) following the last drug dose, males who are sexually active with WOCBP must use a male condom plus partner use of a suitable highly effective method of contraception (see Appendix B).

Pregnancy and Breastfeeding:

It is unknown whether encorafenib or binimetinib, or their metabolites are excreted in human milk. A risk to the newborns/infants cannot be excluded. For patients to be included in DyNAMIC, they must have given birth/stopped breastfeeding 5 days prior to commencing treatment or they will be withdrawn from the trial.

In addition to those required as part of the study procedures, additional pregnancy testing of WOCBP is to be undertaken if clinically indicated according to the investigator criteria. This applies to both treatment arms and is regardless of whether a participant on Arm B is on or off drug.

Fertility:

There are no data on the effects of encorafenib or binimetinib on fertility in humans. Based on findings in animals, the use of encorafenib may impact fertility in males of reproductive potential. As the clinical relevance of this is unknown, male participants should be informed of the potential risk for impaired spermatogenesis and local practice followed to address this. There are no animal data on the effect of binimetinib.

8 TRIAL TREATMENT / INTERVENTIONS

8.1 Introduction

Eligible participants will be randomised between Arm A and Arm B. Participants in Arm A will receive encorafenib 450mg once daily plus binimetinib 45mg twice daily continuous dosing and participants in Arm B will receive encorafenib 450mg once daily plus binimetinib 45mg twice daily adaptive dosing unless a dose reduction is required. If patients require a dose reduction from the start of treatment this should be discussed with the DyNAMIC study team at LCTC.

Arm B Reserve Pack:

DyNAMIC requires the use of “Reserve Packs” in Arm B, this is because participants will need to recommence treatment upon a phone call and therefore it may not be convenient for them to immediately attend hospital to collect study drug. Participants in Arm B need to be dispensed a new reserve pack:

- Prior to re-commencing treatment for the first time
- At each physical visit if the participants have taken any doses from the reserve pack
- If there has been a change to the dose of either encorafenib or binimetinib

The Schedule of Assessments (see Section 99) and Figure 6 flowchart outline the timepoints at which a reserve pack should be dispensed.

DyNAMIC is fully open label and encorafenib and binimetinib are commercially available and should be sourced locally as per standard practice at the investigator sites using products that have a UK or EU license. Both encorafenib and binimetinib are still under a patent and will therefore only be available from current manufacturer Pierre Fabre Limited. Investigator sites will be responsible for using the most recent Summary of Product Characteristics (SmPCs) for products being used at site. A copy of the current SmPC, or a Statement of Location that makes reference to the electronic source, must be maintained in the DyNAMIC Pharmacy Site File.

Descriptive information for the IMPs can be found in the respective package insert. IMPs will be used and stored as detailed on the product label and according to manufacturer’s instructions. There will be no reimbursement to sites for the treatment as this is standard of care.

8.2 Treatment Name / Description

Encorafenib and binimetinib will remain under patent for the duration of this trial, therefore only one brand of each, Braftovi (encorafenib) and Mektovi (binimetinib), are available for use.

Unless a dose reduction is required from the start (see section 8.1), all participants will commence encorafenib 450mg once daily plus binimetinib 45mg twice daily continuous dosing for 1 Cycle of 28 Days. A Baseline 1 blood sample will be taken within 24 hours prior to the initiation of Cycle 1 Day 1 E+B to test for the level of ctDNA (Baseline 1 TAB). Participants who meet all inclusion criteria will be randomised to Arm A or Arm B. Participants will be stratified by ctDNA level at baseline. At week 2 and week 4, ctDNA will be

taken. At week 4 the participant will commence the following depending on which arm they have been randomised to.

Arm A – Continuous encorafenib plus binimetinib targeted therapy

Treatment aligning to (as E+B classed as IMP) standard of care (SOC) continuous dosing of encorafenib 450mg once daily plus binimetinib 45mg twice daily unless a dose reduction is required. Both drugs are administered as part of a 28 day cycle. Participants will have ctDNA testing every 2 weeks and clinical review every 4 weeks with scans every 8 weeks (+/- 1 week). After 6 months patients have the option to reduce the frequency of visits to every 4 weeks. Participants will be treated until confirmed disease progression (RECIST v1.1) or if there remains clinical benefit following confirmed RECIST v1.1 progression, participants are allowed to continue treatment and undergo further scans until clinical deterioration. Clinical deterioration is defined at the discretion of the treating clinician, specifically due to the underlying cancer (examples might include significant increase in pain, increase in ascites or pleural effusions, malignant bowel obstruction or a general decline in ECOG performance status due to cancer progression). The treating clinician should contact the CI, or their representative, via the LCTC if there are any queries relating to defining this. In addition, participants will stop treatment if there is significant toxicity or if it is the participant's choice to stop.

Arm B - Adaptive encorafenib plus binimetinib targeted therapy:

Participants in Arm B are given treatment in cycles of 14 days which commences only after one SOC continuous cycle of 28 days E+B has been completed. Following the SOC cycle, participants will withhold E+B. If the patient has toxicity during the first 28 calendar days that requires treatment to be interrupted, the first drug off period will only commence once the ctDNA TAB is $\leq 50\%$ of baseline 1. This does not impact the timelines for randomisation. They will undergo 2 weekly ctDNA monitoring for TAB level, with results expected to be returned within 5 working days from sample receipt at CRUK NBC (Figure 6). After 6 months patients may reduce the frequency of ctDNA testing only following approval by the MTB and with ongoing monitoring by the trial team as to whether the reduced frequency should continue.

If the ctDNA shows a pre-specified rise in TAB level or there is disease progression (RECIST v1.1), the participant will restart E+B. Following the first period “off-drug”, within 48 hours prior to restarting E+B a new ctDNA sample must be taken. This will establish a new baseline TAB level for the rest of the adaptive cycles (Baseline 2 TAB). If a Baseline 2 is taken outside of this window, a decision as to whether the patient can continue on the study should be discussed by the MTB taking into account the ctDNA results to date.

Following the Baseline 2 timepoint, each time E+B are restarted following an “off-drug” period, is defined as the start of the next targeted adaptive therapy “on-drug” cycle. For each “on-drug” cycle, when the TAB level falls by a pre-specified percentage of Baseline 2 TAB, E+B are withdrawn (considered the start of the “off-drug” period) and they are restarted when the TAB level increases by a prespecified TAB level.

If the ctDNA shows a pre-specified rise in TAB level or there is disease progression (RECIST v1.1), the participant will restart E+B within 24 hours of the result being returned (unless the baseline 2 sample needs to be taken). These values are based on our mathematical modelling of participants undergoing longitudinal ctDNA monitoring and data from the CActUS trial.

Participants will undergo a clinical review every 4 weeks with scans every 8 weeks (+/- 1 week). Participants will be treated until confirmed disease progression (RECIST v1.1) whilst on targeted therapy unless stopped due to toxicity/choice, or if there remains clinical benefit following confirmed RECIST v1.1 progression, participants are allowed to continue treatment and undergo further scans until clinical deterioration. Clinical deterioration is defined at the discretion of the treating clinician, specifically due to the underlying cancer (examples might include significant increase in pain, increase in ascites or pleural effusions, malignant bowel obstruction or a general decline in ECOG performance status due to cancer progression). The treating clinician should contact the CI or their representative within 48 hours via the LCTC if there are any queries relating to defining this. In addition, participants will stop treatment if there is significant toxicity or if it is the participant's choice to stop or it is the end of trial, at which time it will be discussed as to whether they want to continue standard dosing E+B.

Reserve Pack

As participants on Arm B will need to restart treatment upon ctDNA and TAB level rises, a reserve pack needs to be dispensed to the participants when they attend for their Baseline 2 TAB blood test (see below and Section [99](#)).

If participants have a dose limiting toxicity they will undergo dose reduction as per standard of care (see SmPC and Section [8.58.5](#)). If they require withdrawal of drug due to toxicity, this is allowed for a maximum of 6 weeks. If study drug is withdrawn due to toxicity for more than 6 weeks, the participant is to be withdrawn from treatment.

Drug on and off periods will be proactively tracked by manual review. In time, this may be supported by real time visualisations generated by the ACUITY system, managed by NBC Digital Cancer Research Team.

ctDNA samples will continue to be taken at the same 2-weekly schedule regardless of whether the participant is on/off-drug. If the participant has had to stop drug due to toxicity the ctDNA will only be acted upon when they have safely re-started drug as per the treating clinician's judgement. The schedule continues until there is disease progression (RECIST v1.1) when the participant is taking E+B (or clinical deterioration if the participant continues to receive adaptive E+B treatment beyond progression), significant toxicity, participant choice or end of study.

8.3 Manufacturing and Distribution

Regardless of its status as a standard of care treatment, E+B administered within Arm A of DyNAMIC will be classed as IMPs due to the same products also being an experimental regimen within Arm B.

Arm B in this study will be classed as IMP. All IMP will be procured via the local supply chain at each participating hospital site. This study will fall under the exemption to Regulation 37 in Statutory Instrument 2004/1031. Meaning an exemption from the requirement to hold a manufacturing authorisation for investigational medicinal products (IMPs). The regulation specifically applies to “assembly” carried out in a hospital or health centre by a doctor, a pharmacist or a person acting under the supervision of a pharmacist.

Labelling:

The MHRA has approved the justification from sponsor for a Type B trial with adaptations, as a result there is no requirement for Annex 13 labelling.

There is also no requirement for Annex 13 labelling on the reserve pack. To prevent accidental use, reserve packs need to be labelled as per the directions contained in the site dispensing section of the pharmacy manual.

Supply:

E+B will be sourced from routine hospital stock and their handling and management will be subject to standard procedures of the pharmacy and the respective SmPC.

Storage and dispensing:

Descriptive information for E+B can be found in the patient information leaflet within each package. E+B should be stored in accordance with the environmental conditions (temperature, light, and humidity) as determined by the manufacturer. If concerns regarding the quality or appearance of a particular drug arises, the drug should not be dispensed and the manufacturer informed immediately.

There will be no re-imburement to sites for E+B.

E+B should be dispensed according to the standards at each site. E+B for Arm A will be dispensed in cycles of 28 days with Arm B dispensed in cycles of 14 days. Participants on Arm B who have experienced toxicity during the first cycle that resulted in an interruption to treatment may need to delay their first drug off period. If this occurs, the extended treatment should be dispensed in 14 day cycles.


Participants on Arm B will receive a further reserve pack supply of medication, to be used if immediate recommencement of treatment is required following a ctDNA blood test. The reserve pack is to be replaced following its use, or if the dose is reduced (see Figure 6).

If sites are unable to dispense cycles in line with these protocol requirements, then this must be discussed via the DyNAMIC trial team at the LCTC and alternative local arrangements agreed and documented.

All documentation related to the storage and dispensation of E+B must be maintained in the DyNAMIC Pharmacy Site File, or a Statement of Location provided signposting the current document location.

Please refer to the relevant SmPC for further information.

Figure 6: Pharmacy Preferred Dispensing Diagram:



8.4 Preparation and Administration

Preparation and administration of E+B will be according to the SmPC.

Note: If the participant vomits after the administration of either drug, dose(s) should not be re-taken and the next dose(s) taken as scheduled.

8.5 Treatment Modifications

After the participant has been successfully registered to DyNAMIC, the clinician is free to give alternative treatment / intervention to that specified in the protocol, at any stage, if they feel it to be in the best interest of the participant. However, the reason for doing so should be recorded and the participant will remain within DyNAMIC for the purpose of follow-up and data analysis according to the treatment option to which they have been allocated. Similarly, the participant remains free to withdraw at any time from the protocol treatment and follow-up without giving reasons and without prejudicing further treatment.

If treatment-related toxicities occur on either arm, they are to be managed as per local protocols for standard of care and in accordance with the dose modifications guidance within the current SmPC. Both treatments should be simultaneously dose reduced, interrupted or discontinued when possible. Refer to the current SmPC for information on exceptions where dose reductions to one IMP only are necessary. If the participant requires withdrawal of drug due to toxicity, this is allowed for a maximum of 6 weeks. If study drug is withdrawn due to toxicity for more than 6 weeks, the continuation of the participant on treatment should be discussed with the DyNAMIC study team at the LCTC.

8.6 Accountability Procedures

All E+B dispensed must have a sufficient 'use by' date.

A record of individual prescriptions will be maintained as per local practice. Sites will not be required to keep detailed accountability records of batch numbers and expiry dates of the IMPs dispensed. There will be reconciliation between doses recorded as dispensed and doses returned, inclusive of reserve packs.

8.7 Assessment of Compliance

eCRFs will be used to capture information on non-adherence to trial treatment. Study-site personnel will establish information of instances of non-adherence from the participant. If non-adherence was not due to a clinical interruption of treatment, the participant should be counselled on the importance of the adherence to the treatment.

8.8 Product Recall

Participating site pharmacies are to have local policies in place for product recalls. If the need to recall either of the IMPs used in DyNAMIC is identified, investigators should inform the LCTC immediately.

The Sponsor receives email alerts from the MHRA. Should an alert regarding a product recall of either of the IMPs be received, it will be relayed to sites immediately.

8.9 Concomitant Medications/Treatments and Specific Restrictions

Medications Permitted:

- Supportive care for disease-related and drug-related symptoms (anti-emetics, anti-diarrhoeals, and analgesics) may be offered to all participants on DyNAMIC including blood and blood products
- Medications to manage co-morbidities, apart from those listed below, are permitted

Medications Not Permitted/Precautions Required:

The medications listed below are the main interactors with E+B. For further details regarding these and other drug interactions, please refer to the relevant current SmPC.

The following medications are **prohibited** within 28 days or 5 half-lives, whichever is shorter, prior to the start of the initial Cycle 1 Day 1 and during the study treatment period (unless to treat a drug-related adverse event):

- Any concurrent antineoplastic therapy (i.e., chemotherapy, hormonal therapy, immune therapy, non-palliative radiation therapy, or standard or investigational agents for treatment of cancer)
- Illicit drugs
- Antiretroviral drugs
- Herbal remedies (e.g., St. John's wort)

The following medications are **recommended to be stopped** within 28 days or 5 half-lives, whichever is shorter, prior to the start of the initial Cycle 1 Day 1 and during the study treatment period. If they must be continued, then dose reductions and/or additional safety monitoring should be performed as per standard of care:

- CYP3A4 inhibitors (encorafenib is primarily metabolised by CYP3A4). Strong CYP3A4 inhibitors should be avoided (due to increased encorafenib exposure and potential increase in toxicity). Examples of strong CYP3A4 inhibitors include, but are not limited to: ritonavir, itraconazole, clarithromycin, telithromycin, posaconazole and grapefruit juice.
- Strong CYP3A4 inducers (which may compromise the efficacy of encorafenib) should be avoided. Examples strong CYP3A4 inducers include, but are not limited to: carbamazepine, rifampicin, and phenytoin. Alternative agents with no, or moderate, CYP3A induction potential should be considered.
- UGT1A1 inducers (such as rifampicin and phenobarbital) and inhibitors (such as indinavir, atazanavir, sorafenib) should be co-administered with caution as binimetinib is primarily metabolised through UGT1A1 mediated glucuronidation.

The following medications should be **co-administered with caution**.

- OATP1B1, OATP1B3 and BCPR substrates (encorafenib is an inhibitor of these substrates) should

be used with caution. In vivo, these substrates can result in increased concentrations.

- Substrates of renal transporters OAT1, OAT3, OCT2, or hepatic transporters OCT1 and P-gp should be used with caution. In vitro, encorafenib potentially inhibits these transporters, and may increase exposure.

Data on Concomitant Medication:

The investigator must be informed as soon as possible about any medication taken from Screening until 30 days after the last dose of E+B.

Data on concomitant medication(s), including any dietary supplements, taken during the study will not be routinely collected, except during safety event reporting (where indicated). However, the participant will be asked at each visit if they are taking any new medications in order to identify any non-permitted medications.

8.10 Overdose

An overdose may be realised if informed by the participant, or via a check for compliance of treatment. An overdose for encorafenib is defined as the participant taking a dose of >450mg within 18 hours and for binimetinib a total dose of >90mg within 18 hours.

If an apparent overdose is observed, the site team must establish from the participant if extra drug has been taken, or whether the discrepancy is due to another reason e.g. drug misplaced.

If an overdose is observed, it must be recorded as an Adverse Event. Expedited reporting of overdose to LCTC is only required if the criteria for a Serious Adverse Event is met. Specific information on reporting adverse events can be found in Section [1144](#).

At doses of encorafenib between 600 to 800 mg once daily, renal dysfunction (Grade 3 hypercreatinemia) was observed in 3 out of 14 participants. The highest administered dose occurred as a dosing error in one participant who took encorafenib at a dose of 600 mg twice daily for 1 day (total dose 1200 mg). Adverse reactions reported by this participant were Grade 1 events of nausea, vomiting and blurred vision; all subsequently resolved. The highest dose of binimetinib evaluated as single agent in clinical studies was 80 mg administered orally twice daily and was associated with ocular (chorioretinopathy) and skin toxicities (dermatitis acneiform).

Management:

There is no specific treatment for overdose of either encorafenib or binimetinib.

If overdose occurs, the drugs should be interrupted and the participant should be treated supportively with appropriate monitoring as necessary.

Since both E+B are highly bound to plasma proteins, haemodialysis is likely to be ineffective in the treatment of overdose.

9 PARTICIPANT TIMELINES AND ASSESSMENTS

9.1 Identification and Screening

Potential participants will be identified from routine clinics, and their eligibility potential for DyNAMIC assessed by the clinical team responsible for their care. Only following review of the patient's hospital notes for eligibility will an appropriately qualified investigator or study nurse inform participants of the nature of the study and give relevant information, both verbally and in writing, about the objectives of the research, plus the benefits and risks.

Pseudonymised screening logs are to be completed for every patient considered as a potential participant for DyNAMIC, inclusive of those who are not registered as a participant. The screening log should ideally be updated in real time on a patient-by-patient basis. For monitoring purposes, all reasons for non-eligibility are to be recorded on the log. The screening log is maintained within the online DyNAMIC Clinical Study Database.

A separate participant identification log which contains participant direct identifiers should be maintained at site. The participant identification log must link the direct identifiers for participants with the Participant Number generated by the DyNAMIC Clinical Study Database. The participant identification log should be stored in a secure place. It will be required for monitoring visits but must never be sent outside of the site. Sites may use the DyNAMIC Participant Identification Log template or may use a local template.

The Participant Information Sheet & Consent Form (PISC) obtains permission from the patient for screening assessments to be conducted, including ctDNA testing, and must be signed before performing any protocol-related procedures that are not part of normal patient care (see Section [9.29.2](#)).

All potential participants will be assessed for their eligibility in a two-step assessment process. Patients will initially be assessed against the Screening Inclusion Criteria and Screening/Randomisation Exclusion Criteria (see Section [7.7](#)). Participants who meet all Screening Inclusion Criteria will be permitted to begin the initial 28 day cycle of E+B. Once all outstanding test data has been received to meet the requirements of the Randomisation Eligibility Criteria, the participant may be randomised to Arm A or B. Randomisation must be completed prior to Cycle 1 Day 14 of E+B.

To confirm eligibility, the Screening visit must be performed **no more than 42 days** prior to randomisation. See Section [9.49.4](#) for a list of trial assessments.

If the patient's disease is rapidly progressing, or if the patient wants to start treatment immediately, they can start treatment whilst waiting for the ECHO and ctDNA result. Blood for ctDNA must be obtained prior to commencing E+B, and the patient must be randomised within two weeks of commencing E+B. The result of the ECHO must be available prior to randomisation, see schedule of assessments tables for further information.

9.2 Informed Consent

Informed consent is a process initiated prior to an individual agreeing to participate in a trial and continues throughout the individual's participation. Written informed consent is required for all individuals participating in LCTC coordinated studies. The process should involve discussion between the patient and an individual knowledgeable about the research, the presentation of written material (e.g. information leaflet or consent document), and the opportunity for patients to ask questions and have these satisfactorily answered. In obtaining and documenting consent, the research team should comply with applicable regulatory requirements and should adhere to GCP and to the ethical principles that have their origin in the Declaration of Helsinki. Informed consent should be regularly reaffirmed throughout DyNAMIC and all discussions and consent should be documented appropriately in the patient's medical notes. If a patient does not want to provide consent they do not have to give a reason.

A delegated member of the research team who is trained in obtaining informed consent will need to discuss the information contained within the PISC and sufficiently answer any questions the patient may have. The consent discussion will therefore occur during a routine clinic visit; the entire discussion should last for about 30-60 minutes. Written informed consent shall be obtained only when the patient has had sufficient time to consider the information presented and all of their questions have been satisfactorily answered.

Consent should not be discussed with, or requested from the patient until their potential participation has been discussed by the clinical team responsible for their care, and their hospital notes checked for potential eligibility.

Essential trial information should be recorded in the patient's medical notes along with the date the informed consent was obtained. Informed consent must be obtained on a currently approved and localised version of the DyNAMIC PISC, and signed on the same day by both the patient and the delegated member of the research team who conducted the consent discussion.

Informed consent must be obtained before the performance of any protocol-related procedures that are not part of normal patient care, inclusive of screening assessments. Any assessments that have already been conducted as part of standard care may be used instead of repeating the screening tests but must fall within the screening period (up to 28 days prior to starting treatment).

Prospective Informed Consent Process:

Written informed consent will be sought from patients who will be approached by the site team and invited to consider participation.

A written information sheet that forms part of the ethically approved PISC will be provided. This includes a detailed explanation of the DyNAMIC trial and makes clear that the rights and welfare of the participants will be protected; it will be emphasised that consent may be declined or withdrawn at any time in the future without the quality of care being adversely affected. The research staff will facilitate verbal discussions about the research and the consent process, as well as providing answers to any questions that arise.

After verbal and written information has been provided, the investigator seeking consent will ensure that the patient has fully understood all the information and will ask if they are happy to consent to participation in the DyNAMlc trial.

Where this is the case, written informed consent will be obtained by means of a dated signature on the consent form. This should be countersigned and dated by the person who obtained informed consent i.e. the PI or other appropriately qualified member of the research team who has been delegated this responsibility.

The original signed document will be retained in the DyNAMlc Investigator Site File (ISF) and copies will be made:

- One copy provided to the participant for their information
- One copy transferred to the LCTC via a secure and encrypted method. The signed ICF (part of the PISC) must not be sent with any clinical data
- One copy filed in the participant's medical records paper/electronic

N.B. Details of the consent process (date, persons involved, version and type of information sheet and consent form used) must also be recorded directly into the participant's medical records.

Loss of capacity:

If a participant that has consented then becomes unable to give reaffirmed informed consent (as determined by the treating team), they will be immediately withdrawn from DyNAMlc, however previously obtained consent for the collection and use of existing data and translational samples remains valid.

Attempts to elicit information regarding safety events (AE, SAE etc.) that may have occurred to the participant since their incapacitation should be made to the participant's family, chaperone or legal representative etc.

Language Considerations:

If the participant does not have proficient English to provide informed consent, then a translator may be used to verbally translate the PISC according to local practice.

The quality-of-life questionnaires (EORTC QLQ-C30 and PRO-CTCAE) can be made available in a limited number of other languages and will be provided upon request by the LCTC

9.3 DyNAMlc Clinical Study Database

All participants who are considered for inclusion into DyNAMlc, will be recorded within the DyNAMlc Screening Log. The Screening Log is maintained via a secure, 24-hour accessible, web-based system known as the DyNAMlc Clinical Study Database. A unique Participant Number will be generated for every patient logged; this number will be maintained for the entire study duration.

It is the responsibility of the delegated pharmacy staff to ensure there is sufficient supply of study drugs.

A personal login username and password, provided by the LCTC, will be required to access the DyNAMlc Clinical Database System. Designated research staff will be issued with their personal login and password

upon completion of training in the use of the system. This training will be coordinated by the LCTC and an 'eCRF Completion Guidelines' document will be provided to site. These instructions will contain the web-link to the DyNAMIC Clinical Database System, provide step-by-step instructions for participant enrolment and randomisation, and completion of subsequent eCRFs.

System Failure:

Back-up systems/processes are in place to allow for the continuity of enrolment and randomisation in the event of internet failure at participating sites or main server failure at the University of Liverpool.

In the very unlikely event of a network failure at the University of Liverpool, participating sites should contact the DyNAMIC Trial Team at the LCTC to discuss options for the processing of an enrolment or randomisation. LCTC contact details are available from the 'LCTC Contact Details' document.

Back-up randomisation services are available Monday to Friday between 9:00 to 17:00 (excluding public or bank holidays, or University of Liverpool closed days, e.g. working days in between Christmas and New Year).

9.4 Assessments and Confirmation of Eligibility (Screening)

Eligibility can only be confirmed by an appropriately qualified medical professional who is named on the delegation log and must not occur until fully informed consent is documented. Eligibility criteria are described in detail in Section 7.7. Screening can occur multiple times if the condition of the patient changes whereby the investigator feels they might now be eligible. In addition, patients awaiting confirmation of a scan to show progression e.g. on immune therapy can also commence screening, including testing of the ctDNA as long as by the end of the screening period they fulfil all the screening criteria (i.e. have stopped other anti-cancer treatment and have a scan confirming measurable disease within the screening period).

Confirmation of eligibility (Screening) must be documented in the participant's medical notes. As a minimum, details must include who confirmed eligibility, when this was confirmed, and when the participant was formally entered into DyNAMIC (e.g. date of completion for assessment of Screening eligibility).

The following procedures will be performed at the participant's Screening visit:

- Obtain written informed consent prior to any trial specific procedures being conducted.
- Assessment and confirmation of Screening eligibility
- Medical history and physical examination, including height and weight, ECOG performance status and vital signs
- A complete list of all prior anti-cancer therapies will be recorded in the eCRF.
- Retrieve primary tumour block and other tissue from cancer related surgeries
- Blood sampling for the following tests[^]
 - ctDNA (streck tube - inc. extra for research purposes)

- Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH
- Is the participant taking prohibited concomitant medication? If so, these should be stopped and/or alternatives found.
 - Electrocardiograph (ECG)
 - Echocardiogram (ECHO) - standard safety monitoring of participants receiving E+B, not a research requirement – please refer to SmPC. ECHO must however be done as part of Screening and left ventricular function confirmed to be $\geq$ LLN prior to randomisation. If participant has limb disease, a PET-CT (with contrast if no known allergy to it) is to be performed as clinically indicated in place of the CT scan.
 - CT (with or without contrast if known allergy to it) of thorax, abdomen and pelvis (TAP), +/- neck if clinically indicated.
Alternatively, as per local practice if known allergy to contrast, CT without contrast thorax (+/- neck), plus MRI without contrast of abdomen and pelvis.
 - MRI of brain, CT is an acceptable alternative if required according to local practice (both with contrast if no known allergy to it).
Scans are to be reported according to RECIST v1.1. (See Appendix D). It is imperative that the same modality of imaging is performed throughout the participants time on DyNAMIC in order for accurate RECIST reporting and CT TAP (+/- neck) with CT/MRI brain is preferred.
 - Collect tissue from any standard of care indicated surgical resection.
 - Pregnancy test for WOCBP.
 - Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. Images are retained by site as source documents.

[^]Sample may be re-taken if required, e.g. original is lost or becomes unviable in transit, however all other procedures must still be completed within the stipulated timelines for Screening eligibility to be confirmed.

[#]Reporting as AST or ALT is to be consistent throughout the study on a per participant basis.

9.5 Baseline 1 Assessments

Baseline 1 refers to the assessments conducted prior to the commencement of the initial 28 day targeted cycle of E+B and is applicable to all participants.

The Screening and Baseline assessments required at both visits will not need to be duplicated and will only need to be performed once. The Baseline 1 visit ctDNA blood **MUST** be taken on the day of, or day before (- 1 day), commencing the initial target cycle of E+B. If Screening and Baseline visits are the same day, 8 x ctDNA tubes are to be taken.

Where the Screening and Baseline assessments are conducted on separate visits, safety bloods must be taken within 2 days prior to the baseline visit, and all other assessments except for ctDNA collection, must be completed within 28 days of commencing the initial 28 day targeted cycle of E+B.

Blood for ctDNA MUST be taken prior to commencing the initial 28 days targeted cycle of E+B. Failure to collect a ctDNA sample prior to initiating E+B may result in the participant being withdrawn from DyNAMIC and reverting to the local care pathway.

In order to accurately complete the Baseline eCRF and collect the necessary information for the trial analyses, assessments should be completed as detailed below. The following procedures will be performed at the participant's Baseline 1 visit:

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
 - Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH
 - CRP
- Is the participant taking prohibited concomitant medication? If so, these should be stopped and/or alternatives found.
- Physical examination including ECOG performance status and vital signs.
- Pregnancy test for WOCBP (MUST be conducted within 72-hours prior to commencing the initial targeted 28 day cycle of E+B).
- Additional blood for research (EDTA tube).
- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents).
- Dispense initial 28 day supply of E+B medication.
- Optional research biopsy (see Section [9.139-13](#)).
- Quality of life questionnaires (EORTC QLQ-C30 and PRO-CTCAE).

Routinely collected information e.g. medical history / vital signs / relevant blood test results etc. can be transcribed from the participant's medical notes into the eCRF once appropriate consent has been obtained.

The participant may be Randomised once all Screening and Baseline 1 assessments have been completed and Randomisation eligibility is confirmed (see Section [9.79-7](#) for details).

[#]Reporting as AST or ALT is to be consistent throughout the study on a per participant basis.

9.6 Commencement of Initial Targeted 28 day E+B Cycle

IMPORTANT: As noted in Section [9.19-4](#), if the participant is too unwell or wants to start targeted treatment as soon as possible, then the initial targeted 28 day cycle of E+B may commence following confirmation of all Screening Eligibility Criteria, and ctDNA blood has been collected at Baseline 1. Any outstanding Screening or Baseline 1 assessments must be completed within the timeframes stated in Sections 9.4 and 9.5. The participant should however be counselled that they may still not be eligible to continue participation in DyNAMIC and will not be randomised if they do not meet the Randomisation Eligibility Criteria.

9.7 Randomisation

The participant is not required to attend a Randomisation visit.

Participants must be randomised within 14 days of the cycle 1 day 1. All Screening assessments, including the ctDNA result, must be reported prior to randomisation. The week 2 ctDNA test will be compared to the ctDNA taken closest to when the participant commences targeted treatment i.e. Baseline 1.

Eligibility Criteria (see Section [7.27-2](#)) will be finalised upon: receipt of the screening ctDNA result from the CRUK NBC and reporting of an LVEF $\geq 50\%$ or $\geq \text{LLN}$ by ECHO. If there is discrepancy between the screening and Baseline 1 ctDNA TAB levels, this should be discussed via the DyNAMIC trial team at the LCTC. Should the participant satisfy all Screening Inclusion Criteria, the participant will be randomised.

Participants will be randomised to receive either Arm A (Continuous Standard of Care Therapy), or Arm B (Adaptive Therapy) in a ratio of 1:3 (Arm A: Arm B).

Should the ctDNA analysis not be possible for any reason, the participant will be deemed ineligible for randomisation, withdrawn from participation and will revert to the local care pathway.

Once randomisation is complete, an automated email confirmation will be sent to the PI and other relevant research team members at the participating site. Copies will also be sent to the DyNAMIC Trial Team mailbox and to the mailbox used for the management of blood kit supplies.

Following randomisation, participants should be notified of their allocation as soon as possible and then should have their treatment adapted as described in Section [9.89-8](#).

9.8 Intervention

Once the research team are made aware of the treatment arm allocated, the participants will have the treatment plan implemented. All patients will have a run-in of one complete standard of care cycle (i.e. 28 days) of E+B, then at that timepoint Arm A will continue on standard of care therapy and Arm B will withhold drug as described in Section [88](#).

9.9 Schedule for Assessments and Follow-up - Arm A

The assessments listed below are to be performed at the stipulated timepoints, and only for participants randomised to Arm A.

Note that some assessments are independent of the treatment cycle schedule and are required at calendar timepoints e.g. days or weeks, measured from the administration of E+B treatment at cycle 1, day 1.

If two timepoints occur concurrently, any assessments will only need to be completed once, e.g. collection of ctDNA blood.

Treatment Cycle 1, Day 15

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
 - Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH
- Is the participant taking any prohibited concomitant medication? If so, these should be stopped and/or alternatives found.
- Record and report any adverse events and/or toxicity assessments that have occurred since commencing initial treatment.
- Physical examination, including ECOG performance status and vital signs.
- Additional blood for research (EDTA tube).
- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents).
- Pregnancy test for WOCBP (only if clinically indicated).
- Report survival status.

[#]Reporting as AST or ALT is to be consistent throughout the study on a per participant basis.

Treatment Cycle 2 (Onwards), Day 1

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
 - Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH

- Is the participant taking prohibited concomitant medication? If so, these should be stopped and/or alternatives found.
- Record and report any adverse events and/or toxicity assessments that have occurred since commencing initial treatment.
- Physical examination, including weight, ECOG performance status and vital signs.
- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents).
- Pregnancy test for WOCBP (only if clinically indicated).
- Dispense supply of study medication.
- Quality of Life Assessments (Cycle 2 day 1 only)
- Compliance assessment – reconcile quantity of unused tablets.
- Report survival status.

#Reporting as AST or ALT is to be consistent throughout the study on a per participant basis.

Treatment Cycle 2 (Onwards), Day 15

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
- Optional research biopsy (Cycle 2 only) (see Section [9.139-13](#)).
- Pregnancy test for WOCBP (only if clinically indicated).
- Report survival status.

From 6 months post-randomisation, the day 15 visit is optional for Arm A participants

As Per Site Schedule

- Echocardiogram (ECHO) - standard safety monitoring of participants receiving E+B, not a research requirement – please refer to SmPC.
- Electrocardiograph - standard safety monitoring of participants receiving E+B, not a research requirement – please refer to SmPC.

Every 8 Weeks From Cycle 1, Day 1

- If participant has limb disease, a PET-CT (with contrast if no known allergy to it) can be performed as clinically indicated in place of the CT scan.
- CT (with or without contrast if known allergy to it) of thorax, abdomen and pelvis, +/- neck if clinically indicated.

Alternatively, as per local practice if known allergy to contrast, CT without contrast thorax (+/- neck), plus MRI without contrast of abdomen and pelvis.

(Scans to be reported according to RECIST v1.1 (See Appendix D). The same imaging modality, e.g. CT or PET-CT must be used throughout the study to enable RECIST comparison. Comparisons must not be made between PET and CT scans.)

Every 12 Weeks From Cycle 1, Day 1

- Quality of Life Assessments.

Every 26 Weeks From Cycle 1, Day 1

- MRI of brain, CT is an acceptable alternative if required according to local practice (both with contrast if no known allergy to it).

Scans to be reported according to RECIST v1.1 (See Appendix D).

Unscheduled visit

This visit should occur only if there is concern for clinical progression. It can occur multiple times however once the patient is confirmed to have progression with no clinical benefit they should not have any further unscheduled visits.

- Physical examination, including weight, ECOG performance status and vital signs.
- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents).

Discontinuation of Study Treatment

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
 - Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH
- Record and report any adverse events and/or toxicity assessments.
- ECOG performance status
- Weight
- Collect tissue from any standard of care indicated surgical resection
- Pregnancy test for WOCBP 28 days following the final administration of study drug.
- Additional blood for research (EDTA tube).
- Optional research biopsy (see Section [9.139-13](#)).
- Quality of Life Assessments

- Compliance assessment – reconcile quantity of unused tablets.
- Report survival status.

At disease progression (RECIST v1.1) and/or no further clinical benefit)

- Blood tests
 - ctDNA (streck tube – inc. extra for research purposes)
 - Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH
- Record and report any adverse events and/or toxicity assessments.
- Pregnancy test for WOCBP (if clinically indicated) 28 days following the final administration of study drug.
- Physical examination, including weight, ECOG performance status and vital signs.
- Collect tissue from any standard of care indicated surgical resection.
- Additional blood for research (EDTA tube).
- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents).
- Optional research biopsy (see Section [9.139-13](#)).
- Quality of Life Assessments.
- Compliance assessment – reconcile quantity of unused tablets.
- Report survival status.

[#]Reporting as AST or ALT is to be consistent throughout the study on a per participant basis.

Every 12 Weeks Following End of Study Treatment

- Record and report any adverse events and/or toxicity assessments.
- Collect tissue from any surgically indicated surgical resection.
- Record and report data on subsequent treatment(s), date(s) of treatment and any disease progression, and survival status.

End of Participation (excluding death)

- Pregnancy test for WOCBP 28 days following the end of *systemic* treatment (if not captured at End of Treatment or if clinically indicated). If participants are continuing to receive e+b as standard of care treatment following the end of their participation, the pregnancy test is to be deferred to the End of Study Participation visit.
- Record and report any adverse events and/or toxicity assessments.
- ECOG performance status.
- Collect tissue from any standard of care indicated surgical resection.
- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents).
- Compliance assessment – reconcile quantity of unused tablets with treatment regimen.
- Record and report data on any other known treatment(s), date(s) of treatment and any disease progression since the previous visit and survival status.

Arm A ctDNA Real-Time Clinical Decision Making:

As participants on Arm A will receive E+B as a regimen that is equivalent to standard of care, the result of the ctDNA TAB level will not impact the treatment received. Disclosure of the results of ctDNA TAB levels for participants on this Arm will be deferred.

Schedule of Assessments:

All assessments and follow-up are to be conducted in line with the Schedule of Assessment tables, below. The legend for the numerical caveats, indicated in superscript, is in Section [9.119.44](#).

ARM A	Pre-Treatment		Randomisation (no more than 14 days post C1 D1)	Continuous Treatment			As Per Site Schedule	Fixed Timelines			Post-treatment			End of Study Participation	
Study Procedures	Screening (up to 42 days prior to C1 D1) ¹¹	Baseline ¹¹		Cycle 1 D15	Cycle 2 onwards (Optional: omit day 15 from cycle 6 onwards)			Unscheduled visit (concern regarding progression)	Every 8 weeks from C1 D1	Every 12 weeks from C1 D1	Every 26 weeks from C1 D1	Discontinuation of study treatment	Disease Progression and/or no further clinical benefit ⁸		Follow-up every 12 weeks after end of study treatment
					D1	D15 ¹⁴									
Window	- 42d			+/- 4d				+/- 1w	+/- 2w	+/- 2w	+28 days	+28 days	+/- 1w	+/- 1w	
Eligibility check	x	x	x												
Written informed consent prior to any study procedure	x														
Medical history and demographics	x														
Review of prior anti-cancer therapies	x														
Retrieval of archival primary tumour block and any other tissue obtained through prior cancer-related surgery	x														
Haematology and Biochemistry	x	x		x	x						x	x			
Blood for ctDNA (streck tube) inc. additional blood for research	x	x ¹²		x	x	x					x	x			
Confirmation of screening BRAF ctDNA TAB level			x												
Prohibited medication review	x	x		x	x										
AE and toxicity assessment				x	x						x ⁹	x	x ⁹	x ⁹	

ARM A	Pre-Treatment		Randomisation (no more than 14 days post C1 D1) ¹¹	Continuous Treatment				As Per Site Schedule	Fixed Timelines			Post-treatment			End of Study Participation
Study Procedures	Screening (up to 42 days prior to C1 D1) ¹¹	Baseline ¹¹		Cycle 1 D15	Cycle 2 onwards (Optional: omit day 15 from cycle 6 onwards)		Unscheduled visit (concern regarding progression)		Every 8 weeks from C1 D1	Every 12 weeks from C1 D1	Every 26 weeks from C1 D1	Discontinuation of study treatment	Disease Progression and/or no further clinical benefit ⁸	Follow-up every 12 weeks after end of study treatment	
					D1	D15 ¹⁴									
Window	- 42d			+/- 4d					+/- 1w	+/- 2w	+/- 2w	+28 days	+28 days	+/- 1w	+/- 1w
ECOG	x	x		x	x		x					x	x		x
Weight	x				x		x					x	x		
Height	x														
ECG	x							x							
Echocardiogram	x							x							
Confirmation of ECHO results			x												
CT scan (thorax, abdomen pelvis (+/- neck if clinically indicated) or MRI. PET-CT as clinically indicated (NB imaging modality must be consistent)	x								x						
CT or MRI brain	x										x				
Further tissue collection from surgical resection (only if clinically indicated. See Section 9.139.13) ¹	x											x	x	x	x
Pregnancy test	x ⁶	x ⁴		x ⁶								x ⁷	x ⁶		x ⁷
Research blood sample (EDTA tube)		x		x								x	x		
Physical exam ² , vital signs and clinical disease assessment	x	x		x	x		x						x		
Encorafenib plus binimetinib dispensed		x			x										

ARM A	Pre-Treatment		Randomisation (no more than 14 days post C1 D1)	Continuous Treatment				As Per Site Schedule	Fixed Timelines			Post-treatment			End of Study Participation
Study Procedures	Screening (up to 42 days prior to C1 D1) ¹¹	Baseline ¹¹		Cycle 1 D15	Cycle 2 onwards (Optional: omit day 15 from cycle 6 onwards)		Unscheduled visit (concern regarding progression)		Every 8 weeks from C1 D1	Every 12 weeks from C1 D1	Every 26 weeks from C1 D1	Discontinuation of study treatment	Disease Progression and/or no further clinical benefit ⁸	Follow-up every 12 weeks after end of study treatment	
					D1	D15 ¹⁴									
Window	- 42d			+/- 4d					+/- 1w	+/- 2w	+/- 2w	+28 days	+28 days	+/- 1w	+/- 1w
Optional research biopsy		x ⁵				x ¹⁰						x	x		
Quality of Life Assessments (EORTC QLQ-C30 and PRO-CTCAE)		x			x ¹⁵					x		x	x		
Compliance assessment					x							x	x		x
Date of disease progression on each subsequent therapy														x	x
Subsequent treatment data														x	x
Survival status				x	x	x						x	x	x	x

9.10 Schedule for Assessments and Follow-up – Arm B

Treatment assessments

The assessments listed below are to be performed at the stipulated timepoints, and only for participants randomised to Arm B.

Note that some assessments are independent of the treatment cycle schedule and are required at calendar timepoints e.g. days or weeks measured from the administration of E+B cycle 1, day 1.

If two timepoints occur concurrently, any assessments will only need to be completed once, e.g. collection of ctDNA blood.

First Treatment cycle (Cycle 1, Day 15)

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
 - Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH
- Is the participant taking any prohibited concomitant medication? If so, these should be stopped and/or alternatives found.
- Record and report any adverse events and/or toxicity assessments that have occurred since commencing initial treatment.
- Physical examination, including ECOG performance status and vital signs.
- Pregnancy test for WOCBP (only if clinically indicated).
- Additional blood for research (EDTA tube).
- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents).
- Report survival status.

[#]Reporting as AST or ALT is to be consistent throughout the study on a per participant basis.

Treatment Cycle 1, Day 28

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
- Pregnancy test for WOCBP (only if clinically indicated).
- Quality of Life Assessments
- Compliance assessment – reconcile quantity of unused tablets.

- Record and report any adverse events and/or toxicity assessments that have occurred since commencing initial treatment. If the patient has toxicity during the first 28 calendar days that requires treatment to be interrupted, the first drug off period will only commence if the ctDNA TAB at calendar day 28 is $\leq 50\%$ of baseline 1. If the ctDNA TAB at calendar day 28 is $> 50\%$ of baseline 1, the drug on period is to be extended until treatment has been administered for 28 days (non consecutive). This does not impact the timelines for randomisation.
- Report survival status.

During first drug off period

Unless toxicity occurs during cycle 1 (see section 8.2), the first drug off period commences automatically following the completion of the initial 28 day cycle. Central instruction will not be provided.

2 weeks from day 1 drug off period

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
- Optional research biopsy (see Section [9.139-13](#)).
- Pregnancy test for WOCBP (only if clinically indicated).
- Quality of Life Assessments.
- Report survival status.

Every 2 weeks from day 1 first drug off period

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
- Pregnancy test for WOCBP (only if clinically indicated).
- Report survival status.

Every 4 weeks from day 1 first drug off period'

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
 - Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH
- Is the participant taking any prohibited concomitant medication? If so, these should be stopped and/or alternatives found.
- Record and report any adverse events and/or toxicity assessments that have occurred since commencing initial treatment.
- Physical examination, including weight, ECOG performance status and vital signs.
- Pregnancy test for WOCBP (only if clinically indicated).

- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents).
- Report survival status.

#Reporting as AST or ALT is to be consistent throughout the study on a per participant basis.

Baseline 2 (Performed Once Within -2 Days of First Treatment Restart)

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
- Pregnancy test for WOCBP (only if clinically indicated)
- Dispense 14 day supply of study medication and reserve pack

The participant must be directed NOT to use the Reserve Pack unless directed to do so by the research team.

- Report survival status

IMPORTANT - Blood for ctDNA MUST be taken prior to recommencing E+B for the first time. Failure to collect a ctDNA sample within 2 days prior to recommencing E+B may result in the participant being withdrawn from DyNAMIC and reverting to the local care pathway.

Every 2 weeks (from day 1 of re-start)

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
- Pregnancy test for WOCBP (only if clinically indicated)
- Dispense 14 day supply of study medication, plus a reserve pack (if used)
- Report survival status

Every 4 weeks (from day 1 of re-start)

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
 - Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH
- Is the participant is taking prohibited concomitant medication? If so, these should be stopped and/or alternatives found.
- Physical examination, including weight, ECOG performance status and vital signs

- Pregnancy test for WOCBP (only if clinically indicated)
- Dispense 14 day supply of study medication, plus a replacement reserve pack (if used)
- Counsel participant on use of reserve pack
- Compliance assessment – reconcile quantity of unused tablets.
- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents)
- Record and report any adverse events and/or toxicity assessments that have occurred since commencing initial treatment.
- Report survival status

#Reporting as AST or ALT is to be consistent throughout the study on a per participant basis.

From 6 months post-randomisation, the day 15 visit is optional for Arm B participants only if approved by the MTB with regular monitoring as to whether this should continue by the trial team

Additional ctDNA blood tests

- If during the drug off period the investigator has concerns that the patient is clinically progressing (e.g. symptoms/LDH rising/clinical disease assessment) or if the last ctDNA result showed a rise off treatment which was sub-threshold to restart, or if the ctDNA result is uninformative, then, in discussion with the DyNAMIC trial team, an additional ctDNA test can be performed prior to the next protocol mandated ctDNA visit. The timing of this should be agreed with the DyNAMIC trial team and the NBC laboratory informed.
- Physical examination, including weight, ECOG performance status and vital signs should also occur
- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents)
- This visit can occur multiple times, however once the patient is confirmed to have progression with no clinical benefit they should not have any further unscheduled visits.

Delays or extenuating circumstances to ctDNA bloods being taken

- If the quantity of blood drawn for ctDNA analysis is insufficient for any reason, the blood drawn for future research will be re-purposed for this task.
- If there are extenuating circumstances e.g. admission at a different hospital, which result in the participant not being able to attend on the protocol specified day to collect ctDNA bloods for either Arm, then this must be discussed with the CI (or their representative) and the LCTC Trial Manager who will advise when to take the bloods for ctDNA.

As Per Site Schedule

- Echocardiogram (ECHO) - standard safety monitoring of participants receiving E+B, not a research requirement – please refer to SmPC.
- Electrocardiograph - standard safety monitoring of participants receiving E+B, not a research requirement – please refer to SmPC.

Every 8 Weeks From Cycle 1, Day 1

- CT (with or without contrast if known allergy to it) of thorax, abdomen and pelvis, +/- neck if clinically indicated.
Alternatively, as per local practice if known allergy to contrast, CT without contrast thorax (+/- neck), plus MRI without contrast of abdomen and pelvis. Scans to be reported according to RECIST v1.1 (See Appendix D).

Every 12 Weeks From Cycle 1, Day 1

- Quality of Life Assessments

Every 26 Weeks From Cycle 1, Day 1

- MRI of brain, CT is an acceptable alternative if required according to local practice (both with contrast if no known allergy to it). Scans to be reported according to RECIST v1.1 (See Appendix D).

Discontinuation of Study Treatment

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
 - Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH
- Record and report any adverse events and/or toxicity assessments
- Pregnancy test for WOCBP following 28 days following the final administration of study drug.
- Physical examination, including weight, ECOG performance status and vital signs.
- Collect tissue from any surgically indicated surgical resection
- Additional blood for research (EDTA tube).
- Optional research biopsy (see Section [9.139-13](#)).
- Quality of Life Assessments.
- Compliance assessment – reconcile quantity of unused tablets.
- Report survival status

At Disease Progression and/or no clinical benefit

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
 - Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH
- Record and report any adverse events and/or toxicity assessments.
- Pregnancy test for WOCBP (if clinically indicated) following 28 days after administration of the last IMP dose.
- Physical examination, including weight, ECOG performance status and vital signs.
- Collect tissue from any surgically indicated surgical resection.
- Additional blood for research (EDTA tube).
- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents)
- Optional research biopsy (see Section [9.139-13](#)).
- Quality of Life Assessments.
- Compliance assessment – reconcile quantity of unused tablets.
- Report survival status.

[#]Reporting as AST or ALT is to be consistent throughout the study on a per participant basis.

Every 12 Weeks Following End of Study Treatment

- Record and report any adverse events and/or toxicity assessments.
- Collect tissue from any surgically indicated surgical resection.
- Record and report data on subsequent treatment(s), date(s) of treatment and any disease progression and survival status.

End of Trial Participation (excluding death)

- Pregnancy test for WOCBP following 28 days after administration of the last IMP dose. following the end of systemic treatment (only required once at either progression or end of treatment).
- Record and report any adverse events and/or toxicity assessments.
- ECOG performance status.
- Collect tissue from any surgically indicated surgical resection.
- Compliance assessment – reconcile quantity of unused tablets with treatment regimen.
- Record and report data on subsequent treatment(s), date(s) of disease progression of each, and survival status.

- Collect tissue from any standard of care indicated surgical resection.


If the participant is getting benefit from their targeted therapy, they should continue on the treatment as standard care.

Arm B ctDNA Real-Time Clinical Decision Making:

Unless toxicity occurs during cycle 1 (see section 8.2), the first drug off period commences automatically following the completion of the initial 28 day cycle. Central instruction will not be provided.

In order to determine when a participant should subsequently be “On-Drug” or “Off-Drug”, blood for ctDNA will be taken at 2 weekly intervals and a clinical decision made based on the result. The frequency of obtaining blood samples can only be changed by the MTB on a per patient basis unless there are exceptions as outlined in the ‘Additional ctDNA bloods tests’ section, above. Please refer to the DyNAMIC Laboratory Manual for details regarding sample collection tubes. Details regarding the decision as to the thresholds of tumour activity and burden (TAB level) are found in Section 5.15.4 and will be continuously reviewed by the DyNAMIC Trial Team during the study. The current thresholds as defined by the MTB will be used for each timepoint and may change whilst the patient is on the study, depending on the MTB decision. **The decision as to whether a participant should start or stop drug will be taken centrally and the participating site will be informed immediately. The site must then inform the participant immediately.** If the participant is currently “On-drug” and the ctDNA has reached a threshold to stop, then once the result is returned they should be informed to stop treatment the following day. If the participant is currently within a period “Off-Drug” and is being contacted to recommence treatment, they should be directed to use their Reserve Pack of E+B until a new 14 day cycle can be provided at their next clinic appointment. A new Reserve Pack will then need to be provided. Figure 7 shows the process of the blood being taken and actioning the result. The patient should restart drug within 24 hours of the result being returned (unless the baseline 2 needs to be taken).

Figure 7. Flowchart of process of ctDNA being processed and participant being informed as to dosing of E+B.



Schedule of Assessments:

All assessments and follow-up are to be conducted in line with the Schedule of Assessment tables, below. The legend for the numerical caveats indicated in superscript is in Section [9.119-14](#).

ARM B	Pre-Treatment		Randomisation (no more than 14 days post (C1 D1)	Cycle 1 D15 (SOC)	Cycle 1 D28 (SOC)	Drug Off Period		Adaptive treatment			As Per Site Schedule	Fixed Timelines			Discontinuation of study treatment	Disease Progression and/or no further clinical benefit ⁸	Follow-up every 12 weeks after end of study treatment	End of Study Participation	
Study Procedures	Screening (up to 42 days prior to C1 D1) ¹¹	Baseline 1 ¹¹				1 st drug off period	4 wks from D1 off drug 2 wks from D1 off drug	Baseline 2 (within 2 days prior to first restart of e+b)	From D1 of first restart of e+b			Additional ctDNA testing	Every 8 weeks From C1 D1	Every 12 weeks From C1 D1					Every 26 weeks from C1 D1
									2 weekly	4 weekly									
Window	42d			+/- 4d		+/- 4d	2d	+/- 4d				+/-1w	+/-2w	+/-2w	+28 d	+28d	+/-1w	+/-1w	
Eligibility check	x	x	x																
Written informed consent prior to any study procedure	x																		
Medical history and demographics	x																		
Review of prior anti-cancer therapies	x																		
Primary tumour block and any other tissue obtained through prior cancer-related surgery	x																		
Haematology and Biochemistry	x	x		x			x		x						x	x			
Blood for ctDNA (streck tube) including additional blood for research	x	x ¹ ₂		x	x	x	x	x	x	x					x	x			
Confirmation of screening BRAF ctDNA TAB level			x																
Prohibited medication review	x	x		x			x		x										
AE and toxicity assessment				x	x		x		x						x ⁹	x ⁹	x ⁹	x ⁹	
ECOG	x	x		x			x		x	x					x	x		x	
Weight	x						x		x	x					x	x			
Height	x																		
ECG	x										x								
Echocardiogram	x										x								
Confirmation of ECHO results			x																

ARM B	Pre-Treatment		Randomisation (no more than 14 days post (C1 D1))	Cycle 1 D15 (SOC)	Cycle 1 D28 (SOC)	Drug Off Period		Adaptive treatment			As Per Site Schedule	Fixed Timelines			Discontinuation of study treatment	Disease Progression and/or no further clinical benefit ⁸	Follow-up every 12 weeks after end of study treatment	End of Study Participation	
Study Procedures	Screening (up to 42 days prior to C1 D1) ¹¹	Baseline 1 ¹¹				1 st drug off period		Baseline 2 (within 2 days prior to first restart of e+b)	From D1 of first restart of e+b			Additional ctDNA testing	Every 8 weeks From C1 D1	Every 12 weeks From C1 D1					Every 26 weeks from C1 D1
						4 wks from D1 off drug	2 wks from D1 off drug		4 weekly	2 weekly									
Window	42d			+/- 4d		+/- 4d	2d	+/- 4d			+/-1w	+/-2w	+/-2w	+28 d	+28d	+/-1w	+/-1w		
CT scan (thorax, abdomen pelvis (+/- neck if clinically indicated) or MRI. PET-CT if clinically indicated (NB imaging modality must be consistent)	x										x								
CT or MRI brain	x												x						
Further tissue collection from surgical resection (only if clinically indicated) ¹	x													x	x	x	x		
Pregnancy test	x ⁶	x ⁴		x ⁶										x ⁷	x ⁷		x ⁷		
Research blood sample (EDTA tube)		x		x										x	x				
Physical exam ² , vital signs and clinical disease assessment	x	x		x			x		x	x				x	x				
Encorafenib plus binimetinib dispensed if required (on drug)		x						x	x	x									
Reserve pack of E+B dispensed if required ³								x	x	x									
Counsel participant on use of reserve pack								x	x	x									
Optional research Biopsy		x ⁵				x ¹⁰								x	x				
Quality of Life Assessments (EORTC QLQ-C30 and PRO-CTCAE)		x			x	x ¹³							x	x	x				
Compliance assessment					x					x				x	x		x		
Date of disease progression on each subsequent therapy																x	x		
Subsequent treatment data																x	x		
Survival status				x	x	x	x	x	x					x	x	x	x		

9.11 Definitions of Assessments:

Schedule of Assessments – Legend:

1. Further Tissue Collection - Tissue (FFPE and if possible, snap frozen) from any cancer-associated surgical procedure performed as part of routine care to be used for translational research. This can be at first and subsequent progressions following each line of treatment and prior to starting a new line of treatment (see laboratory manual and Section ~~9.139-43~~ for details).
2. Physical Exam - Symptom guided, except at screening.
3. Reserve Pack – Dispensed for the first time at Baseline 2, or if a replacement is required due to usage or dose reduction.
4. Baseline 1 Pregnancy Testing - Regardless of previous pregnancy tests, another must be done within 72 hours prior to commencing study treatment and have a negative result.
5. Optional Research Biopsy – The Baseline 1 Optional Research Biopsy may be performed within **42 days prior** to the initiation of Cycle 1 Day 1.
6. Pregnancy Testing - Serum or urine pregnancy test (minimum sensitivity 25 IU/L or equivalent units of HCG) only required if clinically indicated.
7. Discontinuation of Treatment/End of Study Pregnancy Testing – A serum or urine pregnancy test (minimum sensitivity 25 IU/L or equivalent units of HCG) must be taken 28 days after administration of the last IMP dose. Testing is only required once, i.e. following progression, discontinuation of Treatment or End of Study.
8. Disease Progression and/or no further clinical benefit. Only if there is no further clinical benefit should treatment stop. Procedures should occur within 28 days of confirmed progression and prior to starting any new systemic therapy.
9. SAEs should be actively monitored for 90 days following the final administration of study treatment.
10. Optional Research Biopsy - Required only once at Cycle 2, Day 15 (+/- 5 Days).
11. Baseline 1 Assessments - All assessments, except ctDNA collection, may be completed within 42 days of Cycle 1 Day 1. Where screening and baseline 1 have been conducted on separate visits, safety bloods must be taken within 2 days prior to the baseline visit. The echocardiogram can be performed within the 42 day window, or up to Cycle 1 Day 13, with the result available for randomisation to be conducted prior to Cycle 1 Day 14 of taking E+B.
12. Baseline 1 ctDNA – MUST be collected within 1 day prior to initiating Cycle 1 Day 1.
13. Quality of Life Assessment – to be completed at 2 weeks into the first drug off period.
14. Optional from 6 months post-randomisation
15. Quality of Life Assessment -Cycle 2 , Day 1 only.

General assessments:

Medical History and Demographics:

Medical history should include the following:

- Sex and age
- History of presenting complaint including date first noted symptoms, date of original primary diagnosis and date of disease progression to stage III unresectable, or stage IV disease.
- Date of pathological diagnosis and copy of original pathology/genomics report (confirming melanoma and BRAF point mutation status)
- Date of previous -related surgeries
- Co-morbidities and past medical/surgical history
- Family history including history of cancers, and significant illnesses
- Social history including occupation and smoking history.
- Review of all prior anti-cancer therapies

Concomitant Medications:

All concomitant medication including over-the-counter and herbal medications should be checked against prohibited medications (see Section [8.98-9](#)). If the participant is taking any prohibited medications these should be stopped and/or alternatives found, or the participant may be considered ineligible.

ECOG:

The Eastern Cooperative Oncology Group performance status should be assessed as summarised in Appendix E.

Physical Examination, Vital Signs and Clinical Disease Assessment:

1. Full physical examination will include symptom-guided assessments of the cranial nerve, eye and skin examinations and neck nodes (if prior melanoma on the head), the respiratory, cardiovascular, abdominal, musculoskeletal, neurological, dermatological and lymph nodes systems.
2. Vital signs (heart rate, blood pressure, respiratory rate, oxygenation level) will also be evaluated. Height will be measured at Screening only. Body weight will be observed in each examination.
3. Further targeted physical examinations are to be utilised by the Investigator on the basis of clinical observations and symptomatology.
4. Situations in which physical examination results should be reported as AEs are described in Section [11.711.7](#).

Imaging and clinical assessments:

1. Contrast enhanced CT scan of thorax, abdomen and pelvis every 8 weeks (+/- 1 week).
2. PET-CT as clinically indicated (participants with limb disease).
3. Contrast enhanced CT is the preferred modality for assessment. If participant is allergic to contrast, a non-contrast CT of the thorax and MRI of liver and pelvis is an alternative.
4. CT of the neck to be included as clinically indicated.
5. In addition, a CT or MRI of brain with contrast where possible should be performed every 26 weeks (+/- 2 Weeks).
6. Tumour efficacy (RECIST) assessment dates are not affected by IMP dose delays and remain as originally scheduled, as they are based on the date of Cycle 1 Day 1.
7. Cutaneous lesions can be included for measurable disease if photographed with a ruler depicting clearly the size of target lesion(s). The same method will be used to assess response of the cutaneous lesions identified at baseline throughout the trial, that is, two tumour assessments (one for each method) will be performed at each planned assessment time if the ruler method is used.

Safety Assessments:

Laboratory Assessments – Haematology and Biochemistry:

Table 1 . Haematology Laboratory Tests

Haemoglobin	Mean corpuscular volume
Lymphocytes (ALC)	Platelet count
Eosinophils	Total white cell count
Neutrophils (ANC)	

Table 2 . Clinical Chemistry (Serum or Plasma) Laboratory Tests

Albumin	Gamma glutamyltransferase
Alkaline phosphatase	Glucose
Alanine aminotransferase	Lactate dehydrogenase (hospital normal range should be provided)
Aspartate aminotransferase	Magnesium
Corrected Calcium	Phosphate
C-reactive protein ^a	Potassium
Creatinine	Sodium

Creatinine clearance ^b	Total bilirubin ^a
Creatinine kinase (CK)	Urea or blood urea nitrogen, depending on local practice

a. At baseline 1 and on progression only

b. eGFR will be derived using Cockcroft-Gault equation (see Appendix A)

Pregnancy Test:

Women of childbearing potential (WOCBP) must have a serum or urine pregnancy test at Screening and subsequently within 72 hours prior to commencing Cycle 1 Day 1 of therapy (minimum sensitivity 25 IU/L or equivalent units of HCG). A final pregnancy test must be performed 28 days following administration of the final study dose. All other pregnancy tests are required only if clinically indicated.

Echocardiogram:

This is a standard safety monitoring of participants receiving E+B, not a research requirement – please refer to SmPC. Echo must however be done as part of Screening and left ventricular function confirmed to be $\geq$ LLN prior to Randomisation. All other echo scans may be performed as per site schedule.

If grade 3-4 LVEF decrease or an absolute decrease of LVEF from baseline of $\geq 10\%$ is observed, then E+B should be discontinued and LVEF should be evaluated every 2 weeks until recovery.

ECG:

A single resting 12-lead ECG will be recorded at Screening and as a standard safety monitoring of participants receiving E+B, which is not a research requirement – please refer to SmPC and perform as per local site protocols throughout the study. ECGs should be obtained after the participant has been in a supine position for 5 minutes and recorded while the participant remains in that position. The ECG should be monitored as part of standard of care for any evidence of prolonged QT interval.

Quality of Life Assessments:

The following assessments will be performed every 12 weeks from Cycle 1 Day 1 and at Disease Progression. (An additional completion of the questionnaires will occur at 2 weeks into the first drug off period for participants on Arm B.)

- EORTC QLQ-C30 – global health status score. This will be calculated as per scoring instructions in the EORTC QLQ-C30 manual. A descriptive comparison will be made between the two arms for the change from baseline in global health status score at each timepoint between Arm A and Arm B. In addition, comparisons will be made between participants within Arm B who are “On-Treatment” with those off treatment during an adaptive therapy cycle.
- PRO-CTCAE – a patient reported outcome (PRO) measurement system developed to evaluate symptomatic toxicity. The questions selected will be based on anticipated adverse events observed in the Phase III Columbus study and the encorafenib and binimetinib SmPCs. A descriptive

comparison will be made between the two arms for the change from baseline in global health status score at each timepoint between the standard and the experimental arms. In addition, comparisons will be made between participants who are on therapy with those off therapy during an adaptive therapy cycle.

9.12 Disease progression definition and procedures

Definition of disease progression:

Arm A – Disease progression is defined in accordance with RECIST v1.1 (2009).

Arm B – Disease progression is defined in accordance with RECIST v1.1 (2009) disease progression **when the participant is receiving** targeted therapy “On-treatment”. If RECIST v1.1 disease progression is observed when the participant is not on treatment “Off-Treatment”, they should recommence targeted therapy. If the next scan/clinical assessment when “On-Treatment” shows further disease progression, then that will be the RECIST v1.1 disease progression event. If the participant is still obtaining clinical benefit, they may continue “On-Treatment” and will only stop when no further clinical benefit is achieved. If the next scan shows response then they will continue “On-Treatment”. ctDNA blood tests will continue to be compared against the Baseline 2 TAB to determine ctDNA response as to whether the participant may switch to an “Off-Treatment” period again. This will be determined by the DyNAMIC trial team in discussion with the participating site PI as to the clinical picture.

Procedures following disease progression and/or no further clinical benefit:

The following should be performed within 28 days, and prior to the start of any new systemic treatments, when disease progression is confirmed:

- Blood tests
 - ctDNA (streck tube - inc. extra for research purposes)
 - Full blood count
 - Renal and liver[#] function tests
 - Urea and electrolytes
 - LDH
- Record and report any adverse events and/or toxicity assessments.
- Research bloods (EDTA tube).
- Optional research biopsy (see Section [9.139-13](#)). Ideally from a progressing lesion. Modality as clinically indicated. Biopsies will only be acquired if the lesions are considered to be safely accessible by the interventional radiologist or surgeon. Please refer to the DyNAMIC lab manual for prioritising and initial processing of tissue samples.

- Pregnancy test for WOCBP (if clinically indicated) following 28 days after administration of the last IMP dose.
- Is the participant taking prohibited concomitant medication? If so, these should be stopped and/or alternatives found.
- Physical examination, including weight, ECOG performance status and vital signs.
- Clinical disease assessment – must be measured. If also photographed, a ruler must be shown alongside to assess the scale of the measurements. (Images are retained by site as source documents).
- Quality of Life Assessments.
- Collect tissue from any surgically indicated surgical resection.
- Compliance assessment.
- Report survival status.

#Reporting as AST or ALT is to be consistent throughout the study on a per participant basis.

Follow-up after progression:

Following progression patients are followed up remotely every 3 months until study completion/withdrawal or death, in order to capture the following data:

- Any subsequent anti-cancer therapy taken after study treatment discontinuation should be documented.
- Date of disease progression on each subsequent anti-cancer therapy should be documented.
- Survival data including data of death should be captured.
- Record any adverse events and/or toxicity assessments.
- Collect tissue from any surgically indicated surgical resection.

Radiotherapy

- If a new lesion or RECIST v1.1 progression this should be recorded as a progression event.
- If the treating clinician feels that the patient can still obtain clinical benefit from targeted therapy following the radiotherapy then they can continue on the DyNAMIC treatment schedules – this should be discussed via the DyNAMIC trial team at the LCTC.
- Radiotherapy (including SRS and SABR) to any lesion to improve cancer-related symptoms or to treat isolated progression is allowed for either Arm.
- Local site protocols should be used for dose and fractionation and withholding targeted therapy during treatment period if necessary.
- Information regarding site, dose and fractionation should be collected.

Surgery

- Surgery to any lesion is allowed during the trial to improve cancer-related symptoms or to treat isolated progression is allowed for either Arm.
- If a new lesion or RECIST v1.1 progression, this should be recorded as a progression event.
- If the treating clinician feels that the patient can still obtain clinical benefit from targeted therapy following the surgery then they can continue on the DyNAMlc treatment schedules – this should be discussed via the DyNAMlc trial team at the LCTC.

Samples including FFPE and optionally (but ideally) snap frozen should be taken of the surgical specimen (see section 9.13). **Please refer to the DyNAMlc Laboratory Manual for detailed information and instructions.**

9.13 Sampling

Please refer to the DyNAMlc Laboratory Manual for detailed information and instructions

The following specimens may be collected as part of consent to the trial.

For current covid-19 precautions please refer to the laboratory manual.

Sample/collection procedures include:

- FFPE of archival/primary tumour specimen.
- Blood in EDTA tube for translational studies.
- Blood for ctDNA analysis.
- Fresh biopsy of easily accessible lesions (up to a maximum of three cores per procedure), at baseline 1, when drug is removed on Arm B (C2 D15 +/-5 days) or when drug is present on Arm A (C2 D15 +/-5 days) and if disease progression on therapy. Participants may consent to any or all of the above biopsies.
- If a surgical resection on study occurs for any clinical reason, then provided the participant has consented this tissue should also be made available for translational research with collection as below.

Sample Collection:

An outline for the collection, processing, storage and shipping of samples is specified below. Please refer to the DyNAMlc Lab Manual for more detailed instructions.

If ctDNA samples are missed, the participant will need to be re-called to clinic for a repeat sample to be taken.

If the ctDNA result does not pass quality control testing at the CRUK NBC, or if the sample becomes exhausted prior to the result being known, then the translational sample will be used to repeat the test.

For other missed samples, the reason and which sample/timepoint should be noted on the eCRF.

Tissue:

- Archival FFPE primary tumour tissue and any tissue from previous cancer-related surgeries will be requested for all participants.
- Fresh biopsies (up to a maximum of three cores per procedure) will be obtained if the site is able to perform the biopsy and according to participant consent, as set out in the sample schedule. Biopsies will only be acquired if the lesions are considered to be safely accessible by the interventional radiologist or surgeon. Ideally the same site of disease should be biopsied on each occasion. Please refer to the DyNAMIC Laboratory Manual for prioritising and initial processing of tissue samples. If insufficient material is available participants will not be subjected to an additional biopsy.
- If the participant undergoes a clinically indicated surgical procedure to remove cancer or biopsy of a lesion following consent, then samples will be obtained for translational research purposes. These should include FFPE samples and if possible fresh frozen tissue. Please see the DyNAMIC Laboratory Manual for further details.

Blood:

- Blood collected in Cell-Free DNA BCT Streck tubes for ctDNA analysis
- At scheduled time points, as detailed in Sections [9.99.9](#) and [9.109.10](#). 40-80 ml of blood will be collected in Cell-Free DNA BCT Streck tubes (provided by central trial team).
- Bloods collected in streck tubes must be sent in a mailing box with next day delivery, provided by the trial, on the day of taking – see lab manual for address.
- For samples taken at The Christie, the research nurse (or delegate) will contact NBC to inform that blood samples are ready for collection.
- *Blood collected in EDTA tubes for translational studies*
- At scheduled timepoints, 10ml blood will be collected in EDTA tubes – see schedule of assessments and laboratory manual for further details.

Sample Storage and Handling:

Please refer to the DyNAMIC Laboratory Manual for details regarding processing, handling and storage of biospecimens.

All collected biospecimens will be stored in designated and secure facilities within the NBC prior to analysis using a destructive test, or with the research team using secure sponsor approved facilities pending future use. Frozen tissue and plasma/blood pellet samples will be stored in temperature monitored, -80°C freezers. To preserve participant confidentiality, samples will use only the participant trial ID Number for identification and all samples will catalogued using a tracking system.

Custodianship:

The Chief Investigator will serve as custodian of samples for the duration of this study. Samples for ctDNA analysis will be stored short-term at the NBC laboratory or will be used for translational work, with other samples transferred to the research team for storage using sponsor approved facilities, pending future use. Sample usage will be under the control and jurisdiction of the Chief Investigator and Principal Investigators. Samples will be stored at site as per the Lab Manual for a period of up to the end of study before being transferred by courier or via post to the relevant laboratory. Clinical samples will be kept for up to 25 years and any surplus to the study requirements may be used for future translational research authorised by the trial management committee.

Disposal of biospecimens will be considered under certain circumstances including but not limited to use of the entire specimen; withdrawal of participant consent for storage and analysis of samples (obtained archival samples will be returned to original pathology lab) or reduced specimen integrity. Determination of the integrity of biospecimens is at the discretion of study personnel. Disposed samples will be logged within the tracking system (together with reason for disposal) and physically discarded according to institutional policy.

Sharing of samples with other centres:

Samples may be shared with other centres for research directly in connection with the existing protocol or for future translational work. This may include transfer of samples to collaborators in other countries. Where required, additional approvals will be sought to share samples and a Material Transfer Agreement will be put in place before samples are shared. This will be managed with the assistance of our human tissue governance team and R&D office.

9.14 Intervention Discontinuation and Participant Discontinuation/Withdrawal

In consenting to the trial, participants agree to all trial activities including administration of trial intervention and treatment and follow-up assessments / visits and data collection. Every effort should be made to facilitate the completion of these for every recruited participant. If it is not possible to complete these activities (or it is deemed inappropriate) the reasons why should be documented. The following sub-sections describe the different levels of discontinuation/withdrawal.

Premature Discontinuation of Trial Intervention

Participants may discontinue treatment for reasons including, but not limited to:

- Participant-led i.e. request by the participant
- Unacceptable toxicity (see Section 11 for safety reporting)
- Intercurrent illness preventing further treatment.
- Pregnancy
- Death
- Clinician-led:

- Any change in the participant's condition that justifies the discontinuation of treatment in the clinician's opinion.
 - Reasons of non-adherence or non-compliance with treatment or other trial procedures
 - Participant unable to receive E+B for >6 weeks
- Discontinuation from study intervention does not mean discontinuation of the study altogether, and the remaining study procedures, follow-up assessment / visits and data collection should be completed as indicated in the protocol (unless consent is specifically withdrawn, see section 9.2).

Participants who have discontinued treatment as per protocol will then be transferred into routine clinical care and will be treated as per local practice and guidelines.

Safety follow up is covered in section 11.6.

Participant Withdrawal from Follow-up:

Participants are free to withdraw from follow-up at any time without providing a reason, though a reason should be recorded if one is given. Those who wish to withdraw from further follow-up will have the data collected up to the point of that withdrawal included in the analyses, inclusive of data derived from the analyses of tissue samples. The participant will not contribute further data to the study and the LCTC should be informed via email to the LCTC and via completion of a Withdrawal eCRF to be completed within 7 days.

If participants express a wish to withdraw from follow-up, participants should be given appropriate care under medical supervision for all ongoing adverse events until the symptoms resolve or the participant's condition becomes stable. Any SAEs will be notifiable to the LCTC via processes detailed in Section 11.10 even if a participant has withdrawn from follow-up.

If the participant withdraws their consent for any remaining samples to be included in analyses, all remaining samples originating from that participant will be destroyed. No remaining samples, except archival FFPE tumour samples, that are stored centrally will be returned to the treating hospital.

Participant Transfer:

If a participant moves from the area, every effort should be made for the participant to be followed-up at another participating trial centre and for this trial centre to take over responsibility for the participant or for follow-up via GP.

The participant remains the responsibility of the original site until the Transfer eCRF has been completed.

Loss to Follow-up:

A participant will be considered lost to follow-up if they fail to return for four consecutive scheduled visits and are not contactable by the site research team. Methods of contact are detailed below.

If a participant fails to attend a required study visit the following actions must be taken and the efforts recorded in the participant medical notes.:

- One scheduled visit missed – Site to telephone participant to establish reasons for non-attendance, advise the participant on the importance of maintaining the assigned visit schedule, and attempt to re-arrange visit within allowed timelines.
- Two consecutive scheduled visits missed – Site to telephone participant to establish reasons for non-attendance, advise the participant on the importance of maintaining the assigned visit schedule. An appointment must be made with the participant for treatment safety purposes.
- Three consecutive visits missed – Same as two consecutive visits being missed. If, however, the participant cannot be contacted by telephone, the REC approved reminder letter is to be sent to their last known address.
- Four consecutive visits missed – Participant is classed as a loss to follow-up. Withdraw from study treatment and submit relevant eCRF to LCTC.

9.15 End of Trial

The end of the trial is defined to be the date on which data for all participants is frozen and data entry privileges are withdrawn from the trial system. The trial may be closed prematurely by the Trial Steering Committee (TSC), on the recommendation of the Independent Data and Safety Monitoring Committee (IDSMC).

Site and closure activities will be centrally coordinated and conducted in accordance with LCTC processes regardless of whether the trial closes as planned or prematurely. This includes activities such as:

- End of Trial notification to REC and MHRA
- Trial-related materials reconciled and returned/disposed of as appropriate. See pharmacy manual for further details on the destruction of medication.
- All site data entered onto the study system, discrepancies raised and satisfactory responses received.
- Quality Control checks of the Investigator Site Files, Pharmacy Files and Trial Master File as appropriate.

10 Management of COVID-19

Patients who are COVID-19 positive at the time-point of a scheduled procedure should be advised to not attend hospital for their appointment, and the appointment re-scheduled for 7 days later. In this instance, procedures with a visit tolerance of less than seven days, e.g. bloods for ctDNA +/-2 days, will not be recorded as a protocol deviation.

For participants on Arm B, any changes to their adaptive treatment will also be deferred accordingly.

11 SAFETY REPORTING

Safety reporting in clinical trials is a legal and ethical requirement and it is imperative that all applicable requirements detailed here are followed during the trial.

11.1 Terms and Definitions

Adverse Event (AE):

Any untoward medical occurrence in a patient or clinical trial subject administered a medicinal product and which does not necessarily have a causal relationship with this treatment.

An adverse event can therefore be any unfavourable and unintended sign (including an abnormal laboratory finding, for example), symptom or disease temporally associated with the use of a medicinal product, whether or not considered related to the medicinal product.

Adverse Reaction (AR):

Any untoward and unintended response to an investigational medicinal product related to any dose administered.

Unexpected Adverse Reaction (UAR):

An adverse reaction the nature and severity of which is not consistent with the information about the medicinal product in question set out in the investigator's brochure (IB) or the Summary of Product Characteristics (SmPC), which may be referenced where the IMP in question is a product with a marketing authorisation.

Serious Adverse Event (SAE):

Any untoward medical occurrence or effect that at any dose results in death, is life-threatening, requires hospitalisation or prolongation of existing hospitalisation, results in persistent or significant disability or incapacity, or is a congenital anomaly or birth defect.

Serious Adverse Reaction (SAR):

An adverse reaction which meets the definition of serious (see Section [11.244.2](#)) is a Serious Adverse Reaction. A Serious Adverse Reaction event that has been assessed as 'expected' (see Section [11.544.5](#)) according to the Reference Safety Information (see below) will remain classified as a Serious Adverse Reaction only, however some Serious Adverse Reactions that are considered 'unexpected' will be further classified as a Suspected Unexpected Serious Adverse Reaction (SUSAR) (see below).

Suspected Unexpected Serious Adverse Reaction (SUSAR):

An adverse reaction that is classed in nature as serious and "unexpected" (i.e. not listed within the

Reference Safety Information (RSI) approved for the trial by the MHRA and current at the time of onset of the SUSAR).

11.2 Assessment of Seriousness

The assessment of seriousness of safety events must be performed by an appropriately delegated, medically qualified member of the site research team.

A safety event / reaction is assessed as serious if it:

- Results in death;
- Is life threatening (i.e. the investigator considers the event places the subject at immediate risk of death from the experience as it occurred (this does not include an adverse experience that, had it occurred in a more severe form, might have caused death);
- Requires hospitalisation or prolongation of existing hospitalisation (hospitalisation is defined as an inpatient admission, regardless of length of stay, even if the hospitalisation is a precautionary measure for continued observation. Hospitalisations for a pre-existing condition, including elective procedures that have not worsened, do not constitute an SAE);
- Results in persistent or significant disability or incapacity (substantial disruption of one's ability to conduct normal life functions);
- Consists of a congenital anomaly or birth defect (in offspring of subjects, or their partners, taking the IMP regardless of time of diagnosis);
- Other important medical events (these may not result in death, be life-threatening, overdose, secondary malignancy or require hospitalisation, but may be considered a serious adverse event/experience when, based upon appropriate medical judgment, they may jeopardise the subject and may require medical or surgical intervention to prevent one of the outcomes listed in this definition).

11.3 Severity of Adverse Events

All adverse events should be assessed for severity. The assignment of the severity/grading should be made by the investigator responsible for the care of the participant using the definitions in the table below:

Table 3: Severity Grading

Severity	Description
Mild	Does not interfere with routine activities.
Moderate	Interferes with routine activities.
Severe	Impossible to perform routine activities.
Life-Threatening	Life threatening at the time of the event, i.e. not <i>potentially</i> life-threatening.
Death	Caused by the event.

N.B. A distinction is drawn between **serious** and **severe** AEs. Severity is a measure of intensity (see above) whereas seriousness is defined using the criteria in Section 11.2. Hence, a severe safety event need not necessarily be a “serious” safety event.

11.4 Assessment of “Causality” - Relationship to Trial Treatment/Intervention

The assignment of the causality should be made using the definitions in the table below:

Table 4: Definitions of Causality:

Relationship	Description	Categorisation
Unrelated	There is no evidence of any causal relationship. N.B. An alternative cause for the AE should be given	No reasonable possibility
Unlikely	There is little evidence to suggest there is a causal relationship (e.g. the event did not occur within a reasonable time after administration of the trial medication). There is another reasonable explanation for the event (e.g. the participant’s clinical condition, other concomitant treatment).	No reasonable possibility
Possibly	There is some evidence to suggest a causal relationship (e.g. because the event occurs within a reasonable time after administration of the trial medication). However, the influence of other factors may have contributed to the event (e.g. the	Reasonable possibility

Relationship	Description	Categorisation
	participant's clinical condition, other concomitant treatments).	
Probably	There is evidence to suggest a causal relationship and the influence of other factors is unlikely.	Reasonable possibility
Almost certainly	There is clear evidence to suggest a causal relationship and other possible contributing factors can be ruled out.	Reasonable possibility

Events that are assessed as being possibly, probably or almost certainly related will be reported as having a reasonable possibility of being related, and events assessed as unrelated or unlikely will be reported as having no reasonable possibility of being related.

Assessment of causality should be made based on known safety profiles of the SmPC and known risk profiles of other drugs in the same class. If any doubt about the causality exists, the local investigator should inform the LCTC who will notify the Chief Investigator. In the case of discrepant views on causality between the treating investigator and others, the causality assessment of the treating investigator will never be downgraded and the MHRA and REC will be informed of both points of view.

11.5 Assessment of “Expectedness”

The Chief Investigator and nominated Medical Reviewers for the DyNAMIC trial are responsible for determining whether a safety event is expected or unexpected, however the Chief Investigator or Medical Reviewer will not assess safety events that occur at their own Trust. There is no requirement for a reporting investigator to make an assessment of expectedness.

An event will be considered unexpected if it is not listed within the current and approved Reference Safety Information (RSI) (see section [11.544.5](#)) for the study at the time of onset of the event. The nature, severity, or frequency of the event should be considered – if this is not consistent with that described for the type of event in the RSI the event should be assessed as unexpected.

Reference Safety Information / Information used to Assess Expectedness

The Reference Safety Information (RSI) to be used for this trial for each IMP is as follows. Versions of each are as approved via original application or subsequent amendment:

Summary of Product Characteristics Mektovi 15 mg film-coated tablets Section 4.8

Summary of Product Characteristics Braftovi 50 mg hard capsules Section 4.8

This will be updated as required and implemented only after approval from the appropriate regulatory authorities.

11.6 Time Period for Active Monitoring of Safety Events

IMPORTANT: Any serious adverse event in a trial participant after the end of the “active monitoring” period must continue to be reported by sites to the LCTC if the PI becomes aware of them.

Active monitoring of safety events experienced by trial participants will commence immediately following consent being provided by the participant, and continue until 90 days after the definitive final administration of E+B. Care should be taken in establishing if SAEs for participants on Arm B meet the latter criterion. E.g. if an Arm B participant experiences an SAE on day 95 of being off-treatment, it still needs reporting to LCTC as it is not yet known if E+B will re-commence.

During the course of the study, all AEs and SAEs should be proactively followed up for each subject. Every effort should be made to obtain a resolution for all events, even if the events continue after discontinuation/study completion. All unresolved SAEs and AEs must be followed up until resolution or death. Follow-up must be conducted by the Investigator for as long as medically indicated, and recording in source documents. The Sponsor retains the right to request additional information for any subject with ongoing AE(s)/SAE(s) at the end of the study, if judged necessary

Pregnancies will be followed up at least until the outcome is known (see Section 11.8 for more information on reporting pregnancy).

11.7 Notes on Safety Event Recording

The following events must be recorded for the purposes of the trial:

- An exacerbation of a pre-existing illness
- An increase in frequency or intensity of a pre-existing episodic event/condition
- A condition (even though it may have been present prior to the start of the trial) detected after trial drug administration
- Continuous persistent disease or symptoms present at baseline that worsens following the administration of the study/trial treatment
- Laboratory abnormalities that require clinical intervention or further investigation (unless they are associated with an already reported clinical event).
- Abnormalities in physiological testing or physical examination that require further investigation or clinical intervention
- Injury or accidents
- A new cancer diagnosis

- Overdose of medication with signs or symptoms
- Pregnancy

Do not record:

The events below do not need recording as the trial is considered low risk:

- Medical or surgical procedures - the condition which leads to the procedure is the adverse event
- Pre-existing disease or conditions present before treatment that do not worsen
- Situations where an elective medical occurrence has occurred e.g. cosmetic surgery
- Overdose of medication **without signs or symptoms***
- The disease being treated or associated symptoms/signs unless more severe than expected for the patient's condition

¹ Note that although overdose of medication without signs or symptoms may be excluded from AE reporting it may still require investigation to ensure other protocol and regulatory requirements are met e.g. for IMP management and administration, or to ensure participant safety. If applicable, refer to appropriate part of Treatment section [8.108-10](#) (Overdose).

*N.B. If overdose occurred **with** resulting signs and symptoms that meet the protocol criteria for AE/AR/SAE/SAR/SUSAR then they should be reported accordingly (see section 11.10 for more information).

11.8 Reporting of Pregnancy

Pregnancies of participants, and where consent is obtained to do so partners of male participants, are to be reported to LCTC. If a participant becomes pregnant during the course of the study, the IMPs should be discontinued immediately.

If pregnancy occurs during either the intervention or specified follow-up period of the trial this must be notified to the LCTC and within 24 hours of the research team becoming aware. The pregnancy must be followed up by the site research team until the outcome of the pregnancy and reported to LCTC.

Pregnancies of female participants are reportable if conception is assessed to have occurred within one month of the last administration of E+B. Pregnancies of partners of male participants are reportable if conception is assessed to have occurred within three months of the last administration of E+B (i.e. sperm regeneration cycle).

Any pregnancy outcomes which result in a safety event assessed as "serious" (e.g. birth defect) must also be reported separately on the appropriate Safety Event eCRFs in accordance with processes described in section 11.10. All pregnancies and outcomes reported to LCTC will be notified to the study Sponsor and monitored by trial oversight committees.


11.9 Notification of Deaths

If the research team become aware of the death of a participant (whether related to the trial or not) this should be notified to the LCTC using the appropriate eCRF within 24 hours of becoming aware.

11.10 Reporting Procedures

All safety events which are recorded for the study should be reported following the procedures detailed below. The occurrence of a safety event may come to the attention of research staff during routine study visits, from the participant's notes, directly from the participant or by other means. Note that reporting procedures vary dependent on the nature of the incident (i.e. "serious" events are to be reported to LCTC in an expedited manner). Any questions concerning adverse event reporting should be directed to the LCTC in the first instance. A flowchart is given below to aid in determining reporting procedures for different types of adverse events.

Flowchart for Site Reporting Requirements of Adverse Events:



Reporting Safety Events to the LCTC:

All safety events (whether or not assessed as serious / related / expected) should be recorded on an electronic Adverse Event Form; multiple events can be recorded on one form. When documenting any adverse events the correct medical terminology **must** be used.

Safety events which are assessed as “serious” must **also** be recorded in more detail on electronic Serious Safety Event Forms; a **single form** is used for each individual event (i.e. a single diagnosis), though multiple symptoms can be recorded. Each SAE should have a corresponding record on the participant’s AE form. Where additional information is received by site after initial submission to LCTC, this should be provided on a follow-up form within 5 days. Electronic Serious Safety Event Forms collect data regarding the nature of event, date of onset, severity, corrective therapies given, outcome and causality; all serious events reported to LCTC will be reviewed by the Chief Investigator or Medical Reviewer, and assessed for causality and expectedness.

Follow-up After Adverse Events:

All reportable adverse events should be followed until satisfactory resolution or until the investigator responsible for the care of the participant deems the event to be chronic or the participant to be stable.

When reporting “serious” safety events the investigator responsible for the care of the participant should apply the following criteria to provide information relating to event outcomes:

- resolved
- resolved with sequelae (specifying with additional narrative)
- not resolved/ongoing
- ongoing at final follow-up
- fatal or unknown.

11.11 Investigator Reporting Responsibilities

The PI is responsible for ensuring that all safety events which the local research team becomes aware of, and which require recording on this study are reported to LCTC. It is the responsibility of the PI and other medically qualified doctors as recorded on the Delegation Log to assess the seriousness and causality of events.

All safety events must be recorded on the electronic AE form, which is to be reported to LCTC within 7 days following any updates.

Unless the SAE is specified in the protocol as not requiring immediate reporting (see Section 11.7), safety events which meet the definition of “serious” must be reported in more detail to the LCTC on an electronic

SAE form and reported **immediately and in no circumstances later than 24 hours from becoming aware**, where they will be appropriately processed.

The eSAE form should be completed by an appropriately delegated member of the research team; the assessments of seriousness and causality must be performed by an appropriately medically qualified person. Minimum reporting information must be provided in all initial reports.

The minimum dataset required for a preliminary report should include the following:

Minimum information required for reporting:

- Study site identifier
- Participant study number
- Name of reporting site staff member
- Description of the event including date of onset
- Suspect IMP
- Seriousness assessment
- Causality assessment

N.B. In the absence of a delegated medically qualified person, the eSAE form should be completed and signed by an alternative member of the research site trial team and submitted to the LCTC. As soon as possible thereafter, the responsible investigator should check the eSAE form, make amendments as appropriate, sign and re-send to the LCTC. The initial report shall be followed by detailed follow-up reports as appropriate.

Safety events should be reported to the site R&D team in accordance with local policy.

Reporting an initial or follow-up SAE:

The investigator should ensure the actions below are completed for all reportable SAEs:

1. Research sites should telephone the appropriate DyNAMIC Trial Manager (see DyNAMIC Contact List) to advise that an SAE report has been submitted as soon as possible.
2. **The eSAE form should be reported via the electronic clinical database system within 24 hours of becoming aware.**
3. The responsible investigator must notify their R&D department of the event (as per standard local governance procedures).
4. The participant must be identified by participant trial number, age and initials **only**. The participant's name **should not** be used on any correspondence.

5. SAEs must be subsequently followed up in line with the processes below:
 - a. Follow-up must continue until clinical recovery is complete and laboratory results have returned to normal, or until the event has stabilised (see Section 11.6). N.B. Follow-up may continue after completion of protocol treatment if necessary.
 - b. Follow-up information is noted on a new eSAE form to be reported to the LCTC as soon as more information becomes available.
 - c. Indicate on the eSAE form the type of report; this is dependent on resolution status of the SAE e.g. follow-up / final.
6. Extra, annotated information and/or copies of anonymised test results should only be submitted when requested by the medical reviewer via LCTC.
7. Blank Serious Adverse Event forms can be obtained from the LCTC by contacting the DyNAMIC Trial Manager. Contact details are available within the LCTC contacts document.

11.12 LCTC Responsibilities

The trial Sponsor, The Christie NHS Foundation Trust, have delegated to LCTC the duty of onward reporting of safety events to REC, regulatory authorities and the Sponsor. The Sponsor is responsible for the reporting of safety events to Pierre Fabre Ltd, who are also the marketing authorisation holder for both encorafenib and binimetinib. LCTC SOPs will be followed to ensure appropriate reporting as detailed below.

All “serious” safety events will be forwarded to the Chief Investigator or Medical Reviewer by LCTC within 24 hours of receiving the minimum information from site. The CI (or delegate) will review information provided by site and for all events assessed as “related” will provide an assessment of “expectedness”.

Safety events which are assessed as serious, related and unexpected adverse events will be expedited to the REC and MHRA within the following timeframes (in accordance with Part 5 of the Medicines for Human Use (Clinical Trials) Regulations SI1031 (Sections 32-25)):

- SUSARs which are fatal or life-threatening – **as soon as possible and in any case no later than 7 days** after the LCTC is first aware of the event. If the initial report is incomplete, a complete report will be submitted within an additional 8 days.
- SUSARs that are not fatal or life-threatening – within **15 days** of the LCTC first becoming aware of the event.

A copy of the report will be provided to the Sponsor in parallel with correspondence to the REC and MHRA. Additionally, SUSARs will be reported to the Principal Investigators of participating sites in line with the requirements of LCTC Standard Operating Procedures (SOP).

The LCTC will submit a Development Safety Update Report to the MHRA on an annual basis.

The following safety issues should also be reported in an expedited fashion:

- An increase in the rate of occurrence or a qualitative change of an expected serious adverse reaction, which is judged to be clinically important.
- Post-study SUSARs that occur after the participant has completed a clinical trial and are notified by the Investigator to the LCTC.
- New events related to the conduct of the trial or the development of the IMPs and likely to affect the safety of the subjects, such as:
 - a. A SAE which could be associated with the trial procedures and which could modify the conduct of the trial;
 - b. A significant hazard to the subject population, such as lack of efficacy of an IMP used for the treatment of a life-threatening disease;
- Recommendations of the Independent Data and Safety Monitoring Committee where relevant for the safety of the subjects.

The PIs at all institutions participating in the trial will be notified of any SUSARs at a frequency defined by LCTC SOP.

Any concerns raised by the TSC or IDSMC or inconsistencies regarding safety reporting noted as part of central monitoring may prompt additional training at sites, with the potential for the LCTC to carry out site visits if there is suspicion of unreported AEs / ARs and SARs / SAEs in participant case notes. Additional training will also be provided if there are unacceptable delays in safety reporting timelines.

Maintenance of Blinding in Adverse Event Reporting:

Not applicable, open label study.

Safety Reports:

Safety reports will be generated during the course of the trial which allows for monitoring of safety event including reporting rates and safety events by site / arm. The LCTC will send annual developmental safety update reports (DSURs) containing a list of all SAEs and SARs to the IDSMC and MHRA. If any safety reports identify issues that have implications for the safety of trial participants, the PIs at all institutions participating in the trial will be notified.

The DSUR will present a comprehensive annual review and evaluation of pertinent safety information collected during the reporting period relating to the Investigational Medicinal Products. It will cover the following 4 areas:

1. Examine whether the information obtained by the Sponsor during the reporting period is in accordance with previous knowledge of the investigational drugs' safety.
2. Describe new safety issues that could have an impact on the protection of clinical trial subjects.

3. Summarise the current understanding and management of identified and potential risks.
4. Provide an update on the status of the clinical investigation/development programme and study results have implications for the safety of trial participants, the PIs at all institutions participating in the trial will be notified.

Urgent Safety Measures (USMs):

An urgent safety measure (USM) is a procedure to protect clinical trial participants from any immediate hazard to their health and safety but has not previously been defined by the protocol. It can be put in place prior to authorisation by the REC and the MHRA.

The Sponsor/LCTC will notify the MHRA and REC immediately and, in any event, within 3 days that such a measure has been taken and the reasons why it has been taken. The initial notification to the REC and MHRA will be by telephone (ideally within 24 hours) and a notice in writing will be sent within 3 days, setting out the reasons for the USM and the plan for further action. After discussion with the REC and MHRA, further action will be agreed, which may include submission of a substantial amendment, a temporary halt, or permanent termination of the trial.

Following notification, if a substantial amendment is required this must be submitted as soon as possible to the REC and ideally within two weeks to the MHRA. If the study is temporarily halted it may not recommence until authorised to do so by the REC and MHRA. If the study is permanently terminated before the date specified for its conclusion (in the original applications to REC and MHRA), the Sponsor should notify the REC and MHRA within 15 days of the date of termination by submitting the formal End of Trial Notification.

11.13 Contact Details and Out-of-hours Medical Cover

E+B are both currently utilised within standard NHS practice and both E+B have a well-established safety profile emergency and out-of-hours medical care will be in line with usual NHS arrangements and local standard practice; no special provision is required for DyNAMIC participants. All participants will be provided with a copy of the information sheet which includes information about their participation and contact details for the local research team who may be contacted if necessary. During office hours, the CI or delegate are able to provide medical advice in relation to participation using the contact details listed at the beginning of this document.

12 DATA MANAGEMENT AND TRIAL MONITORING

For the DyNAMIC trial the responsibilities for Data Management and monitoring are delegated to the LCTC. Separate Data Management and Trial Monitoring Plans have been developed which provide detail regarding the internal processes that will be conducted at the LCTC throughout the trial. Justification for the level of monitoring is provided within those documents and the trial-specific risk assessment. All data will be managed as per local LCTC processes and in line with all relevant regulatory, ethical and legal obligations.

12.1 Source Documents

The case report form (eCRF) will be considered the source document for data where no prior record exists and which is recorded directly in the bespoke eCRF, e.g. participant completed quality of life questionnaires. A DyNAMIC source document list will be produced for each site to be kept in the ISF and provide detail of what constitutes DyNAMIC-specific source data.

Date(s) of informed consent processes (including date of provision of participant information sheet(s), randomisation number and the fact that the participant is participating in a clinical trial (including possible treatment arms) should be added to the participant's medical record chronologically.

12.2 Data Collection Methods

Data are to be entered into an electronic secure web-based trial specific system by approved members of the research team at site. This is the primary data collection instrument for the study so all data requested **must** be recorded and all missing data must be explained. Missing or inconsistent data will be queried by LCTC, the system will be updated based on the response.

12.3 Monitoring

Monitoring is conducted to ensure protection of patients participating in the trial and all aspects of the trial (procedures, laboratory, trial intervention administration and data collection) are of high quality and conducted in accordance with sponsor and regulatory requirements.

A detailed Trial Monitoring Plan will be developed and agreed by the TMG and CI to describe who will conduct the monitoring, at what frequency monitoring will be done, whether central or on site and what level of detail monitoring will be conducted.

The purposes of site monitoring visits include, but are not limited to:

- assessing compliance with the study protocol;
- discussing any emerging problems that may have been identified prior to the visit;
- source data verification.

12.4 Risk Assessment

In accordance with the LCTC SOPs, a risk assessment has been completed in partnership with:

- Representatives of the Trial Sponsors
- Representatives of CRUK National Biomarker Centre
- CI and Co-Investigators
- Trial Manager
- Trial Statistician
- LCTC Senior Trial Manager
- LCTC Information Systems
- LCTC Data Management
- LCTC Quality Assurance

In conducting this risk assessment, the contributors considered potential participant, organisational and study hazards, the likelihood of their occurrence and resulting impact should they occur.

The outcome of the risk assessment for a Clinical Trial of an Investigational Medicinal Product (CTIMP) is expressed as an overall risk level as set out below, assigned according to one of the following categories:

CTIMP Type A = Comparable to the risk of standard medical care

CTIMP Type B = somewhat higher than the risk of standard medical care

CTIMP Type C = markedly higher than the risk of standard medical care

The risk assessment resulted in a **CTIMP Type B**.

12.5 Confidentiality

This trial will collect personal data (e.g. participant names), including special category personal data (i.e. participant medical information) and this will be handled in accordance with all applicable data protection legislation. Data (including special category) will only be collected, used and stored if necessary for the trial (e.g. evidencing provision of consent, for data management and central monitoring, statistical analysis, regulatory reporting, etc.). At all times, this data will be handled confidentially and securely.

ECRFs will be assigned a unique Participant Number. Verification that appropriate informed consent is obtained will be enabled by the provision of copies of participant's signed informed consent forms being supplied to the LCTC by recruiting sites. This transfer of identifiable data is disclosed in the PISC.

N.B. Consent forms must be transferred separately to any other trial documentation to ensure the pseudonymisation of special category data is maintained.

Site-specific study-related information will be stored securely and confidentially at sites and all local relevant data protection policies will be adhered to.

The LCTC as part of The University of Liverpool will preserve the confidentiality of participants taking part in the study.

The data controller for this study is The Christie NHS Foundation Trust, which is also the sponsor, and is registered as a data controller with the Information Commissioners Office. The University of Liverpool and The University of Manchester are data processors and will preserve the confidentiality of participants taking part in the study.

Breaches of data protection principles or regulations identified by LCTC will be notified promptly to the trial Sponsor and The University of Liverpool's Data Protection Officer and appropriate processes followed.

12.6 Quality Assurance and Control

To assure protocol compliance, ethical standards, regulatory compliance and data quality, as a minimum, the following will occur:

- The PI and other key staff from each centre will attend initiation training, which will incorporate elements of trial-specific training necessary to fulfil the requirements of the protocol.
- The TMG will determine the minimum key staff required to be recorded on the delegation log in order for the centre to be eligible to be initiated.
- The Trial Manager (or delegate) at the LCTC will verify appropriate approvals are in place prior to initiation of a centre and the relevant personnel have attended the trial specific training. A greenlight checklist will verify all approvals are in place prior to trial initiation at LCTC and the individual centre.
- The trial will be conducted in accordance with procedures identified in the protocol.
- The IDSMC and independent members of the TSC will provide independent oversight of the trial.
- The TMG will monitor screening, randomisation and consent rates between centres and compliance with the protocol.
- Data quality checks and monitoring procedures will be undertaken in line with the trial Data Management Plan.

12.7 Records Retention

The retention period for the DyNAMIC data and information is 25 years from the official End of Trial date (defined in section [9.159-15](#) above).

The PI at each investigational site must make arrangements to store the essential trial documents (as defined by ICH GCP guidelines) including the Investigator Site File, the applicable participant medical records and

Pharmacy Site File, for the full length of the trial's retention period and will arrange for confidential destruction at the end of this period as instructed by the LCTC.

The PI is also responsible for archiving all relevant source documents so that the trial data can be compared against source data after completion of the trial (e.g. in case of inspection from authorities). They must ensure the continued storage of the documents, even if they, for example, leave the clinic/practice or retire before the end of required storage period. Delegation of responsibility for this must be documented in writing.

All other persons and organisations involved in the trial will be responsible for storing and archiving the parts of the TMF relevant to their delegated duties (e.g. laboratories, IMP manufacturers and distributors, third-party vendors providing randomisation and IMP allocation systems, etc.).

The LCTC undertakes to archive as per their contractual requirements; documents will be archived in compliance with the principles of GCP. All electronic CRFs and trial data will be archived onto an appropriate media for long term accessible storage. Hard copies of data will be boxed and transferred to secure premises where unique reference numbers are applied to enable confidentiality, tracking and retrieval.

13 REGULATORY AND ETHICAL CONSIDERATIONS

13.1 Statement of Compliance

This study is designed to comply with the guideline developed by the International Conference on Harmonisation (ICH) for Good Clinical Practice (GCP) and will be conducted in compliance with the protocol, LCTC Standard Operating Procedures and the UK Statutory Instrument 2004 No 1031: Medicines for Human Use (Clinical Trials) Regulations 2004 and all subsequent amendments. Local sites must conduct research in compliance with the UK Policy Framework for Health and Social Care Research.

13.2 Ethical Considerations

The DyNAMIC trial will be conducted in accordance with, but not limited to, the UK Data Protection Act (DPA) 2018, Freedom of Information Act 2000 subject to the provisions of Sections 41 and 43 thereof, the Medicines for Human Use (Clinical Trials) Regulations 2004 (as amended), the Medicines Act 1968, the Human Tissue Act 2004, the principles of the World Medical Association Declaration of Helsinki 1996 and the UK Policy Framework for Health and Social Care Research, as amended from time to time.

Participants will be asked to consent that data are recorded, collected, stored and processed and may be transferred to other countries, in accordance with the UK DPA 2018 and UK General Data Protection Regulation (GDPR) and to allow a copy of their completed signed consent form to be sent to the LCTC.

This study may be terminated at the request of the CI, IDSMC, Independent Research Ethics Committee (REC) or the MHRA if, during the course of the study, concerns about the safety of further dosing emerge.

The CI will update the ethics committee of any new information related to the study drug when appropriate and this will also be disseminated to the Principal Investigators at each trial centre.

13.3 Approvals

The protocol, PISC and any proposed public-facing material will be submitted to an appropriate Research Ethics Committee (REC), MHRA, Health Research Authority (HRA) and host institution(s) for written approval.

Any substantial amendments to the original approved documents will be submitted and, where necessary, approved by the above parties before use.

13.4 Non-Compliances

Deviations from, breaches or violations of, or non-compliance to either the protocol, the conditions or principles of GCP, and relevant regulatory and ethical e.g. MHRA and REC requirements are handled based on their nature and severity, thus:

Non-Serious breaches:

Protocol deviations and other non-serious breaches of GCP etc. will be managed according to local site and LCTC procedures as appropriate. They will be reported to trial oversight committees.

Serious breaches:

A breach of the protocol or GCP is 'serious' if it meets the definition of being "likely to affect to a significant degree the safety or physical or mental integrity of the trial participants, or the scientific value of the trial". This assessment can only be determined by the Sponsor.

If any persons involved in the conduct of the trial become aware of a potential serious breach, they must immediately report this to the LCTC who will in turn notify the Sponsor. The Sponsor will assess the breach and determine if it meets the criteria of a 'serious' breach.

The Sponsor may seek advice from medical expert members of the TMG and/or of the independent oversight committees (IDSMC and TSC) in determining whether or not the breach is likely to affect to a significant degree the safety, physical or mental integrity of participants.

The Sponsor may seek advice from the Trial Statistician in determining whether or not the breach is likely to significantly affect the scientific value of the trial. However, the Sponsor retains responsibility for the assessment of whether or not a breach meets the definition of 'serious' and is subject to expedited reporting to MHRA and REC.

Breaches confirmed as 'serious' will be reported to MHRA and REC within 7 days by the LCTC on behalf of the Sponsor and notified to the TMG, IDSMC and TSC at their next meeting.

Any requests for additional information from the Sponsor, TMG, TSC, IDSMC, MHRA or REC, will be promptly actioned by the relevant member(s) of the research team and open communication will be maintained to ensure appropriate corrective actions are taken and documented.

Incidents of protocol non-compliance will be recorded as protocol deviations, the incidence of which are monitored and reported to trial oversight committees.

14 INDEMNITY

The Christie NHS Foundation Trust holds insurance against claims from participants for harm caused by their participation in this clinical study. However, the treating hospital continues to have a duty of care to the participant and the Sponsor does not accept liability for any breach in the hospital's duty of care, or any negligence of the part of hospital employees. In these cases, clinical negligence indemnification will rest with the participating NHS Trust or Trusts under standard NHS arrangements.

15 PUBLICATION AND DISSEMINATION

15.1 Publication Policy

The results from different participating sites will be analysed together and published as soon as possible, maintaining participant confidentiality at all times. Individual clinicians must undertake not to submit any part of their individual data for publication without the prior consent of the Trial Management Group (TMG).

The TMG will form the basis of the writing committee and will advise on the nature of publications. The Uniform Requirements for Manuscripts Submitted to Biomedical Journals (<http://www.icmje.org/>) will be respected. All publications shall include a list of participants and if there are named authors these should include the trial's Chief Investigator Paul Lorigan and Sub investigator Rebecca Lee, representative (s) from NBC, Statistician(s) and Trial Manager(s) involved as a minimum. If there are no named authors (i.e. group authorship) then a writing committee will be identified that would usually include these people, at least. The ISRCTN allocated to this trial will be attached to any publications resulting from this trial and members of the TSC and IDSMC should be acknowledged.

Any publications arising from this research will be reviewed appropriately prior to publication.

Authorship:

Contributors to all four of (i) the design, conduct, data analysis and interpretation, (ii) writing, (iii) manuscript approval and (iv) accountability for the integrity of the work will, depending on their contribution and journal requirements, be included by name at the manuscript head or listed at the end in a by-line as members of the DyNAMIC TMG which will also be named at the manuscript head.

15.2 Dissemination to Key Stakeholders

On completion of the research, a Final Trial Report will be prepared and submitted to the MHRA and REC. The results of DyNAMIC will be published regardless of the magnitude or direction of effect.

15.3 Data Sharing

At the end of the trial, after the primary results have been published, the Sponsor will hold responsibility for the production of an anonymised individual participant data (IPD) set. All requests for access to the IPD will be considered by the Sponsor.

16 CHRONOLOGY OF PROTOCOL AMENDMENTS

16.1 Version 1.0 (05/05/2022)

- Original Approved version

16.2 Version 2.0

- This version was never released. All changes are implemented into version 3.

16.3 Version 3.0 (28/11/2022)

- Expansion of exclusion criteria
- Removal of scans of the brain at every 8 weeks
- Re-classification of trial as a 'Type B'
- Addition of pregnancy tests for WOCBP following the end of systemic treatment and if clinically indicated
- Clarification that E+B are classed as IMP, regardless of the treatment arm
- Initial ctDNA thresholds for stopping and re-starting treatment included, along with a summary of how these will be monitored
- Update to co-enrolment guidelines
- Correction to inconsistencies in the timelines for the first re-start of treatment on Arm B
- Correction to timeline for the collection of data for adverse events
- Correction to inconsistencies in terminology, i.e. brain vs head and chest abdomen pelvis (CAP) vs thorax abdomen pelvis (TAP)

16.4 Version 4.0 (27/03/2023)

- Dose reduction tables removed and replaced with reference to the SmPC
- Elaboration to the proposed translational work on samples
- Formalisation of the Molecular Tumour Board as an Oversight Committee
- Requirement added to confirm consistency of scanning modality for the duration of a participants involvement, and that comparisons should not be made between images from differing modalities
- Removal of CRP blood test at all visits except baseline and progression
- Elaboration to the requirements for a physical examination of the head
- Update to advise translational blood sample will be used as a reserve for ctDNA analysis if required
- Section added on the management of participants with COVID-19
- Minor administrative changes

16.5 Version 5.0 (21/05/2024)

- ISRCTN and CPMS reference added.
- Statement of use for generic terminology for participant trial ID inserted.
- Contact details and Glossary updated.
- A two-step process for confirming eligibility added.
- Timeline for exclusion criterion 1 has been updated from 6 months to 26 weeks.
- Definition of participant and study duration has been clarified.
- Secondary Objective 4 amended to define clinical deterioration.
- Secondary Objective 5 updated to clarify that overall survival will be compared in Arm A vs Arm B.
- A new secondary objective added to assess melanoma specific overall survival in Arm A vs Arm B.
- Minor amendments made to Secondary Objective 10 “To assess quality of life” and Secondary Objective 11 “To assess toxicity (all grade adverse events)”.
- Secondary Objective 12 has been added to the summary page.
- 2.1 schematic of the study design has been removed.
- Funder’s contact details removed.
- Reference to CRFs amended to eCRF throughout.
- Table of protocol contributors updated and moved to appendices.
- Section 4.4 amended to remove duplication and the table updated to align with the objectives on the summary page.
- Section 5.1 updated to add a statement to clarify MTB decision making and to give instructions to PIs to notify the CI if a scan shows progression.
- Figure 5: Adaptive treatment schedule updated.
- Section 7.5 guidance on use of barrier method of contraception updated to align with main guidance.
- Section 8.1 table removed to reduce duplication. Labelling details have moved to Section 8.3.
- Added guidance for treatment continuation following detection of clinical deterioration (Section 8.2).
- Diagram in Section 8.2 removed to reduce duplication.
- The pharmacy dispensing diagram updated to clarify that the reserve pack is given in Cycle 1 day 28.
- Section 8.5 updated to clarify that if study drug is withdrawn due to toxicity for more than 6 weeks, the participant is to be withdrawn from the trial treatment.
- Removal of the requirement for use of Patient Diaries. Compliance will be monitored via pill counts.
- Clarifications and additions to Section 8.9 “Medications Not Permitted/Precautions Required”.
- Section 9.2 updated to confirm availability of quality of life questionnaires in multiple languages
- Details regarding the study database have moved to section 9.3
- Additional sentence added to confirm that screening may only occur once.
- Change to echocardiogram timing at screening (Section 9.4).
- Addition of sentence confirming that blood samples taken for ctDNA analysis may be re-drawn in the event of loss or lack of sample viability (Section 9.4).
- Section 9.6 “Commencement of Initial Targeted 28 day E+B Cycle” added
- Section 9.7 updated to specify randomisation requirements.
- Schedule for ECHO and ECG amended to allow sites to follow their local schedule.
- The requirement to do pregnancy tests every 8 weeks, 12 and 26 weeks has been removed as they occur every 2 weeks.
- Survival status will be collected throughout the trial.
- Data collection requirements for discontinuation of treatment have been added to 9.9 and 9.10.

- Disease relapse replaced with “Disease progression” (RECIST v1.1) and/or no further clinical benefit.
- Requirement to capture prohibited medications has been removed from timepoint of disease progression and every 12 weeks following end of study treatment. Compliance data will be captured.
- Requirement to record adverse events for Arm B cycle 1 day 28 has been added.
- Requirement for a compliance check every 2 weeks for Arm B has been amended to every 4 weeks.
- A section has been added for Arm B participants experiencing delays or extenuating circumstances to ctDNA bloods being taken.
- Arm B participants now also complete QoL two weeks into first drug off period.
- Tables for Arm A and B updated in line with the changes outlined above.
- The legend in section 9.11 has been simplified to reduce duplication of information.
- Removed requirement to collect ethnicity (Section 9.11).
- Change to use of mail-boxes for transport of blood samples.
- Removed reference to meeting the exclusion criteria as reason for discontinuing treatment.
- Section 9.14 gives guidance if a participant missing 2 or more consecutive follow-up appointments.
- SAE Reporting Investigator guidance updated to reflect the use of eSAE reporting.
- Section 11.6 updated to clarify the period of active monitoring of safety events.
- Minimum information required for safety reporting has been updated.
- Section 12.3 updated to remove duplicated information that is detailed in the Monitoring Plan.
- Section 15.3 updated to reflect responsibility for dealing with data sharing access requests.
- Minor administrative changes made throughout for clarification purposes.

16.6 Version 6.0 (05/03/2025)

- Contact details, Institutions - Trial pharmacist email address updated.
- Contact details, Institutions (and throughout) - Central laboratory details updated.
- Contact details, Individuals - Addition to pool of Medical Experts who will evaluate SAE reports.
- Glossary table updated.
- Sections 2 and 7.1 - Erroneous reference to measurable brain disease removed from inclusion criterion 5, and alternative abbreviations for BRAF mutation points added to inclusion criterion 6. An appendix for the latter has also been added.
- Section 2 - Running order of eligibility criteria updated to align with section 7.2.
- Sections 2, and 7.3 - Criterion 3 updated to clarify that only current metastasis to the brain/leptomeninges is excluded.
- Section 4.3 – Updated to include the above.
- Section 5 - Trial Schema diagram updated to differentiate between the processes for commencing treatment pre- vs post- randomisation.
- Section 5.1 - The Molecular Tumour Board (MTB) will also convene ad hoc.
- Section 5.1 (and throughout) - Clarification the scheduled convening of the MTB will occur after every 6th occurrence of recommenced therapy, and not every 6th participant on Arm B, e.g. 6, 12, 18.
- Section 8.1 (and throughout) – Change to the timeline for the dispensing of the first Arm B reserve pack. Supporting advice on counselling the participant on its use also relocated to new timepoint within sections 9.9 and 9.10.
- Section 8.2 (repeated in 9.10) – Management information added for when treatment stops for the first time on Arm B following an interruption to treatment in cycle 1 due to toxicity.
- Section 8.2 - Correction to the timeline for the use of ACUTY.

- Section 8.3 - Caveat added for local pharmacy arrangements to be allowed if the site cannot accommodate the requirements for Arm B.
- Section 8.3 - Pharmacy dispensing diagram updated to accommodate the change to the timeline of dispensing the first reserve pack
- Section 8.7 - Assessment of compliance updated so only instances of non-compliance will need capturing.
- Sections 9.1 and 9.2 - Review of eligibility by the multi disciplinary team updated to the clinical team
- Section 9.1 - Clarification the result of the eligibility ECHO must be available prior to randomisation.
- Sections 9.4, 9.5, 9.9, 9.10 and 9.12 - Clarification that the method of reporting liver functionality (AST or ALT) must be consistent on a per participant basis.
- Sections 9.5, 9.9, 9.10 and 9.12 – Deletion of erroneous texts about the requirement of CT scans.
- Section 9.9 - From 6 months post-randomisation, visits for participants on Arm A may reduce to every 4 weeks.
- Section 9.9 - Information added on the timeline for the final study specific pregnancy test for WOCBP, and text on procedures at End of Participation corrected.
- Section 9.9 and 9.10 - Caveats added to the schedule of events tables. Legend in section 9.11 also updated.
- Section 9.10 – Text added to clarify a ctDNA is required as a specific timeline
- Section 9.10 - Clarification added that the Baseline 2 ctDNA blood sample must be taken prior to recommencing treatment for the first time. Typographical error of '-2' updated to '2' days prior.
- Section 9.10 - Expansion to the delays or extenuating circumstances to ctDNA bloods being taken.
- Section 9.10 - Flowchart for ctDNA processes updated to include the reduction of visits.
- Section 9.10 -Correction to the fixed timeline for brain scans on the schedule of events table for Arm B.
- Section 9.11 - Update to legend to the timeline for dispensing the reserve pack, and the Baseline 1 optional research biopsy. Clarification added to the timeline for obtaining the results of the baseline echocardiogram, and addition that 2 weekly visits on Arm A are optional from 6 months post-randomisation.
- Section 9.12 - Addition of allowance of radiotherapy during follow-up after progression.
- Section 9.13 - Update to local logistics at The Christie for transportation of ctDNA blood samples to NBC. Method of delivery for non ctDNA samples from all sites updated to include posting as an option.
- Section 9.15 - Removal of the Annual Progress Report to REC.
- Section 11.1 - Correction to remove pregnancies as an SAE.
- Throughout - Minor updates to e.g. grammar, job roles, abbreviations.

16.7 Version 7.0 (23/06/2025)

- Contact details- minor formatting changes
- Section 2. Protocol overview, updated to reflect changes to main text
- Section 4.4. clarification to secondary outcome measures from radiological progression to just progression.
- Section 5.0, removal of requirement for disease progression to be shown on scan
- Section 5.1, removal of requirement for disease progression to be shown on scan
- Section 7.1 update to inclusion criteria 2 to include acral
- Section 7.2 change to inclusion criteria 13: BRAF ctDNA TAB level reduced to ≥ 7 copies/ml of plasma

- Section 7.3 exclusion criteria 3, text updated to active metastasis of the brain and additional guidance added.
- Section 8.2 removal of requirement for disease progression to be confirmed on imaging/radiological.
- Section 8.2 Arm B, clarification on when treatment commences.
- Section 8.2 Arm B, additional wording added to confirm participant will restart E+B within 24hrs of the ctDNA result being returned,
- Section 9.4 additional text added to allow re-screening and also clarification on when a patient can start screening.
- Section 9.6 removal of erroneous text on timing of screening echocardiogram.
- Section 9.9 Treatment Cycle 2(onwards), day 1 addition of Quality of Life Assessments.
- Section 9.9 addition of an unscheduled visit and the required assessments.
- Section 9.9 Discontinuation of Study treatment, addition of assessments to reflect changes to schedule of assessments table.
- Section 9.9 Schedule of assessments table. Addition of unscheduled visit and related assessments to table. Updates to windows for study procedures. Option to omit day 15 from cycle 6 onwards added.
- Section 9.9 Schedule of assessments table. Quality of life assessments added at cycle 2 day 1.
- Section 9.9 Schedule of assessments table. Haematology and Biochemistry, weight, tissue collection from surgical resection, research blood sample, optional research biopsy and quality of life assessments added to discontinuation of study treatment time point.
- Section 9.10 Treatment Cycle 1, Day 28 Quality of Life Assessments added.
- Section 9.10 section added on additional ctDNA blood tests.
- Section 9.10 Discontinuation of Study treatment, addition of assessments to reflect changes to schedule of assessments table.
- Section 9.10 Arm B ctDNA real-time clinical decision-making section. Text added to clarify the use of the current thresholds and when the participant should stop/restart treatment,
- Section 9.10 Schedule of assessments table. Additional ctDNA testing time point added and associated assessment. Updates to windows for study procedures.
- Section 9.10 Schedule of assessments table. Haematology and biochemistry, weight, tissue collection from surgical resection, research blood sample, physical exam, vital signs, clinical disease assessment, optional research biopsy and quality of life assessments added to discontinuation of study treatment time point.
- Section 9.10 Schedule of assessments table. Quality of life assessments added at cycle 1 day 28.
- Section 9.11 definitions of assessments. Definition 8 updated, definition 15 added.
- Section 9.12 update to definition of disease progression.
- Section 9.12 procedures following disease progression. Addition of requirement to record and report and adverse events and/or toxicity assessments. Clarification of timing of pregnancy test for WOCBP. Addition of requirement to report survival status.
- Section 9.12: Addition of surgery section and associated guidance.

16.8 Version 8.0 (14/01/2026)

- Minor admin to:
 - add a public registry reference,
 - update to the job title of the person authorising on behalf of sponsor, and address of LCTC,
 - reformat the table of contact information of those who will assess SAE Reports,
 - correct throughout the erroneous texts where applicable of '≥15 copies/ml plasma' to '≥7 copies...' that were overlooked in the previous amendment,

- correct throughout the erroneous texts of 'radiological disease progression' to 'disease progression' that were overlooked during the previous update.
- Correct throughout the erroneous texts of 'Participants will be treated and undergo scans until confirmed disease progression...' to 'Participants will be treated until confirmed disease progression...'.
- correction to erroneous outcome measures texts of 'Maximal radiological response' to 'Maximal response', and 'sites of progression on imaging' to 'sites of progression' that were overlooked during the previous update.
- Correction to the lower range of the lowest stratification factor from '15-50' to '7-50', that was overlooked in the previous amendment.
- References to appendices. Updated following the removal of appendix C, affiliated to the now deleted eligibility criteria on pre-existing conditions.
- References of '...discuss with the DyNAMIC trial team at the LCTC.' for medical queries, updated to '...discuss via the DyNAMIC trial team at the LCTC'.
- Section 2: Eligibility criteria updated to align with changes in sections 7.1 to 7.3.
- Section 2: Quantity of study centres updated from 11 to 11-15, with the same update made to all references in later sections.
- Section 2: Follow-up period reduced from 52 to 26 weeks (-6 months), off-set by extending the recruitment period from 24 to 30 months (+6 months), with the same update made to all references in later sections.
- Section 5: Figure 4 amended to remove dosing information from columns.
- Section 5.1: Clarified that current ctDNA threshold levels will be used in establishing if treatment stops or re-starts,
- Section 5.1: Remit of Molecular Tumour Board (MTB) expanded to assess whether the frequency of ctDNA testing should be changed on a per participant basis.
- Section 5.1: Inclusion of conversations with the MTB of individual participants outside of formal meetings (decisions will be documented).
- Section 6.2: Update to average recruitment required per site from 1 to 1-3 participants/year/site.
- Section 6.5: Deletion of paragraph relating to missing data.
- Section 7.1: Inclusion at screening criterion 5 expanded to also allow disease that is clinically accurately measurable,
- Section 7.1: Inclusion criterion 9 updated to remove the blood results that define adequate organ function, replaced with '...organ function enabling safe administration of the study treatments. (The footnotes related to the blood results were also deleted.)
- Section 7.2: Section header reworded from 'Inclusion criteria on or prior to randomisation' to 'Inclusion criteria prior to randomisation'.
- Section 7.2: Removal of the 'at screening or prior to randomisation' timelines for compliance with eligibility. Remaining text reworded.
- Section 7.2: Criterion 13 expanded to advise that discrepancies between screening 1 and baseline ctDNA samples should be discussed with the trial team.
- Section 7.2: Relocation of footnotes to suit revision to texts.
- Section 7.3: Exclusion criterion 3 updated so the designation of active brain metastasis is determined by the local clinical team, informed by recent imaging, and not based specifically on the results of a scan.
- Section 7.3: Addition of a caveat to exclusion criterion 7 to allow discussion with the central trial team regarding eligibility related to historical other malignancies.
- Section 7.3: Deletion of exclusion criteria 10-16 related to specific pre-existing medical conditions, which will be captured in criteria 4 and 8 related to contraindications to treatment and serious or unstable pre-existing medical conditions.

- Section 7.3: Renumbering of subsequent exclusion criteria 17-19 to 10-12.
- Section 7.4: Update to co-enrolment guidelines. Timeline for eligibility updated and clarified it relates to days since administration of IMP on previous CTIMP that must have elapsed to be considered for DyNAMIC.
- Section 8.1: Dose reductions due to toxicity replaced with information on commencing trial treatment at a reduced dose.
- Section 8.1: Removal of references to the reserve pack needing to contain only a 10 day supply of IMP.
- Section 8.2: Text added relating to commencing trial treatment at a reduced dose.
- Section 8.2: Text added to indicate extra visits for ctDNA testing may reduce to every 4 weeks after 6 months on trial. For those on Arm B, this needs approval from the MTB.
- Section 8.2: Clarification that first re-starts of treatment will occur with 24 hours of the result being made available.
- Section 8.2: Change to running order of text within the section.
- Section 8.3: Separate texts to discuss site inability to comply with dispensing requirements replaced with a single sentence that covers all scenarios.
- Section 8.3: Correction to formatting errors on the pharmacy dispensing diagram.
- Section 8.5: Relaxation of the requirement for participants to withdraw from the study if withdrawal of treatment due to toxicity exceeds 6 weeks.
- Section 9.1: Screening window extended from 28 to 42 days to enable better recruitment and sites having to repeat scans which are just outside of the original window.
- Section 9.5: Removal of stipulation to confirm that the participant still complies with eligibility criteria at baseline.
- Section 9.7: Updated so baseline ctDNA TAB result needs to have been reported, not that the value of the result re-confirms eligibility. Any discrepancy needs to be discussed.
- Section 9.7: Text added to advise that discrepancies between screening 1 and baseline ctDNA samples should be discussed with the trial team.
- Section 9.9: Addition of text advising that reporting of ctDNA TAB results for those on Arm A will be deferred.
- Section 9.9: Table of procedures amended to align with updates.
- Section 9.10: Extra instruction added regarding treatment stopping following cycle 1. (Repeated within the real-time clinical decision making text of the same section.)
- Section 9.10: Missing the baseline 2 blood sample updated from 'will' result in withdrawal' to 'may' result in withdrawal.
- Section 9.10: Deletion of erroneous text related to participant not using reserve pack prior to this visit.
- Section 9.10: Texts added to indicate extra visits for ctDNA testing may reduce to every 4 weeks after 6 months on trial. This will require approval by the MTB.
- Section 9.10: Expansion to the reasons for additional ctDNA blood tests being required.
- Section 9.10: Table of procedures amended to align with updates.
- Section 9.11: Numerous updates to references to the screening timeline, from 28 to 42 days to match the expanded screening period.
- Section 11.5: Updated to advise that medical reviewers cannot assess reports of SAEs for participants at their Trust, not just those reports where they are also the reporting clinician.
- Section 11.6: Minor change to running order of text

17 REFERENCES

1. Thompson JF, Scolyer RA, Kefford RF. Cutaneous melanoma. *Lancet* [Internet]. 2005 Jan [cited 2015 Jan 13];365(9460):687–701. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15721476>
2. Cancer Research UK. Skin cancer statistics [Internet]. Cancer Research UK; 2015 [cited 2015 Jan 13]. Available from: <http://www.cancerresearchuk.org/cancer-info/cancerstats/types/skin/>
3. WHO | Skin cancers. WHO [Internet]. 2017 [cited 2019 Aug 5]; Available from: <https://www.who.int/uv/faq/skincancer/en/index1.html>
4. Cancer Research UK. Melanoma skin cancer mortality statistics [Internet]. 2021 [cited 2021 Apr 2]. Available from: <https://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/melanoma-skin-cancer/mortality>
5. Cancer Research UK. Melanoma skin cancer statistics [Internet]. [cited 2020 Oct 11]. Available from: <https://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/melanoma-skin-cancer#heading-One>
6. Gershenwald JE, Scolyer RA, Hess KR, Sondak VK, Long G V., Ross MI, et al. Melanoma staging: Evidence-based changes in the American Joint Committee on Cancer eighth edition cancer staging manual. *CA Cancer J Clin* [Internet]. 2017 Nov 1 [cited 2018 Feb 24];67(6):472–92. Available from: <http://doi.wiley.com/10.3322/caac.21409>
7. Ugurel S, Röhm J, Ascierto PA, Becker JC, Flaherty KT, Grob JJ, et al. Survival of patients with advanced metastatic melanoma: The impact of MAP kinase pathway inhibition and immune checkpoint inhibition - Update 2019. *Eur J Cancer*. 2020 May 1;130:126–38.
8. Hodi FS, O'Day SJ, McDermott DF, Weber RW, Sosman J a, Haanen JB, et al. Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* [Internet]. 2010 Aug 19;363(8):711–23. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23439861>
9. Robert C, Thomas L, Bondarenko I, O'Day S, M D JW, Garbe C, et al. Ipilimumab plus dacarbazine for previously untreated metastatic melanoma. *N Engl J Med* [Internet]. 2011 Jun 30;364(26):2517–26. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21639810>
10. Ishida Y, Agata Y, Shibahara K, Honjo T. Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. *EMBO J* [Internet]. 1992 Nov 1 [cited 2021 May 5];11(11):3887–95. Available from: <https://www.embopress.org/doi/full/10.1002/j.1460-2075.1992.tb05481.x>
11. Robert C, Long G V., Brady B, Dutriaux C, Maio M, Mortier L, et al. Nivolumab in Previously Untreated Melanoma without BRAF Mutation. *N Engl J Med* [Internet]. 2014 Nov 16 [cited 2014 Nov 16];141116004513004. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25399552>
12. Robert C, Schachter J, Long G V., Arance A, Grob JJ, Mortier L, et al. Pembrolizumab versus Ipilimumab in Advanced Melanoma. *N Engl J Med* [Internet]. 2015 Jun 25 [cited 2017 Feb 24];372(26):2521–32. Available from: <http://www.nejm.org/doi/abs/10.1056/NEJMoa1503093>
13. Larkin J, Chiarion-Sileni V, Gonzalez R, Grob JJ, Cowey CL, Lao CD, et al. Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma. *N Engl J Med* [Internet]. 2015 May 31 [cited 2015 Jun 2];373:23–34.

Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26027431>

14. Michielin O, Atkins MB, Koon HB, Dummer R, Ascierto PA. Evolving impact of long-Term survival results on metastatic melanoma treatment [Internet]. Vol. 8, Journal for ImmunoTherapy of Cancer. BMJ Publishing Group; 2020 [cited 2021 May 22]. p. 948. Available from: <http://jitc.bmj.com/>
15. Larkin J, Chiarion-Sileni V, Gonzalez R, Grob J-J, Rutkowski P, Lao CD, et al. Five-Year Survival with Combined Nivolumab and Ipilimumab in Advanced Melanoma. *N Engl J Med* [Internet]. 2019 [cited 2019 Oct 7];381(16). Available from: <http://www.nejm.org/doi/10.1056/NEJMoa1910836>
16. Wolchok JD, Chiarion-Sileni V, Gonzalez R, Grob J-J, Rutkowski P, Lao CD, et al. CheckMate 067: 6.5-year outcomes in patients (pts) with advanced melanoma. *J Clin Oncol* [Internet]. 2021 May 20 [cited 2021 Jun 9];39(15_suppl):9506–9506. Available from: https://ascopubs.org/doi/10.1200/JCO.2021.39.15_suppl.9506
17. Lipson EJ, Tawbi HA-H, Schadendorf D, Ascierto PA, Matamala L, Gutiérrez EC, et al. Relatlimab (RELA) plus nivolumab (NIVO) versus NIVO in first-line advanced melanoma: Primary phase III results from RELATIVITY-047 (CA224-047). *J Clin Oncol* [Internet]. 2021 May 20 [cited 2021 Jun 9];39(15_suppl):9503–9503. Available from: https://ascopubs.org/doi/10.1200/JCO.2021.39.15_suppl.9503
18. Marshall C. J. Specificity of receptor tyrosine kinase signaling: Transient versus sustained extracellular signal-regulated kinase activation. *Cell* [Internet]. 1995 Jan 27 [cited 2015 Jan 18];80(2):179–85. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/7834738>
19. Heidorn SJ, Milagre C, Whittaker S, Nourry A, Niculescu-Duvas I, Dhomen N, et al. Kinase-dead BRAF and oncogenic RAS cooperate to drive tumor progression through CRAF. *Cell* [Internet]. 2010 Jan 22 [cited 2014 Dec 2];140(2):209–21. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2872605&tool=pmcentrez&rendertype=abstract>
20. Siroy AE, Boland GM, Milton DR, Roszik J, Frankian S, Malke J, et al. Beyond BRAF(V600): Clinical Mutation Panel Testing by Next-Generation Sequencing in Advanced Melanoma. *J Invest Dermatol* [Internet]. 2015 Feb [cited 2015 Jan 18];135(2):508–15. Available from: <http://dx.doi.org/10.1038/jid.2014.366>
21. Chapman PB, Hauschild A, Robert C, Haanen JB, Ascierto P, Larkin J, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med* [Internet]. 2011 Jun 30;364(26):2507–16. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22646765>
22. Flaherty KT, Infante JR, Daud A, Gonzalez R, Kefford RF, Sosman J, et al. Combined BRAF and MEK inhibition in melanoma with BRAF V600 mutations. *N Engl J Med* [Internet]. 2012 Nov [cited 2013 Nov 7];367(18):1694–703. Available from: <http://www.nejm.org/doi/abs/10.1056/NEJMoa1210093>
23. Schadendorf D, Flaherty KT, Hersey P, Nathan PD, Garbe C, Milhem MM, et al. Overall survival (OS) update on METRIC (NCT01245062), a randomized phase 3 study to assess efficacy of trametinib (T) compared with chemotherapy (C) in patients (pts) with BRAFV600E/K mutation positive (+) advanced or metastatic melanoma (MM). *Pigment Cell Melanoma Res*. 2013;26(6):997.
24. Long G V., Stroyakovskiy D, Gogas H, Levchenko E, de Braud F, Larkin J, et al. Combined BRAF and MEK Inhibition versus BRAF Inhibition Alone in Melanoma. *N Engl J Med* [Internet]. 2014 Nov 13 [cited 2018 Jul 28];371(20):1877–88. Available from: <http://www.nejm.org/doi/10.1056/NEJMoa1406037>
25. Larkin J, Ascierto PA, Dréno B, Atkinson V, Liskay G, Maio M, et al. Combined Vemurafenib and Cobimetinib in BRAF-Mutated Melanoma. *N Engl J Med* [Internet]. 2014 Sep 29;371(20):1867–76. Available from:

<http://dx.doi.org/10.1056/NEJMoa1408868>

26. Dummer R, Ascierto PA, Gogas HJ, Arance A, Mandala M, Liskay G, et al. Encorafenib plus binimetinib versus vemurafenib or encorafenib in patients with BRAF -mutant melanoma (COLUMBUS): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* [Internet]. 2018 May [cited 2018 Jul 1];19(5):603–15. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S1470204518301426>
27. Long G V, Stroyakovskiy D, Gogas H, Levchenko E, de Braud F, Larkin J, et al. Dabrafenib and trametinib versus dabrafenib and placebo for Val600 BRAF-mutant melanoma: a multicentre, double-blind, phase 3 randomised controlled trial. *Lancet* [Internet]. 2015 May 29 [cited 2015 Jun 7]; Available from: <http://www.thelancet.com/article/S0140673615608984/fulltext>
28. Ascierto PA, Dummer R, Gogas HJ, Flaherty KT, Arance A, Mandala M, et al. Update on tolerability and overall survival in COLUMBUS: landmark analysis of a randomised phase 3 trial of encorafenib plus binimetinib vs vemurafenib or encorafenib in patients with BRAF V600–mutant melanoma. *Eur J Cancer* [Internet]. 2020 Feb 1 [cited 2021 Apr 3];126:33–44. Available from: <http://creativecommons.org/licenses/by-nc-nd/4.0/>
29. Dummer R, Flaherty K, Robert C, Arance AM, Groot JW de, Garbe C, et al. Five-year overall survival (OS) in COLUMBUS: A randomized phase 3 trial of encorafenib plus binimetinib versus vemurafenib or encorafenib in patients (pts) with BRAF V600-mutant melanoma. https://doi.org/10.1200/JCO20213915_suppl9507. 2021 May 28;39(15_suppl):9507–9507.
30. Center for Drug Evaluation and Research. Approved Drugs - Hematology/Oncology (Cancer) Approvals & Safety Notifications [Internet]. Center for Drug Evaluation and Research; [cited 2015 Jan 18]. Available from: <http://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm279174.htm>
31. Hauschild A, Grob J-J, Demidov L V, Jouary T, Gutzmer R, Millward M, et al. Dabrafenib in BRAF-mutated metastatic melanoma: a multicentre, open-label, phase 3 randomised controlled trial. *Lancet* [Internet]. 2012 Jul 28 [cited 2013 Nov 8];380(9839):358–65. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22735384>
32. Zhang C, Spevak W, Zhang Y, Burton EA, Ma Y, Habets G, et al. RAF inhibitors that evade paradoxical MAPK pathway activation. *Nature* [Internet]. 2015 Oct 22 [cited 2021 Apr 3];526(7574):583–6. Available from: <https://pubmed.ncbi.nlm.nih.gov/26466569/>
33. Wong DJ, Robert L, Atefi MS, Lassen A, Avarappatt G, Cerniglia M, et al. Antitumor activity of the ERK inhibitor SCH722984 against BRAF mutant, NRAS mutant and wild-type melanoma.
34. Ugurel S, Röhm J, Ascierto PA, Flaherty KT, Grob JJ, Hauschild A, et al. Survival of patients with advanced metastatic melanoma: the impact of novel therapies–update 2017. *Eur J Cancer* [Internet]. 2017 Sep [cited 2018 Jun 2];83:247–57. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28756137>
35. Liu C, Peng W, Xu C, Lou Y, Zhang M, Wargo JA, et al. BRAF inhibition increases tumor infiltration by T cells and enhances the antitumor activity of adoptive immunotherapy in mice. *Clin Cancer Res*. 2013;19(2):393–403.
36. Hu-Lieskovan S, Mok S, Homet Moreno B, Tsoi J, Robert L, Goedert L, et al. Improved antitumor activity of immunotherapy with BRAF and MEK inhibitors in BRAFV600E melanoma. *Sci Transl Med* [Internet]. 2015 Mar 18;7(279):279ra41–279ra41. Available from: <http://stm.sciencemag.org/content/7/279/279ra41.abstract>
37. Boni A, Cogdill AP, Dang P, Udayakumar D, Njauw CNJ, Sloss CM, et al. Selective BRAFV600E inhibition enhances T-cell recognition of melanoma without affecting lymphocyte function. *Cancer Res*. 2010;70(13):5213–9.

38. Khalili JS, Liu S, Rodriguez-Cruz TG, Whittington M, Wardell S, Liu C, et al. Oncogenic BRAF(V600E) promotes stromal cell-mediated immunosuppression via induction of interleukin-1 in melanoma. *Clin Cancer Res*. 2012;18(19):5329–40.
39. Luke JJ, Flaherty KT, Ribas A, Long G V. Targeted agents and immunotherapies: optimizing outcomes in melanoma. *Nat Rev Clin Oncol* [Internet]. 2017 Apr 4 [cited 2017 Jul 29];14(8):463–82. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28374786>
40. Frederick DT, Piris A, Cogdill AP, Cooper ZA, Lezcano C, Ferrone CR, et al. BRAF inhibition is associated with enhanced melanoma antigen expression and a more favorable tumor microenvironment in patients with metastatic melanoma. *Clin Cancer Res* [Internet]. 2013 Mar 1 [cited 2017 Apr 10];19(5):1225–31. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23307859>
41. Tumeu PC, Harview CL, Yearley JH, Shintaku IP, Taylor EJM, Robert L, et al. PD-1 blockade induces responses by inhibiting adaptive immune resistance. *Nature* [Internet]. 2014 Nov 26 [cited 2014 Nov 26];515(7528):568–71. Available from: <http://dx.doi.org/10.1038/nature13954>
42. First report of efficacy and safety from the phase II study SECOMBIT (SEquential COMBo Immuno and Targeted therapy study) | OncologyPRO [Internet]. [cited 2020 Dec 8]. Available from: <https://oncologypro.esmo.org/meeting-resources/esmo-virtual-congress-2020/first-report-of-efficacy-and-safety-from-the-phase-ii-study-secombit-sequential-combo-immuno-and-targeted-therapy-study>
43. Immunotherapy With Ipilimumab and Nivolumab Preceded or Not by a Targeted Therapy With Encorafenib and Binimetinib - Full Text View - ClinicalTrials.gov [Internet]. [cited 2020 Dec 8]. Available from: <https://clinicaltrials.gov/ct2/show/NCT03235245>
44. Spatalizumab plus dabrafenib and trametinib (Sparta-DabTram) in patients (pts) with previously untreated BRAF V600-mutant unresectable or metastat... | OncologyPRO [Internet]. [cited 2020 Oct 22]. Available from: <https://oncologypro.esmo.org/meeting-resources/esmo-virtual-congress-2020/spatalizumab-plus-dabrafenib-and-trametinib-sparta-dabtram-in-patients-pts-with-previously-untreated-braf-v600-mutant-unresectable-or-metastat>
45. Gutzmer R, Stroyakovskiy D, Gogas H, Robert C, Lewis K, Protsenko S, et al. Atezolizumab, vemurafenib, and cobimetinib as first-line treatment for unresectable advanced BRAFV600 mutation-positive melanoma (IMspire150): primary analysis of the randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* [Internet]. 2020 Jun 13 [cited 2020 Jun 17];395(10240):1835–44. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S014067362030934X>
46. Zhang J, Cunningham JJ, Brown JS, Gatenby RA. Integrating evolutionary dynamics into treatment of metastatic castrate-resistant prostate cancer. *Nat Commun* [Internet]. 2017 Dec 28 [cited 2018 Jul 13];8(1):1816. Available from: <http://www.nature.com/articles/s41467-017-01968-5>
47. West JB, Dinh MN, Brown JS, Zhang J, Anderson AR, Gatenby RA. Multidrug Cancer Therapy in Metastatic Castrate-Resistant Prostate Cancer: An Evolution-Based Strategy. *Clin Cancer Res* [Internet]. 2019 Jul 15 [cited 2019 Aug 30];25(14):4413–21. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30992299>
48. Gatenby RA, Silva AS, Gillies RJ, Frieden BR. Adaptive Therapy. *Cancer Res* [Internet]. 2009 Jun 1 [cited 2019 Aug 31];69(11):4894–903. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19487300>
49. Enriquez-Navas PM, Wojtkowiak JW, Gatenby RA. Application of Evolutionary Principles to Cancer Therapy.

- Cancer Res [Internet]. 2015 Nov 15 [cited 2019 Aug 31];75(22):4675–80. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26527288>
50. Strobl MAR, West J, Viossat Y, Damaghi M, Robertson-Tessi M, Brown JS, et al. Turnover modulates the need for a cost of resistance in adaptive therapy. *Cancer Res*. 2021 Feb 15;81(4):1135–47.
 51. Hansen E, Woods RJ, Read AF. How to Use a Chemotherapeutic Agent When Resistance to It Threatens the Patient. *PLoS Biol*. 2017 Feb 9;15(2).
 52. Bacevic K, Noble R, Soffar A, Wael Ammar O, Boszonyik B, Prieto S, et al. Spatial competition constrains resistance to targeted cancer therapy. *Nat Commun* [Internet]. 2017 Dec 8 [cited 2018 Jul 13];8(1):1995. Available from: <http://www.nature.com/articles/s41467-017-01516-1>
 53. Viossat Y, Noble R. The logic of containing tumors. *bioRxiv*. bioRxiv; 2020.
 54. Kim E, Brown JS, Eroglu Z, Anderson ARA. Adaptive therapy for metastatic melanoma: Predictions from patient calibrated mathematical models. *Cancers (Basel)* [Internet]. 2021 Feb 2 [cited 2021 May 10];13(4):1–15. Available from: <https://doi.org/10.3390/cancers13040823>
 55. Gallaher JA, Enriquez-Navas PM, Luddy KA, Gatenby RA, Anderson ARA. Spatial heterogeneity and evolutionary dynamics modulate time to recurrence in continuous and adaptive cancer therapies. *Cancer Res*. 2018 Apr 15;78(8):2127–39.
 56. Maley CC, Aktipis A, Graham TA, Sottoriva A, Boddy AM, Janiszewska M, et al. Classifying the evolutionary and ecological features of neoplasms. *Nat Rev Cancer* [Internet]. 2017 Oct 15 [cited 2019 Sep 2];17(10):605–19. Available from: <http://www.nature.com/articles/nrc.2017.69>
 57. Siravegna G, Mussolin B, Buscarino M, Corti G, Cassingena A, Crisafulli G, et al. Clonal evolution and resistance to EGFR blockade in the blood of colorectal cancer patients. *Nat Med* [Internet]. 2015 Jul 1 [cited 2019 Aug 30];21(7):795–801. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26030179>
 58. Smalley I, Kim E, Li J, Spence P, Wyatt CJ, Eroglu Z, et al. Leveraging transcriptional dynamics to improve BRAF inhibitor responses in melanoma. *EBioMedicine*. 2019 Oct;
 59. Müller J, Krijgsman O, Tsoi J, Robert L, Hugo W, Song C, et al. Low MITF/AXL ratio predicts early resistance to multiple targeted drugs in melanoma. *Nat Commun* [Internet]. 2014 Dec 15 [cited 2019 Apr 7];5(1):5712. Available from: <http://www.nature.com/articles/ncomms6712>
 60. Hoek KS, Goding CR. Cancer stem cells versus phenotype-switching in melanoma. *Pigment Cell Melanoma Res* [Internet]. 2010 Dec 1 [cited 2018 Sep 2];23(6):746–59. Available from: <http://doi.wiley.com/10.1111/j.1755-148X.2010.00757.x>
 61. Garraway LA, Widlund HR, Rubin MA, Getz G, Berger AJ, Ramaswamy S, et al. Integrative genomic analyses identify MITF as a lineage survival oncogene amplified in malignant melanoma. *Nature* [Internet]. 2005 Jul 7 [cited 2017 Sep 12];436(7047):117–22. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16001072>
 62. Valpione S, Carlino MS, Mangana J, Mooradian MJ, McArthur G, Schadendorf D, et al. Rechallenge with BRAF-directed treatment in metastatic melanoma: A multi-institutional retrospective study. *Eur J Cancer* [Internet]. 2018 Mar 1 [cited 2018 Jul 28];91:116–24. Available from: <https://www.sciencedirect.com/science/article/pii/S0959804917314867>
 63. Schreuer M, Jansen Y, Planken S, Chevolet I, Seremet T, Kruse V, et al. Combination of dabrafenib plus

- trametinib for BRAF and MEK inhibitor pretreated patients with advanced BRAF V600 -mutant melanoma: an open-label, single arm, dual-centre, phase 2 clinical trial. *Lancet Oncol* [Internet]. 2017 Apr [cited 2019 Aug 31];18(4):464–72. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28268064>
64. Tietze JK, Forschner A, Loquai C, Mitzel-Rink H, Zimmer L, Meiss F, et al. The efficacy of re-challenge with BRAF inhibitors after previous progression to BRAF inhibitors in melanoma: A retrospective multicenter study. *Oncotarget*. 2018 Sep 1;9(76):34336–46.
 65. Gebhardt C, Ascierto P, Atkinson V, Corrie P, Dummer R, Schadendorf D. The concepts of rechallenge and retreatment in melanoma: A proposal for consensus definitions. *Eur J Cancer*. 2020 Oct 1;138:68–76.
 66. Wan JCM, Massie C, Garcia-Corbacho J, Mouliere F, Brenton JD, Caldas C, et al. Liquid biopsies come of age: Towards implementation of circulating tumour DNA. Vol. 17, *Nature Reviews Cancer*. Nature Publishing Group; 2017. p. 223–38.
 67. Schwarzenbach H, Hoon DSB, Pantel K. Cell-free nucleic acids as biomarkers in cancer patients. *Nat Rev Cancer* [Internet]. 2011 Jun 12 [cited 2014 Jul 9];11(6):426–37. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21562580>
 68. Sorenson G, Pribish D, Valone F, Memoli V, Bzik D, Yao S. Soluble normal and mutated DNA sequences from single-copy genes in human blood. *Cancer Epidemiol Biomarkers Prev* [Internet]. 1994 Jan 1 [cited 2014 Dec 14];3(1):67–71. Available from: <http://cebp.aacrjournals.org/content/3/1/67.abstract>
 69. Choi J-J, Reich CF, Pisetsky DS. The role of macrophages in the in vitro generation of extracellular DNA from apoptotic and necrotic cells. *Immunology* [Internet]. 2005 May [cited 2014 Dec 16];115(1):55–62. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1782131&tool=pmcentrez&rendertype=abstract>
 70. Smith B, Selby P, Southgate J, Pittman K, Bradley C, Blair GE. Detection of melanoma cells in peripheral blood by means of reverse transcriptase and polymerase chain reaction. *Lancet*. 1991 Nov 16;338(8777):1227–9.
 71. Murtaza M, Dawson S-J, Tsui DWY, Gale D, Forshew T, Piskorz AM, et al. Non-invasive analysis of acquired resistance to cancer therapy by sequencing of plasma DNA. *Nature* [Internet]. 2013 May 2 [cited 2014 Jul 9];497(7447):108–12. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23563269>
 72. Syeda MM, Wiggins JM, Corless BC, Long G V., Flaherty KT, Schadendorf D, et al. Circulating tumour DNA in patients with advanced melanoma treated with dabrafenib or dabrafenib plus trametinib: a clinical validation study. *Lancet Oncol*. 2021;22(3):370–80.
 73. Diehl F, Schmidt K, Choti MA, Romans K, Goodman S, Li M, et al. Circulating mutant DNA to assess tumor dynamics. *Nat Med* [Internet]. 2008 Sep [cited 2014 Oct 9];14(9):985–90. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2820391&tool=pmcentrez&rendertype=abstract>
 74. Dawson S-J, Tsui DWYY, Murtaza M, Biggs H, Rueda OM, Chin S-F, et al. Analysis of circulating tumor DNA to monitor metastatic breast cancer. *N Engl J Med* [Internet]. 2013 Mar 28 [cited 2014 Jul 9];368(13):1199–209. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23484797>
 75. Santiago-Walker A, Gagnon R, Mazumdar J, Casey M, Long G V., Schadendorf D, et al. Correlation of BRAF Mutation Status in Circulating-Free DNA and Tumor and Association with Clinical Outcome across Four BRAFi and MEKi Clinical Trials. *Clin Cancer Res*. 2016;22(3).
 76. Santiago-Walker A, Gagnon R, Mazumdar J, Casey M, Long G V., Schadendorf D, et al. Correlation of BRAF

- mutation status in circulating-free DNA and tumor and association with clinical outcome across four BRAFi and MEKi clinical trials. *Clin Cancer Res* [Internet]. 2015;1–9. Available from: <http://clincancerres.aacrjournals.org/cgi/doi/10.1158/1078-0432.CCR-15-0321>
77. Chang GA, Tadeipalli JS, Shao Y, Zhang Y, Weiss S, Robinson E, et al. Sensitivity of plasma BRAF mutant and NRAS mutant cell-free DNA assays to detect metastatic melanoma in patients with low RECIST scores and non-RECIST disease progression. *Mol Oncol*. 2016 Jan;10(1):157–65.
 78. Schreuer M, Meersseman G, Van Den Herrewegen S, Jansen Y, Chevolet I, Bott A, et al. Quantitative assessment of BRAF V600 mutant circulating cell-free tumor DNA as a tool for therapeutic monitoring in metastatic melanoma patients treated with BRAF/MEK inhibitors. *J Transl Med*. 2016 Dec;14(1):95.
 79. Haselmann V, Gebhardt C, Brechtel I, Duda A, Czerwinski C, Sucker A, et al. Liquid Profiling of Circulating Tumor DNA in Plasma of Melanoma Patients for Companion Diagnostics and Monitoring of BRAF Inhibitor Therapy. *Clin Chem* [Internet]. 2018 May [cited 2018 May 17];64(5):830–42. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29483107>
 80. Braune J, Keller L, Schiller F, Graf E, Rafei-Shamsabadi D, Wehrle J, et al. Circulating Tumor DNA Allows Early Treatment Monitoring in BRAF- and NRAS-Mutant Malignant Melanoma. *JCO Precis Oncol*. 2020 Sep;(4):20–31.
 81. Girotti MR, Gremel G, Lee R, Galvani E, Rothwell D, Viros A, et al. Application of sequencing, liquid biopsies, and patient-derived xenografts for personalized medicine in melanoma. *Cancer Discov* [Internet]. 2016; Available from: <http://www.scopus.com/inward/record.url?eid=2-s2.0-84960075353&partnerID=MN8TOARS>
 82. Gray ES, Rizos H, Reid AL, Boyd SC, Pereira MR, Lo J, et al. Circulating tumor DNA to monitor treatment response and detect acquired resistance in patients with metastatic melanoma. *Oncotarget* [Internet]. 2015 Dec 8 [cited 2018 May 19];6(39):42008–18. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26524482>
 83. Das Thakur M, Salangsang F, Landman AS, Sellers WR, Pryer NK, Levesque MP, et al. Modelling vemurafenib resistance in melanoma reveals a strategy to forestall drug resistance. *Nature*. 2013 Feb;494(7436):251–5.
 84. Hong A, Moriceau G, Sun L, Lomeli S, Piva M, Damoiseaux R, et al. Exploiting Drug Addiction Mechanisms to Select against MAPKi-Resistant Melanoma. *Cancer Discov* [Internet]. 2018 Jan 18 [cited 2018 Jun 10];8(1):74–93. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28923912>
 85. Kong X, Kuilman T, Shahrabi A, Boshuizen J, Kemper K, Song J-Y, et al. Cancer drug addiction is relayed by an ERK2-dependent phenotype switch. *Nature* [Internet]. 2017 Oct 4 [cited 2017 Nov 19];550(7675):nature24037. Available from: <http://www.nature.com/doi/10.1038/nature24037>
 86. Lee RJ, Marais R. Tumours addicted to drugs are vulnerable. *Nature* [Internet]. 2017 Oct 4 [cited 2018 Jul 19];550(7675):192–3. Available from: <http://www.nature.com/doi/10.1038/nature24148>
 87. Algazi AP, Othus M, Daud AI, Lo RS, Mehnert JM, Truong TG, et al. Continuous versus intermittent BRAF and MEK inhibition in patients with BRAF-mutated melanoma: a randomized phase 2 trial. *Nat Med* [Internet]. 2020 Oct 1 [cited 2021 May 9];26(10):1564–8. Available from: <https://doi.org/10.1038/s41591-020-1060-8>
 88. Gremel G, Lee RJ, Girotti MR, Mandal AK, Valpione S, Garner G, et al. Distinct subclonal tumour responses to therapy revealed by circulating cell-free DNA. *Ann Oncol* [Internet]. 2016 Oct [cited 2017 Aug 3];27(10):1959–65. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27502704>

89. Lee RJ, Gremel G, Marshall A, Myers KA, Fisher N, Dunn JA, et al. Circulating tumor DNA predicts survival in patients with resected high-risk stage II/III melanoma. *Ann Oncol* [Internet]. 2018 Feb 1 [cited 2018 Feb 18];29(2):490–6. Available from: <https://academic.oup.com/annonc/article/29/2/490/4589598>
90. Tan L, Sandhu S, Lee RJ, Li J, Callahan J, Ftouni S, et al. Prediction and monitoring of relapse in stage III melanoma using circulating tumor DNA. *Ann Oncol* [Internet]. 2019 Feb 6 [cited 2019 May 18]; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30838379>
91. Girotti MR, Gremel G, Lee R, Galvani E, Rothwell D, Viros A, et al. Application of Sequencing, Liquid Biopsies, and Patient-Derived Xenografts for Personalized Medicine in Melanoma. *Cancer Discov*. 2016 Mar;6(3):286–99.
92. Wong SQ, Raleigh JM, Callahan J, Vergara IA, Ftouni S, Hatzimihalis A, et al. Circulating Tumor DNA Analysis and Functional Imaging Provide Complementary Approaches for Comprehensive Disease Monitoring in Metastatic Melanoma. *JCO Precis Oncol* [Internet]. 2017 Jul 27 [cited 2018 Jul 2];(1):1–14. Available from: <http://ascopubs.org/doi/10.1200/PO.16.00009>
93. Chang-Hao Tsao S, Weiss J, Hudson C, Christophi C, Cebon J, Behren A, et al. Monitoring response to therapy in melanoma by quantifying circulating tumour DNA with droplet digital PCR for BRAF and NRAS mutations. *Sci Rep* [Internet]. 2015 Jan 22 [cited 2015 Jun 27];5(1):11198. Available from: <http://www.nature.com/articles/srep11198>
94. Koelblinger P, Thuerigen O, Dummer R. Development of encorafenib for BRAF-mutated advanced melanoma. Vol. 30, *Current Opinion in Oncology*. Lippincott Williams and Wilkins; 2018. p. 125–33.
95. Robert C, Grob JJ, Stroyakovskiy D, Karaszewska B, Hauschild A, Levchenko E, et al. Five-year outcomes with dabrafenib plus trametinib in metastatic melanoma. *N Engl J Med*. 2019 Aug 15;381(7):626–36.
96. Marsavela G, Johansson PA, Pereira MR, McEvoy AC, Reid AL, Robinson C, et al. The prognostic impact of circulating tumour dna in melanoma patients treated with systemic therapies—beyond braf mutant detection. *Cancers (Basel)*. 2020;12(12):1–13.
97. Wong SQ, Raleigh JM, Callahan J, Vergara IA, Ftouni S, Hatzimihalis A, et al. Circulating Tumor DNA Analysis and Functional Imaging Provide Complementary Approaches for Comprehensive Disease Monitoring in Metastatic Melanoma. *JCO Precis Oncol*. 2017 Nov 27;(1):1–14.

18 DOCUMENTS SUPPLEMENTARY TO THE PROTOCOL

Documents referenced within the protocol are separately maintained and version controlled. Any of the supplementary documents subject to MHRA or Research Ethics Committee review are submitted as separate version controlled documents.

19 APPENDICES

Appendix A: Cockcroft-Gault formula:

Cockcroft Gault Formula

$$C_{Cr} = \{((140 - \text{age}) \times \text{weight}) / (72 \times S_{Cr})\} \times 0.85 \text{ if female}$$

Appendix B: Contraceptive Guidance:

Woman of Childbearing Potential (WOCBP)

A woman is considered fertile following menarche and until becoming post-menopausal unless permanently sterile (see below).

Women in the following categories are not considered WOCBP:

- Premenarchal
- Premenopausal female with 1 of the following:
 - Documented hysterectomy
 - Documented bilateral salpingectomy
 - Documented bilateral oophorectomy

Note: Documentation can come from the site personnel's review of the participant's medical records, medical examination, or medical history interview.

- Postmenopausal female
 - A postmenopausal state is defined as no menses for 12 months without an alternative medical cause.
 - A high follicle stimulating hormone (FSH) level in the postmenopausal range may be used to confirm a postmenopausal state in women not using hormonal contraception or hormone replacement therapy (HRT). However, in the absence of 12 months of amenorrhea, confirmation with two FSH measurements in the postmenopausal range is required.
 - Females on HRT and whose menopausal status is in doubt will be required to use one of the non-hormonal highly effective contraception methods if they wish to continue their HRT during the study. Otherwise, they must discontinue HRT to allow confirmation of postmenopausal status before study enrolment.

Contraception Requirements:

Male Participants:

Male participants with partners who are WOCBP are eligible to participate if they agree to one of the following during the protocol defined time frame outlined in Section [7.57-5](#)

- Be abstinent from penile-vaginal intercourse as their usual and preferred lifestyle (abstinent on a long term and persistent basis) and agree to remain abstinent

- Use a male condom plus partner use of a contraceptive method with a failure rate of <1% per year as described in Table 5 when having penile-vaginal intercourse with a WOCBP who is not currently pregnant.
- Note: Men with a pregnant or breastfeeding partner must agree to remain abstinent from penile-vaginal intercourse or use a male condom during each episode of penile penetration.

Female Participants:

Female participants of childbearing potential are eligible to participate if they agree to use a highly effective method of contraception consistently and correctly as described in Table 5 during the protocol-defined time frame outlined in Section 7.5.

Table 5 - Highly Effective Contraception Methods:

Highly Effective Contraceptive Methods That Are User Dependent ^a <i>Failure rate of <1% per year when used consistently and correctly.</i>
• Combined (oestrogen- and progestogen- containing) hormonal contraception ^{b, c} ○ Oral ○ Intravaginal ○ Transdermal ○ Injectable
• Progestogen-only hormonal contraception ^{b, c} ○ Oral ○ Injectable
Highly Effective Methods That Have Low User Dependency <i>Failure rate of <1% per year when used consistently and correctly.</i>
• Progestogen- only contraceptive implant ^{b, c} • Intrauterine hormone-releasing system (IUS) ^b • Intrauterine device (IUD) • Bilateral tubal occlusion

- Vasectomized partner

A vasectomized partner is a highly effective contraception method provided that the partner is the sole male sexual partner of the WOCBP and the absence of sperm has been confirmed. If not, an additional highly effective method of contraception should be used.

- Sexual abstinence

Sexual abstinence is considered a highly effective method only if defined as refraining from heterosexual intercourse during the entire period of risk associated with the study treatment. The reliability of sexual abstinence needs to be evaluated in relation to the duration of the study and the preferred and usual lifestyle of the participant.

Notes:

Use should be consistent with local regulations regarding the use of contraceptive methods for participants of clinical studies.

^a Typical use failure rates are lower than perfect-use failure rates (i.e. when used consistently and correctly).

^b If hormonal contraception efficacy is potentially decreased due to interaction with study treatment, condoms must be used in addition to the hormonal contraception during the treatment period and for at least 5 months, corresponding to time needed to eliminate study treatment after the last dose.

^c If locally required, in accordance with Clinical Trial Facilitation Group guidelines, acceptable hormonal contraceptives are limited to those which inhibit ovulation.

Appendix C: Clinical and Scientific contributors

Name	Affiliations	Contribution to protocol
Prof Trevor Graham	Barts Cancer Institute	Study design and input regarding tumour evolution and mathematical analyses.
Dr Rebecca Lee	The University of Manchester, The Christie NHS Foundation Trust	Clinical input, study design, protocol review.
Prof Paul Lorigan	The Christie NHS Foundation Trust	Lead on clinical input, study design, protocol review and approval.
Alkesh Patel	The Christie NHS Foundation Trust	Lead on Pharmaceutical aspects of the study and pharmacy approval.
Dr Dominic Rothwell	CRUK National Biomarker Centre	Development of the ctDNA assays to GCP standards.
Mr Nigel Smith	CRUK National Biomarker Centre	Development of the ctDNA assays to GCP standards.
Prof Andrea Sottoriva	Human Technopole	Study design and input regarding tumour evolution and mathematical analyses.
Dr Benjamin Werner	Barts Cancer Institute	Study design and input regarding tumour evolution and mathematical analyses.

Appendix D: Scans and RECIST criteria:

Scan modalities by anatomy

Colour legend

Green = Required / Preferred	Amber = Acceptable alternative	Red = Not applicable
------------------------------	--------------------------------	----------------------

Anatomy	Modality			
	PET-CT	CT only	MRI	Ultrasound
Research Biopsy	N/A	Only if clinically indicated as per local practice.	N/A	Only if clinically indicated as per local practice.
Limbs (only if disease present)	Head to toe, with contrast if no known allergy.	N/A	N/A	N/A
Brain (if limb disease)	By default, if limb disease.	N/A	With contrast if no known allergy.	N/A
Brain (if no limb disease)	N/A	With contrast if no known allergy. Acceptable alternative to MRI, as per local practice	With contrast if no known allergy	N/A
Neck (if indicated)	By default, if limb disease.	With contrast if no known allergy.	N/A	N/A
Thorax	By default, if limb disease.	With contrast if no known allergy.	N/A	N/A
Abdomen & Pelvis	By default, if limb disease.	With contrast if no known allergy. If known allergy to contrast, MRI can be used as an alternative, as per local practice	Acceptable alternative to CT if known allergy to contrast, as per local practice.	N/A

For participants who have measurable disease at baseline, the table below provides a summary of the overall time point response: participants with target (+/- non-target) disease.

Target lesions	Non-Target lesions	New lesions	Overall response
CR	CR	No	CR
CR	Non-CR/non-PD	No	PR
CR	Not evaluated	No	PR
PR	Non-PD or not all evaluated	No	PR
SD	Non-PD or not all evaluated	No	SD
Not all evaluated	Non-PD	No	NE
PD	Any	Yes or No	PD
Any	PD	Yes or No	PD
Any	Any	Yes	PD

Legend:

CR = Complete Response,

NE = Not Evaluable,

PD = Progressive Disease,

PR = Partial Response,

SD = Stable Disease.

Missing Assessments and Not Evaluable Designation:

When no imaging/measurement is done at all at a particular time point, the participant is not evaluable (NE) at that time point. If only a subset of lesion measurements are made at an assessment, usually the case is also considered NE at that time point, unless a convincing argument can be made that the contribution of the individual missing lesion(s) would not change the assigned time point response.

For full RECIST version 1.1 criteria please refer to:

https://ctep.cancer.gov/protocoldevelopment/docs/recist_guideline.pdf

Appendix E: ECOG performance status*:

Grade	ECOG status
0	Fully active, able to carry on all pre-disease performance without restriction
1	Restricted in physically strenuous activity but ambulatory and able to carry out work of a light or sedentary nature, e.g., light house work, office work
2	Ambulatory and capable of all selfcare but unable to carry out any work activities. Up and about more than 50% of waking hours
3	Capable of only limited selfcare, confined to bed or chair more than 50% of waking hours
4	Completely disabled. Cannot carry on any selfcare. Totally confined to bed or chair
5	Dead

*Oken M, Creech R, Tormey D, et al. Toxicity and response criteria of the Eastern Cooperative Oncology Group
 Am J Clin Oncol. 1982;5:649-655

Appendix F: Amino acids and abbreviations:

Amino acid	Abbreviation (1 letter)	Abbreviation (3 letters)
Aspartic acid	D	Asp
Glutamic acid	E	Glu
Lysine	K	Lys
Arginine	R	Arg