An international multicentre randomised trial comparing general anaesthetic versus local anaesthetic for carotid surgery

Submission date Recruitment status Prospectively registered 12/09/2002 No longer recruiting [X] Protocol Statistical analysis plan Registration date Overall study status 12/09/2002 Completed [X] Results Individual participant data Last Edited Condition category 31/12/2008 Surgery

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Ms Bridget Colam

Contact details

Neurosciences Trials Unit Bramwell Dott Building Western General Hospital Crewe Road Edinburgh United Kingdom EH4 2XU +44 (0)131 537 2930 gala@skull.dcn.ed.ac.uk

Additional identifiers

Protocol serial number 2268/1979

Study information

Scientific Title

Acronym

GALA

Study objectives

A prospective, randomised trial of local versus general anaesthesia for carotid endarterectomy is proposed to determine whether the type of anaesthesia does indeed influence peri-operative morbidity and mortality, quality of life and long term outcome in terms of stroke-free survival.

Please note that as of 29/05/2008 this record was updated to include a longer recruitment period. All changes can be found under the relevant fields, with the update date of 29/05/2008. The anticipated end date of this trial was extended to 30/04/2008. The previous anticipated end date of this trial was 31/12/2006. Randomisation stopped on 31/10/2007 and the presentation of results is expected on 06/09/08.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received from the Northern and Yorkshire MREC on the 28th August 1998 (pilot phase) and 22nd April 2003 (main phase).

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Carotid stenosis (at risk from stroke)

Interventions

Anaesthetic for surgery - local versus general. Randomisation will be by telephone to the trial office. Data will be collected at baseline (prior to randomisation) and this will include demographic details, known risk factors, diagnostic procedures and findings, and indications for surgery. The follow-up data will be collected when the patient is discharged from acute care, and operative details and any early complications will be recorded. At 1 month post-surgery, a stroke physician, blind to the type of anaesthesia, will review the patient. Also the patient will be asked to complete a health related quality of life questionnaire including the widely used EuroQol and Short Form 36, plus a carotid endarterectomy (CEA) specific questionnaire designed to capture patient satisfaction levels with the method of anaesthesia. Length of time spent in the recovery room, intensive care unit (ICU) and high dependency unit (HDU), and length of overall acute hospital stay will be recorded. Annual follow up will be by post to the family doctor and to the patient, seeking details of any strokes.

Intervention Type

Procedure/Surgery

Phase

Not Specified

Primary outcome(s)

The proportion of patients alive, stroke free (including retinal infarction) and without myocardial infarction (MI) 30 days post-surgery.

Key secondary outcome(s))

Proportion alive and stroke free at one year and in the longer term, a comparison of health related quality of life at 30 days and any surgical adverse events, re-operation and re-admission rates, the relative cost of the two methods of anaesthesia, length of stay and intensive and high dependency bed occupancy.

Completion date

30/04/2008

Eligibility

Key inclusion criteria

The main criterion for entry into this study is that, in the opinion of the responsible clinician, the patient requiring an endarterectomy is suitable for either local or general anaesthesia, and there is no clear indication for either type. All patients with either symptomatic or asymptomatic internal carotid stenosis for whom surgery is advised are eligible. There is no upper age limit.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

- 1. Failure to obtain informed consent
- 2. Patient unable to co-operate with awake testing during local anaesthesia
- 3. Patient considered unfit for general anaesthesia
- 4. Patient considered unfit for local anaesthesia
- 5. Patient requiring simultaneous bilateral carotid endarterectomy
- 6. Carotid endarterectomy combined with another operative artery bypass surgery
- 7. Patient has been randomised into the trial previously

Date of first enrolment

01/06/1999

Date of final enrolment

30/04/2008

Locations

Countries of recruitment United Kingdom
Scotland
Australia
Austria
Belarus
China
Croatia
Czech Republic
Estonia
Georgia
Germany
Greece
Hungary
Israel
Italy
Latvia
Netherlands
Poland
Portugal
Saudi Arabia
Slovakia
Spain

Sweden

Türkiye

Ukraine

Study participating centre Neurosciences Trials Unit Edinburgh United Kingdom EH4 2XU

Sponsor information

Organisation

The Health Foundation (UK)

ROR

https://ror.org/02bzj4420

Funder(s)

Funder type

Charity

Funder Name

The Health Foundation (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	20/12/2008		Yes	No
Protocol article	Protocol	21/05/2008		Yes	No
	Study website				

<u>Study website</u> 11/11/2025 11/11/2025 No Yes