# Gastric bypass, adjustable gastric banding or sleeve gastrectomy surgery to treat severe and complex obesity

Submission date 01/09/2011	<b>Recruitment status</b> No longer recruiting	[X] Prospectively registered			
		[X] Protocol			
Registration date	Overall study status	Statistical analysis plan			
05/09/2011	Completed	[X] Results			
Last Edited 04/04/2025	<b>Condition category</b> Nutritional, Metabolic, Endocrine	Individual participant data			

# Plain English summary of protocol

#### Background and study aims

Obesity is an increasing health problem in the UK and is predicted to get worse. The three recognised operations in bariatric surgery are laparoscopic adjustable gastric banding (small inner tube placed around the stomach to reduce its capacity), laparoscopic gastric bypass (operation to make the stomach smaller and the digestive system shorter) and sleeve gastrectomy (operation to make the stomach smaller). This study aims to compare these three types of operation to see which one is better.

#### Who can participate?

The By-Band-Sleeve study is in follow up and is no longer recruiting new participants (added 24 /08/2022). Obese males and females over the age of 18 years who are referred for obesity surgery (at the recruiting centres) under current government guidelines can participate in the study. Participants should have a body mass index (BMI) of 40kg/m2 or more, or a BMI of 35 kg /m2 to 40 kg/m2 with other risk factors (e.g. type 2 diabetes), that could improve with weight loss.

# What does the study involve?

Participants are randomly allocated to be treated with either gastric band surgery, gastric bypass surgery or sleeve gastrectomy surgery. Participants are also asked to complete a series of questionnaires at regular intervals up to three years after the operation about their quality of life, and some participants are invited to be interviewed about their experiences of treatment decisions. Researchers also ask participants to provide two blood samples in addition to the samples they would give as part of their normal care for future research into obesity.

# What are the possible benefits and risks of participating?

There is no direct benefit for people enrolling in the study, although some people find that interviews help them talk through their situation and this is comforting. There should be no additional physical risk to you because all three operations offered are standard treatments for severe and complex obesity. Where is the study run from? University of Bristol (UK).

When is the study starting and how long is it expected to run for? January 2012 to September 2022

Who is funding the study? The National Institute of Health Research, Health Technology Assessment Programme (UK)

Who is the main contact? Dr Graziella Mazza g.mazza@bristol.ac.uk

**Study website** www.by-band-sleeve.bristol.ac.uk

# **Contact information**

**Type(s)** Scientific

**Contact name** Dr Graziella Mazza

# **Contact details**

Clinical Trials and Evaluation Unit School of Clinical Sciences University of Bristol Bristol Royal Infirmary Marlborough Street Bristol United Kingdom BS2 8HW

# Additional identifiers

EudraCT/CTIS number

**IRAS number** 

ClinicalTrials.gov number NCT02841527

Secondary identifying numbers HTA 09/127/53, 1279

# Study information

Scientific Title

Gastric bypass, adjustable gastric banding or sleeve gastrectomy surgery to treat severe and complex obesity: a multi-centre randomised controlled trial

### Acronym

**By-Band-Sleeve** 

# **Study objectives**

Current hypothesis as of 23/09/2015:

The By-Band-Sleeve study will compare the effectiveness, cost-effectiveness and acceptability of Band versus Bypass versus Sleeve surgery for treatment of severe and complex obesity. We will test the following joint hypotheses:

Bypass is non-inferior to Band with respect to excess weight loss of more than 50% at three years and that Bypass is superior to Band with respect to HRQOL at three years
Sleeve is non-inferior to Band with respect to excess weight loss of more than 50% at three years and that Sleeve is superior to Band with respect to HRQOL at three years
Sleeve is non-inferior to Bypass with respect to excess weight loss of more than 50% at three

years and that Sleeve is superior to Bypass with respect to HRQOL at three years

In the primary analysis both outcomes will be considered collectively, i.e. both hypotheses must be supported to conclude that Bypass is more effective than Band, or that Sleeve is more effective than Band or Sleeve is more effective than Bypass.

Specific objectives are to estimate:

1. The difference between groups in the proportion of patients achieving >50% excess weight loss at three years

2. The difference between groups in their average EQ-5D-5L health state score at three years 3. The difference between groups with respect to a range of secondary outcomes including generic, disease specific and gastro-intestinal symptom specific measures of HRQOL, adverse events, and resolution of co-morbidities; to explore, in a sub-sample, patients' experiences of management, outcome and eating behaviour change

4. The cost effectiveness of Band, Bypass and Sleeve

Previous hypothesis:

The joint hypotheses is that gastric BYpass is non-inferior to gastric BAND with respect to excess weight loss of more than 50% at three years and that BYpass is superior to BAND with respect to quality of life at three years. In the primary analysis both outcomes will be considered collectively, i.e. both hypotheses must be supported to conclude that BYPASS is more effective than BAND.

Specific objectives are to estimate:

1. The difference between groups in the proportion of patients achieving >50% excess weight loss at three years

2. The difference between groups in their average EQ-5D health state score at three years 3. The difference between groups with respect to a range of secondary outcomes including generic, disease specific and gastro-intestinal symptom specific measures of HRQOL, adverse events, and resolution of co-morbidities; to explore, in a sub-sample, patients' experiences of management, outcome and eating behaviour change.

4. The cost effectiveness of BAND and BYPASS

# Ethics approval required

Old ethics approval format

**Ethics approval(s)** South West Frenchay Research Ethics Committee, 14/10/2011

**Study design** Multi-centre randomised controlled trial

**Primary study design** Interventional

**Secondary study design** Randomised controlled trial

**Study setting(s)** Hospital

**Study type(s)** Treatment

# Participant information sheet

Not available in web format, please use the contact details to request a patient information sheet

Health condition(s) or problem(s) studied Obesity

# Interventions

Current interventions as of 23/09/2015:

All three surgical procedures (Band, Bypass and Sleeve) will be carried out laparoscopically. The methods used to create a pneumoperitoneum, the placement of the laparoscopic ports, and retractors are at the discretion of the surgeon. The type of instruments used is also at the surgeon's discretion. Undertaking a hiatal hernia (<5cm) repair and cholecystectomy are permitted but not compulsory. An apronectomy is prohibited at the time of surgery. Placement of drains is optional.

1. Laparoscopic adjustable gastric banding (Band surgery)

The type and size of adjustable gastric band is at the discretion of the surgeon. It is mandatory to i) dissect the lesser curve using the 'Pars flaccida' technique, ii) use non-absorbable gastro-gastric tunnelling sutures and iii) to fix the adjustable port to the anterior abdominal wall.

2. Laparoscopic gastric bypass surgery (Bypass surgery)

Methods used to create the biliary and gastric limbs are flexible, although upper limits of 75 cm and 150 cm are recommended for the biliary and gastric limbs, respectively. Routing of the Roux limb (antecolic or retrocolic) is flexible. The pouch can be created according to the surgeon's usual practice, except that a horizontal gastric pouch that includes fundus is prohibited. Use of a bougie is optional. Anastomoses can be performed as the surgeon chooses (e.g. stapled or sutured, single or double layer). Testing integrity of the anastomoses, and the closure of the mesenteric defects, is optional.

### 3. Laparoscopic sleeve gastrectomy surgery (Sleeve surgery)

It is mandated to visualise the left crus after dissection of the fundus. The type of bougie used is flexible although should be between 32 and 40 Fr. The type of stapler used is flexible and the use of additional sutures, clips, reinforcement of the staple line is at the discretion of the surgeon. Testing the integrity of the staple line is optional.

## 4. Concomitant interventions

Procedures will be carried out under general anaesthesia. All patients will receive peri-operative antibiotics and thromboprophylaxis in accordance with local policy. The use of nasogastric tubes, central and arterial lines, and urinary catheters is optional. After surgery, oral intake will be commenced according to local policy. The day of discharge will be chosen at the surgeon's discretion. The use of post-operative contrast swallows is optional.

### Previous interventions:

Both BYPASS and BAND surgical procedures will be carried out laparoscopically in a standard fashion under general anaesthesia with all patients receiving antibiotic and DVT prophylaxis. For the purposes of this pragmatic trial each intervention will be allowed to be implemented according to the standard local policy. Particular aspects of each intervention that are considered mandatory or prohibited are listed below. Fidelity to the surgical interventions will be monitored by completion of an operative manual.

# 1. Laparoscopic adjustable gastric banding (BAND surgery)

The procedure will involve placement of laparoscopic ports, creation of a pneumoperitoneum and placement of retractors as the surgeon chooses. The choice about the type and size of adjustable gastric band will be made by the surgeon. If a hiatal defect is present it may be repaired and closure of pre-existing umbilical hernia or undertaking concomitant cholecystectomy is at the discretion of the surgeon. It is considered mandatory to dissect the lesser curve using the 'Pars flaccida' technique, to use gastro-gastric tunnelling sutures and to fix the adjustable port to the anterior abdominal wall. An apronectomy is prohibited.

# 2. Laparoscopic gastric bypass surgery (BYPASS surgery)

The laparoscopic ports, creation of a pneumoperitoneum and placement of retractors may be performed as the surgeon chooses. Creation of the biliary and gastric limbs and formation of the gastric pouch is performed as the surgeon chooses. An upper limit of 100cm and 200cm is recommended for the biliary and gastric limbs. Testing integrity of the anastomoses, closure of pre-existing umbilical, internal hernia and hiatal hernia defects are optional, as is undertaking a cholecystectomy. Formation of a horizontal gastric pouch that includes fundus and undertaking an apronectomy is prohibited.

# Intervention Type

Procedure/Surgery

# Primary outcome measure

Current primary outcome measures as of 23/09/2015:

There are two primary endpoints:

1. The proportion achieving loss of greater than 50% of excess weight at three years (calculated as 100×[BMI at 3 years – BMI at randomisation\*] / [BMI at randomisation – 25])

2. HRQOL at three years (EQ-5D-5Lhealth state score)

\*Calculated using the participant weight recorded at baseline, after consent and before randomisation

Previous primary outcome measures:

The BY-BAND trial will compare weight loss and quality of life after the laparoscopic adjustable gastric banding operation (BAND) and the laparoscopic gastric bypass operation (BYPASS) to treat morbid obesity.

There are two co-primary outcomes:

 The proportion achieving loss of greater than 50% of excess weight at three years (calculated as 100×[BMI at 3 years - BMI at randomisation] / [BMI at randomisation - 25])
HRQOL at three years (EQ-5D health state score).

The hypothesis that the trial is testing, that the Bypass operation is superior to the Band operation, will be supported if:

1. Bypass is not inferior with respect to weight loss at 3 year AND

2. Bypass is superior to Band with respect to health-related quality of life

### Secondary outcome measures

Current primary outcome measures as of 23/09/2015:

To compare the cost effectiveness of the three operations, their acceptability and quality of life after surgery, nutritional outcomes, time to resolution of co-morbidities and complications. We will explore, in a sample, patients' experiences of surgery, outcomes and lifestyle change. We will also develop a core set of outcomes to use in future trials of morbid obesity surgery.

The secondary endpoints include:

- 1. Change in BMI over time adjusted for BMI at randomisation
- 2. % weight loss at 3 years
- 3. Waist circumference at 3 years

4. Time taken from randomisation to reach first loss of at least 50% of excess BMI

5. Time taken from first losing 50% excess BMI to first relapse (defined as weight re-gain such that the target of at least 50% of excess weight loss is no longer met)

6. Generic and symptom specific (i.e. obesity and GI specific) HRQOL: SF12, EQ5D, IWQOL-Lite, and GIQLI to three years

7. Resource use to three years

8. Standard NHS nutritional blood tests will be performed at each assessment including; full blood count, electrolytes, creatinine, glucose, HbA1c, liver function tests, iron, ferritin, vitamin B12, folate/red cell folate, lipid profile, 25-hydroxyvitamin D, calcium, parathyroid hormone
9. Measures of 24 hour recall eating using a standardised and validated interview process
10. Binge eating behaviour using a validated questionnaire

11. Adverse health events including the need for re-operation and cross over between interventions

12. Resolution of co-morbidities at 3 years, including sleep apnoea, non-alcoholic fatty liver disease, type 2 diabetes, hypertension and hyperlipidaemia

13. Time to resolution of sleep apnoea, type 2 diabetes, hypertension and hyperlipidaemia

Previous primary outcome measures:

To compare the cost-effectiveness of the two operations, their acceptability and quality of life after surgery, nutritional outcomes, time to resolution of co-morbidities and complications. We will explore, in a sample, patients' experiences of surgery, outcomes and lifestyle change. We will also develop a core set of outcomes to use in future trials of morbid obesity surgery.

- 1. Change in BMI over time adjusted for BMI at randomisation
- 2. Percentage weight loss at 3 years
- 3. Waist circumference at 3 years

4. Time taken from randomisation to reach first loss of at least 50% of excess BMI

5. Time taken from first losing 50% excess BMI to first relapse (defined as weight re-gain such that the target of at least 50% of excess weight loss is no longer met),

6. Generic and symptom specific (i.e. obesity and GI specific) HRQOL: SF12, EQ5D, IWQOL-Lite, and GIQLI to three years

7. Resource use to three years,

8. Standard NHS nutritional blood tests will be performed at each assessment including; full blood count, electrolytes, creatinine, fasting glucose, HbA1c, liver function tests, iron, ferritin, vitamin B12, folate/red cell folate, lipid profile, 25-hydroxyvitamin D, calcium, parathyroid hormone.

9. Measures of 24 hour recall eating using a standardised and validated interview process 10. Binge eating behaviour using a validated questionnaire

11. Adverse health events including the need for re-operation and cross over between interventions,

12. Resolution of co-morbidities at 3 years including sleep apnoea, non alcoholic fatty liver disease, type-2 diabetes, hypertension and hyperlipidaemia

13. Time to resolution of co-morbidities listed in 12 above

# Overall study start date

01/01/2012

# **Completion date**

30/09/2022

# Eligibility

# Key inclusion criteria

Current inclusion criteria as of 23/09/2015:

Participants may enter study if ALL of the following apply:

1. Male or female patients

2. Over 18 years of age

3. Referred for bariatric surgery according to NICE guidelines - BMI of 40kg/m2 or more, OR BMI of 35 kg/m2 to 40 kg/m2 and other significant disease (e.g. type 2 diabetes or high blood pressure) OR BMI of 30 kg/m2 or more and recent onset diabetes OR Asian family origin with lower BMI and recent onset diabetes, that could improve with weight loss

4. Has been or is willing to receive intensive management in a specialist tier 3 obesity service

- 5. Fit for anaesthesia and surgery
- 6. Committed to follow-up and able to complete quality of life questionnaires

7. Able to provide written informed consent

# Previous inclusion criteria:

The participant may enter study if ALL of the following apply:

1. Male or female patients

2. Over 18 years of age

3. Referred for bariatric surgery according to NICE guidelines - BMI of 40kg/m2 or more, OR BMI of 35 kg/m2 to 40 kg/m2 and other co-morbidities (e.g. type 2 diabetes), that could improve with weight loss

- 4. Willing to receive intensive management in a specialist obesity service
- 5. Fit for anaesthesia and surgery
- 6. Committed to follow-up and able to complete quality of life questionnaires
- 7. Able to provide written informed consent

# Participant type(s)

Patient

# Age group

Adult

# Lower age limit

18 Years

#### **Sex** Both

Target number of participants

1341

# Total final enrolment

1351

# Key exclusion criteria

Current exclusion criteria as of 23/09/2015:

Participants may not enter study if ANY of the following apply (assessed by patient history and clinical examination)

1. Previous gastric surgery or surgery for severe and complex obesity

2. Previous abdominal surgery or gastro-intestinal (GI) condition that precludes one or more of Band, Bypass or Sleeve

3. Large abdominal ventral hernia

4. Pregnancy (women who have given birth and women planning pregnancy will NOT be excluded)

5. Crohn's disease

- 6. Liver cirrhosis and portal hypertension
- 7. Systemic lupus erythematosis
- 8. Known silicone allergy

9. Hiatus hernia >5cm

10. Other clinical/psychological reason, to be specified

11. Active participation in another interventional research study which might interfere with By-Band-Sleeve

Previous exclusion criteria:

The participant may not enter study if ANY of the following apply:

- 1. A history of previous gastric surgery or surgery for morbid obesity
- 2. Large abdominal ventral hernia
- 3. Hiatus hernia more than 5 cm

4. Pregnancy (women who have given birth and women planning pregnancy will NOT be excluded)

- 5. Crohn's disease
- 6. Liver cirrhosis and portal hypertension
- 7. Systemic lupus erythematosis
- 8. Known silicone allergy
- 9. Surgeon unwilling to for patient to be randomised (reason to be specified)

**Date of first enrolment** 01/11/2012

**Date of final enrolment** 30/09/2019

# Locations

**Countries of recruitment** England

United Kingdom

**Study participating centre Musgrove Park Hospital** Taunton United Kingdom TA1 5DA

**Study participating centre University Hospital Southampton** Southampton United Kingdom SO16 6YD

**Study participating centre Royal Bournemouth and Christchurch Hospital** Bournemouth United Kingdom BH7 7DW

**Study participating centre St James University Hospital** Leeds United Kingdom LS9 7TF

Study participating centre

**Sunderland Royal Hospital** Sunderland United Kingdom SR4 7TP

**Study participating centre Royal Cornwall Hospital** Truro United Kingdom TR1 3LQ

**Study participating centre Homerton University Hospital** London United Kingdom E9 6SR

**Study participating centre Royal Derby Hospital** Derby United Kingdom DE22 3NE

**Study participating centre Queen Alexandra Hospital** Portsmouth United Kingdom PO6 3LY

**Study participating centre Heart of England NHS Foundation Trust** Birmingham United Kingdom B9 5SS

**Study participating centre Imperial College Healthcare NHS Trust** Salton House St Mary's Hospital London United Kingdom W2 1NY

Study participating centre North Bristol Trust Brunel building Southmead Hospital Westbury-on-Trym Bristol United Kingdom BS10 5NB

# Sponsor information

**Organisation** University of Bristol (UK)

**Sponsor details** c/o Dr Birgit Whitman Senate House Tyndall Avenue Clifton Bristol England United Kingdom BS8 1TH

**Sponsor type** University/education

Website http://www.bristol.ac.uk/

ROR https://ror.org/0524sp257

# Funder(s)

**Funder type** Government **Funder Name** Health Technology Assessment Programme

Alternative Name(s) NIHR Health Technology Assessment Programme, HTA

**Funding Body Type** Government organisation

Funding Body Subtype National government

**Location** United Kingdom

# **Results and Publications**

# Publication and dissemination plan

The trial results will be published and disseminated at a later date.

### Intention to publish date

30/09/2023

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from the study mailbox address (by-band-sleeve@bristol.ac.uk).

#### IPD sharing plan summary

Available on request

#### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient- facing?
<u>Other</u> publications	recruiter perspectives	06/01 /2014		Yes	No
<u>Protocol</u> <u>article</u>	protocol	11/02 /2014		Yes	No
<u>Other</u> publications	most effective operation for adults with severe and complex obesity:	14/03 /2014		Yes	No
<u>Other</u> publications	development of a core outcome set	01/01 /2015		Yes	No
<u>Results article</u>	qualitative case study results from By-Band	28/12 /2015		Yes	No
<u>Other</u> publications	NHS access to bariatric surgery	11/05 /2016		Yes	No
<u>Other</u> publications	use of adjustable gastric bands for management of severe and complex obesity:	01/06 /2016		Yes	No
<u>Other</u> publications	conveying equipoise during recruitment:	18/10 /2016		Yes	No

<u>Other</u> publications	health professionals' and patients' views of the importance of outcomes	01/11 /2016		Yes	No
<u>Other</u> publications	core outcome set	29/11 /2016		Yes	No
<u>Other</u> publications	adaptation of By-Band randomized clinical trial	01/08 /2017		Yes	No
<u>Other</u> publications	costs of bariatric surgery	01/08 /2017		Yes	No
<u>Results article</u>	By-Band pilot study results and explanation of By-Band- Sleeve design	01/08 /2017		Yes	No
<u>Other</u> publications	enabling recruitment success	01/11 /2017		Yes	No
<u>Other</u> publications	EQ-5D-5L to measure health-related quality of life:	18/12 /2017		Yes	No
<u>Other</u> publications	QuinteT Recruitment Intervention	01/02 /2019		Yes	No
<u>Other</u> publications	micro-costing of procedures	01/02 /2019		Yes	No
<u>Other</u> publications	prevalence, and severity, of anxiety and depressive symptoms	03/01 /2024	08/01 /2024	Yes	No
<u>Results article</u>		31/03 /2025	04/04 /2025	Yes	No