Evaluation of lifespan in AN69ST with two different heparinization strategies

| Submission date | Recruitment status | Prospectively registered |
|-------------------|---------------------------------|-----------------------------|
| 13/02/2008 | No longer recruiting | ☐ Protocol |
| Registration date | Overall study status | Statistical analysis plan |
| 13/03/2008 | Completed | Results |
| Last Edited | Condition category | Individual participant data |
| 14/03/2008 | Urological and Genital Diseases | Record updated in last year |

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Beng-Åke Henriksson

Contact details

Sahlgrenska University Hospital Department of Intensive Care (CIVA) Gothenburg Sweden SE- 413 45

Additional identifiers

Protocol serial number 1446

Study information

Scientific Title

Study objectives

Reduction of heparin dose in acute renal failure treated with hemodiafiltration.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Local research ethics committee, Göteborg, Sweden. Date of approval: 17/12/2007 (Dnr. 576-07)

Study design

Monocentric, open, randomised and controlled study with two parallel groups.

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Acute renal failure/fluid overload

Interventions

The same dialysis membrane, AN69ST, will be used for both the intervention and control treatment. Two different heparinization strategies will be compared:

Control group (usual heparin doses): Bolus at start (between 1,000 IU - 2,500 IU) + usual infusion dose (10 IU/kg/h)

Intervention group (Reduction of heparin dose): No bolus at start + infusion decreased by 50% (5 IU/kg/h)

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Filter lifespan, defined as the time period between patient's connection and filter disconnection due to filter clotting.

Key secondary outcome(s))

Adverse events, monitored for up to 24 days of treatment

Completion date

01/01/2010

Eligibility

Key inclusion criteria

- 1. Patients requiring continuous renal replacement therapies (CRRT)
- 2. Aged 18 and over
- 3. Patients weighing 30-120 kg
- 4. Patients having signed a written consent to participate in the study

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Αll

Key exclusion criteria

- 1. History of heparin antibodies or heparin-induces thrombocytopenia
- 2. Known hypertensive to any dialysis membrane
- 3. Current enrolment in another trial which could impact the successful completion of this study
- 4. Patients under guardianship
- 5. Patient anticipated to require transportation outside the unit for diagnostic or therapeutic procedures in the coming first week .These patients can eventually be included in a later more stable phase
- 6. Unconscious patients for whom no relative or person of trust can give consent to treatment. In the absence of any relative or person of trust, the patient in question cannot be included in the study
- 7. Patient requiring anticoagulation for other indications e.g., valvular surgery or extracorporeal ventricular assist devices
- 8. Patients with active bleeding who does not require heparinization

Date of first enrolment

12/02/2008

Date of final enrolment

01/01/2010

Locations

Countries of recruitment

Sweden

Study participating centre
Sahlgrenska University Hospital
Gothenburg
Sweden

SE-41345

Sponsor information

Organisation

Gambro Lundia AB (Sweden)

ROR

https://ror.org/05mw5ed57

Funder(s)

Funder type

Industry

Funder Name

Gambro Lundia AB (Sweden)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet Participant information sheet 11/11/2025 11/11/2025 No Yes