

Prevention of Atrial Arrhythmias by Infusion of Magnesium Sulphate after Lung Resection for Cancer

| | | |
|--|---|--|
| Submission date 28/09/2007 | Recruitment status No longer recruiting | <input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol |
| Registration date 28/09/2007 | Overall study status Completed | <input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results |
| Last Edited 02/08/2012 | Condition category Cancer | <input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year |

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Mr N Chaudhuri

Contact details
Department of Thoracic Surgery
The Cardiothoracic Centre
Thomas Drive
Liverpool
United Kingdom
L14 3PE

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
N0054186615

Study information

Scientific Title

Study objectives

We hypothesize that the prophylactic infusion of magnesium sulphate significantly reduces incidences of postoperative atrial fibrillation in patients undergoing thoracotomy, pneumonectomy and lobectomy.

This is a single-centre, controlled and randomised clinical trial. Study patients will be recruited into the trial from participating surgeons waiting lists and those consenting for the trial will be included in the trial register. Atrial arrhythmias are common after pulmonary procedures with reported incidences ranging from 10% to 20% after lobectomy and up to 40% after pneumonectomy. To date only one randomised study published nearly a decade ago showed that postoperative atrial tachyarrhythmias, mainly atrial fibrillation could be significantly reduced when magnesium sulphate was administered. A general consensus is still lacking on the efficacy of magnesium sulphate as anti-arrhythmic drug for patients undergoing lung cancer operation. We plan to investigate in this by carrying out a randomised, parallel, blinded, controlled trial involving a total of 240 patients (n=120 per group).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Single-centre controlled and randomised clinical trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Prevention

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Cancer: Lung

Interventions

Patients will be randomly allocated in equal numbers to treatment with or without magnesium sulphate and followed-up during the in-hospital perioperative period.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Magnesium Sulphate

Primary outcome measure

Atrial fibrillation will be assessed using a standard 12-lead ECG measurement from the onset of the Q wave to the end of the S wave.

Secondary outcome measures

1. Myocardial Infarction will be defined by two of the following three criteria if present: Unequivocal ECG changes. Elevation of cardiac enzyme(s): 3 times upper limit of creatinine kinase CK/CKMB) and above twice the upper limit of normal or elevated troponin (T/I). Chest pain typical of ischaemia lasting for more than 20 minutes.
2. Respiratory complications as determined by the requirement for postoperative mechanical ventilation greater than 24 hours or reintubation for ventilatory support after the day of surgery; pneumonia - defined as fever leukocytosis, pulmonary infiltrate requiring antibiotic therapy; air leak from thoracostomy tubes for more than six days postoperatively; lobar collapse on postoperative chest radiograph; empyema and bronchopleural fistula. Sputum retention defined as failure to clear bronchial secretions that can result in: bronchial obstruction, atelectasis, lobar collapse, secondary pulmonary infection.
3. Renal Complications - postoperative rise in serum creatinine above 200µmol/litre or requirement of postoperative dialysis support in a patient with normal preoperative renal function was considered to be a renal complication.
4. Neurological complications - a postoperative new cerebrovascular accident or transient ischaemic attack was considered to be a neurological complication.
5. Pulmonary embolus defined as a blockage of an artery in the lungs by fat, air, tumor tissue, or blood clot.
6. Death is defined as all-cause mortality and the cause of death will have to be specified. In the analysis, all deaths during the study period will be compared between the two groups.

Overall study start date

01/09/2006

Completion date

01/09/2008

Eligibility

Key inclusion criteria

240 subjects will be recruited. The investigator will be responsible for screening all patients listed for lung surgery at the CTC. Evidence from Terzi et al [32], had demonstrated that incidences of AF occurred in 10.7% of patients on magnesium sulphate as compared with 26.7%

of control patients. We predict that at 90% power (2-sided $\alpha=0.05$), 240 patients allowing for drop outs, deaths etc) are required to detect at least a 16% reduction in incidences of AF.

Inclusion Criteria:

1. Elective for non-cardiac thoracic operations for lung cancer (lobectomy and pneumonectomy)
2. Ability to provide informed consent

Participant type(s)

Patient

Age group

Not Specified

Sex

Not Specified

Target number of participants

240

Key exclusion criteria

1. Patients not undergoing major lung resection (see above)
2. Impaired renal function, preoperative creatinine >200 mol/L
3. Myocardial infarction within the last 6 months
4. Urgent or emergency operations
5. Video assisted thoracic surgery
6. Currently on drugs with antiarrhythmic properties
7. Patients with a history of preoperative cardiac arrhythmias

Date of first enrolment

01/09/2006

Date of final enrolment

01/09/2008

Locations

Countries of recruitment

England

United Kingdom

Study participating centre

Department of Thoracic Surgery

Liverpool

United Kingdom

L14 3PE

Sponsor information

Organisation

Record Provided by the NHSTCT Register - 2007 Update - Department of Health

Sponsor details

The Department of Health, Richmond House, 79 Whitehall
London
United Kingdom
SW1A 2NL
+44 (0)20 7307 2622
dhmail@doh.gsi.org.uk

Sponsor type

Government

Website

<http://www.dh.gov.uk/Home/fs/en>

Funder(s)**Funder type**

Government

Funder Name

The Cardiothoracic Centre Liverpool NHS Trust (UK), NHS R&D Support Funding

Results and Publications**Publication and dissemination plan**

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration