A randomised trial comparing hormonal treatment versus combination of hormonal treatment and radiotherapy in locally advanced prostate cancer

Submission date	Recruitment status No longer recruiting	Prospectively registered		
29/05/2008		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
13/06/2008	Completed	[X] Results		
Last Edited	Condition category	[] Individual participant data		
08/04/2009	Cancer			

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Anders Widmark

Contact details

Department of Radiation Sciences, Oncology Umeå University, Hospital Umeå Sweden 90185 Anders.Widmark@onkologi.umu.se

Additional identifiers

Protocol serial number

Version 1996 01 09

Study information

Scientific Title

Randomised trial of locally advanced/aggressive prostatic cancer: T3 diff grad 1-3, T1b - T2 diff grad 2-3 (optional), NO, MO antiandrogen treatment with or without radiotherapy

Acronym

SPCG-7/SFUO-3

Study objectives

To evaluate if the addition of radiotherapy improves the outcome in hormonally-treated, locally advanced/aggressive, node-negative and non-metastasised prostatic cancer.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received from Umeå University, Medical Faculty Ethical Committee in 1995 (ref: paragraph 247/95; diary no. 95-179).

Study design

A randomised open, comparative, parallel design trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Local or locally advanced prostate cancer, pN0, M0

Interventions

After randomisation all patients were treated with total androgen blockade (TAB) with an LHRH-agonist (Procren Depot; Abbott), for three months in combination with an oral anti-androgen (AA) Eulexin (Schering-Plough) 250 mg x 3. Thereafter all patients continued on the anti-androgen alone, T Eulexin 250 mg x 3 daily, continuously until progression. After three months patients in the Radiotherapy+Hormone arm started radiotherapy whereas patients in the Hormone-only arm had no local treatment. Minimal radiation dose to the prostate will be 70 Gy and the seminal vesicles will be included up to a minimum dose of 50 Gy.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Leuprolide (Procren Depot), flutamide (Eulexin)

Primary outcome(s)

To evaluate if cause-specific seven-year survival can be improved in patients treated with a combination of radiotherapy and anti-androgen as compared to anti-androgen therapy only.

Primary and secondary endpoints were planned to be analysed after seven years.

Key secondary outcome(s))

- 1. To evaluate:
- 1.1. Time to biochemical progression (PSA)
- 1.2. Time to symptoms related to local progression
- 1.3. Time to symptoms related to distant progression
- 2. To evaluate quality of life (QOL) with special focus on sexual function, urinary and gastrointestinal morbidity

Primary and secondary endpoints were planned to be analysed after seven years.

Completion date

30/12/2002

Eligibility

Key inclusion criteria

- 1. Men less than 76 years of age and, as judged by the doctor, a life expectancy of less than 10 years (except for cancer) at time of randomisation with performance status World Health Organization (WHO) 0 2
- 2. Patients with histologically/cytologically verified prostatic cancer
- 3. Patients with prostatic cancer of clinical category T1b-T2; G2-G3 and T3; G1-G3 according to the TNM classification system of 1992. Inclusion of patients with T1b-T2; G3 and T2; G2 is optional.
- 4. The patients should have no evidence of metastases by clinical investigation, bone scan or pulmonary x-ray
- 5. Patients should be lymph node negative
- 6. Patients should be suitable for radiotherapy and anti-androgen treatment

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Male

Key exclusion criteria

- 1. Patients who earlier have undergone any other treatment against prostatic cancer except transurethral resection of the prostate (TUR-P)
- 2. Patients with a prostate specific antigen (PSA) greater than 70 ng/ml
- 3. Patients unable to cooperate or suffering from any other form of disease that would interfere with the planned treatment (e.g. colitis)
- 4. Liver function that would interfere with the anti-androgen treatment (a bilirubin and/or alanine aminotransferase [ALAT] value above the upper normal limit)

5. Patients with previous diagnosis of other malignant disease. Exceptions could be made for basal cell carcinoma of the skin or progression free survival at least 10 years after any previous tumour (this requires discussion with the study co-ordinator).

Date of first enrolment

27/02/1996

Date of final enrolment

30/12/2002

Locations

Countries of recruitment

Denmark

Norway

Sweden

Study participating centre

Department of Radiation Sciences, Oncology

Umeå

Sweden

90185

Sponsor information

Organisation

Scandinavian Prostate Cancer Group (SPCG) (Sweden)

Funder(s)

Funder type

Research organisation

Funder Name

Scandinavian Prostate Cancer Group (SPCG) (Sweden) - after receiving an unrestricted grant from Schering-Plough Inc. and Abbott Scandinavia Inc.

Funder Name

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created Date added	Peer reviewed?	Patient-facing?
Results article	results	24/01/2009	Yes	No
Results article	four-year follow-up results	01/04/2009	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	No	Yes