The effect of participatory womens groups on birth outcomes in Bangladesh: does coverage matter?

Submission date Recruitment status Prospectively registered 28/04/2011 No longer recruiting [X] Protocol [] Statistical analysis plan Registration date Overall study status 15/06/2011 Completed [X] Results [] Individual participant data **Condition category** Last Edited Pregnancy and Childbirth 22/05/2013

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Anthony Costello

Contact details

UCL Institute of Child Health Centre for International Health and Development London United Kingdom WC1N 1EH +44 (0)20 7905 2883 a.costello@ich.ucl.ac.uk

Additional identifiers

Protocol serial number 1VTP/07CH05

Study information

Scientific Title

The effect of participatory womens groups on birth outcomes in Bangladesh: does coverage matter? A cluster randomised controlled trial

Acronym

Perinatal Care Project (PCP) Diabetic Association of Bangladesh (BADAS)

Study objectives

We hypothesise that the intervention will lead to a 30% lower neonatal mortality in intervention clusters compared with control clusters, during the last 24 months of the study.

Ethics approval required

Old ethics approval format

Ethics approval(s)

- 1. UCL Research Ethics Committee approved in March 2008; annually reviewed (ref:ID number: 1488/001)
- 2. Diabetic Association of Bangladesh Ethical Review Committee approved on 7th July 2010 (but approval was granted before the start of the trial, as part of our ongoing women's group work that had started in 2005)

Study design

Cluster randomised controlled trial

Primary study design

Interventional

Study type(s)

Screening

Health condition(s) or problem(s) studied

Maternal and Child Health

Interventions

The intervention is a participatory learning action cycle with womens groups. In each intervention cluster, facilitators convene womens groups that meet on a monthly basis. The participatory learning and action cycle has 4 phases: First, the groups identify and prioritise health problems, then plan strategies to address these problems, subsequently they put these strategies into practice, and finally, they evaluate their strategies.

As we aim to study the effect of an intervention with a high coverage of womens groups in the population, 648 new groups were formed by newly recruited facilitators and started to meet from January 2009 onwards, in addition to the 162 womens groups that were already set up in the intervention areas as part of an earlier trial in the same study area. These old groups have continued to meet on a monthly basis from late 2004 onwards. The 648 new groups will go through a cycle of monthly meetings on maternal and newborn health (Cycle 1), while from April 2010 the 162 old groups will proceed to a cycle of meetings on under-5 and womens health (Cycle 2), while periodically continuing to review maternal and newborn health issues. The combined 810 womens groups constitute a coverage of 1 group per 300 population, in comparison with 1 group per 1414 population in the previous trial.

The role of the facilitator is to activate and strengthen groups, support them in identifying problems, help to plan possible solutions and support the implementation and monitoring of solution strategies in the community. Although this role requires a grasp of health issues and some knowledge of potential interventions, she needs to be a facilitator rather than a teacher. As such, the facilitator may act as a broker of information and communication but her prime importance is as a catalyst for community mobilisation.

All clusters, control and intervention, receive health system strengthening activities.

Patient information sheets are not used because this was a community trial of a social intervention (i.e. not a clinical trial). The intervention consists of women's groups that discuss and design their own strategies to improve newborn and maternal health. All the women in these women's groups participate voluntarily. At the start of the women's groups, there was extensive discussion of what the aims and structure of the women's groups are. By voluntarily joining a women's group, the participants consent to the intervention (i.e. women's groups). Oral consent was obtained from the respondents in the monitoring and surveillance interviews.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Neonatal mortality (deaths in the first 28 complete days after birth per 1,000 live births), during the last 24 months of the study.

Key secondary outcome(s))

- 1. Early and late neonatal mortality rate
- 2. Stillbirth rate
- 3. Pregnancy-related mortality ratio and maternal mortality ratio
- 4. Health care use
- 5. Home care practices

Completion date

30/06/2011

Eligibility

Key inclusion criteria

- 1. Women, living in the study area, who are permanent residents in the union in which their delivery or their death was identified.
- 2. Women and their newborn infants are included after birth, or, if a woman dies during pregnancy, after her death.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Key exclusion criteria

- 1. Women who are temporary residents in the union in which their delivery or death was identified
- 2. Women who decline to be interviewed
- 3. Women who reside outside the study area

Date of first enrolment

01/01/2009

Date of final enrolment

30/06/2011

Locations

Countries of recruitment

United Kingdom

England

Bangladesh

Study participating centre UCL Institute of Child Health

London United Kingdom WC1N 1EH

Sponsor information

Organisation

University College London (UK)

ROR

https://ror.org/02jx3x895

Funder(s)

Funder type

Charity

Funder Name

Big Lottery Fund Strategic Grant (UK) IS/2/010281409

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/09/2013		Yes	No
Protocol article	protocol	26/09/2011		Yes	No
Other publications	coverage estimates	29/06/2012		Yes	No