

The effect of the laparoscopic reconstruction of a caesarean scar (Niche)

Submission date 23/04/2013	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 27/02/2014	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 14/09/2023	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

In Western countries, caesarean rates are rising. A niche is a cavity that can develop at the site of an old caesarean section scar which occurs in approximately 60% of the women who had a caesarean section. A niche can cause abnormal uterine blood loss, intracavitary accumulation of fluid and inaccessibility of the uterine cavity (i.e. for intra-uterine insemination or embryo transfer). Surgical therapy can possibly reduce these complaints. In general, a hysteroscopic resection (surgery using a thin telescope called a hysteroscope and no incision is made) and/or coagulation of the vessels in the niche will be sufficient. However, in the case of a large niche, a laparoscopic resection (surgery in which only small incisions are made) and closure of the niche are preferred. It is assumed that in these cases the risk of bladder injury during hysteroscopic niche resection is too high. The main aim is to study the effectiveness of laparoscopic niche resection in the treatment of niche-related bleeding disorders. Other aims are to study the effect of laparoscopic niche resection on menstrual pain, quality of life and fertility, intracavitary accumulation of fluid and niche characteristics.

Who can participate?

Women with niche-related abnormal uterine blood loss, dysmenorrhoea (painful periods) or accumulation of intracavitary fluid with residual myometrium (middle layer of the uterine wall) < 3mm

What does the study involve?

All patients will undergo surgery (laparoscopic niche resection) and will be followed up for 6 months after the surgery.

What are the possible benefits and risks of participating?

We expect that laparoscopic reconstruction of a niche will reduce prolonged menstrual bleeding, and associated menstrual pain and as a consequence will improve related quality of life. In addition, we expect the laparoscopic niche reconstruction to reduce the accumulation of intra-uterine fluid. We additionally expect the reconstruction to increase the thickness of the remaining myometrium between the bladder and the niche. Risks of the intervention include bleeding and bladder injury during the surgery and incomplete niche reconstruction.

Where is the study run from?
VU University Medical Center (Netherlands).

When is the study starting and how long is it expected to run for?
The study started in September 2009 and will continue until December 2015.

Who is funding the study?
VU University Medical Center (Netherlands).

Who is the main contact?
Prof. Dr. H.A.M. Brolmann, h.brolmann@vumc.nl

Contact information

Type(s)
Scientific

Contact name
Prof H.A.M. Brölmann

Contact details
Department of Obstetrics and Gynaecology
VU University Medical Center
Postbus 7057
Amsterdam
Netherlands
1007 MB

Additional identifiers

Protocol serial number
ABR: NL37922.029.11

Study information

Scientific Title
The effect of the laparoscopic reconstruction of a caesarean scar (Niche): an observational pilot study

Acronym
LapNiche

Study objectives
Laparoscopic niche resection will reduce the complaints and possibly increase fertility chances in women with a niche with residual myometrium < 3 mm and complaints of abnormal uterine blood loss / dysmenorrhoea or intracavitary accumulation of fluid.

Ethics approval required
Old ethics approval format

Ethics approval(s)

Ethical Board VUmc, 02/10/2011, ref.:2011/297

Study design

Observational study of an intervention (prospective cohort study)

Primary study design

Observational

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Abnormal Uterine Bleeding / Niche resection

Interventions

This is an observational study of an intervention, laparoscopic niche resection.

There is only one study arm. The laparoscopic niche resection/reconstruction is an operative laparoscopy where the roof of the niche is resected and the myometrial wound is closed in two layers with Vicryl. The operation is performed under hysteroscopic guidance and lasts about 2-3 hours. The patient is followed up for 6 months after the intervention. Symptoms of postmenstrual spotting are assessed and a transvaginal ultrasound is made and compared to baseline values.

Intervention Type

Procedure/Surgery

Phase

Not Applicable

Primary outcome(s)

Reduction of abnormal uterine blood loss in days, measured at baseline and after 3, 6, 9 and 12 months

Key secondary outcome(s)

1. Quality of life (SF36)
2. Dysmenorrhoe
3. Fertility/pregnancy
4. Intracavitary accumulation of fluid

All above outcomes measured at baseline and after 3, 6, 9 and 12 months

Completion date

31/01/2018

Eligibility**Key inclusion criteria**

1. A niche with residual myometrium < 3 mm (as measured with SIS / GIS)
2. Complaints of abnormal uterine blood loss, dysmenorrhoea or accumulation of intracavitary fluid
3. Premenopausal women aged above 18

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Female

Total final enrolment

133

Key exclusion criteria

1. Pregnancy
2. Age below 18
3. Other disorders that may induce spotting [polyps (everywhere in the uterus/cervix are exclusion criteria except those in the niche)], fibroids, known atypical endometrial cells, cervical dysplasia, cervical or pelvic infection, assumed malignancy, irregular cycle (>35 days or intercycle variation of 2 weeks or more)
4. Contra-indications for spinal or general anaesthesia

Date of first enrolment

01/01/2009

Date of final enrolment

12/12/2015

Locations**Countries of recruitment**

Netherlands

Study participating centre

Department of Obstetrics and Gynaecology

Amsterdam

Netherlands

1007 MB

Sponsor information

Organisation

VU University Medical Center (Netherlands)

ROR

<https://ror.org/00q6h8f30>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Time of the researchers is paid by the institution VU University Medical Center (VUmc) (Netherlands).

Funder Name

As it is an observational study of usual care the direct costs are reimbursed by the insurance company.

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	Long-term follow-up	12/09/2023	14/09/2023	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes