Imaging-based Thrombolysis trial in Acute Ischemic Stroke -III

Submission date	Recruitment status	Prospectively registered
14/10/2007	No longer recruiting	Protocol
Registration date	Overall study status	Statistical analysis plan
30/10/2007	Completed	Results
Last Edited	Condition category	[] Individual participant data
30/10/2007	Circulatory System	Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Yongjun Wang

Contact details

Neurology Department
Beijing Tiantan Hospital
Tiantan Xili 6
Chongwen District
Beijing
China
100050
yongjunwang111@yahoo.com.cn

Additional identifiers

Protocol serial number

5

Study information

Scientific Title

Acronym

ITAIS- III

Study objectives

Whether the patients without occlusion or significant stenosis of the large vessels should receive thrombolysis had been focused on and questioned for a long time. However, there is still no large Randomised Controlled Trial (RCT) to prove if these patients can benefit from thrombolysis. The objective of this trail is to investigate:

- 1. For the selected acute ischemic stroke patients without large vessels occlusion or significant stenosis, whether the IntraVenous (IV) thrombolysis can improve the clinical outcome if the mismatch exists on CTP/Computed Tomography Angiography-Source Images (CTA-SI) scan in 3-9 hr time-window
- 2. For the selected acute ischemic stroke patients without large vessels occlusion or significant stenosis, whether the IV thrombolysis can improve the blood perfusion of brain if the mismatch exists on CT perfusion (CTP)/CTA-SI scan in 3-9 hr time-window
- 3. For the selected acute ischemic stroke patients without large vessels occlusion or significant stenosis, but the mismatch exists on CTP/CTA-SI scan in 3-9 hr time-window, whether the safety of IV thrombolysis is better than the NINDS trial which was in 3 hr window or implemented on the patients with large vessels occlusion or significant stenosis in 3-9 hr time-window.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approval gained from the Beijing Tiantan Hospital Capital Medical University Ethics Board on October 10, 2006.

Study design

A prospective, multi-center, assessor-blind controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Acute ischemic stroke

Interventions

Patients included were divided into recombinant tissue Plasminogen Activator (rt-PA) (0.9 mg /kg) intravenous thrombolysis group and regular treatment group according to the intention of doctors and patients.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Recombinant tissue Plasminogen Activator (rt-PA)

Primary outcome(s)

Proportion of participants who achieve the following:

- 1. Reperfusion improvement, assessed 24 to 48 hours post-treatment and defined as either greater than or equal to 30% reduction of Mean Transit Time (MTT) volume of abnormality
- 2. Good clinical outcome at 90 days defined as a modified Rankin score (mRS) of 0-1
- 3. Intracerebral haemorrhage within 24-36 h after thrombolysis

Key secondary outcome(s))

Proportion of participants who achieve the following:

- 1. mRS 0 to 2 at 90 days
- 2. Barthel Index (BI) score 75 to 100 at 90 days
- 3. NIHSS 4 points improvement or 0-1 at 2 hours after treatment
- 4. NIHSS 4 points improvement or 0-1 at 24 to 48 hours
- 5. NIHSS 4 points improvement or 0-1 at day 7

Completion date

01/06/2010

Eligibility

Key inclusion criteria

- 1. Female or male inpatients
- 2. Age 18-80 years
- 3. Clinical diagnosis of ischaemic stroke
- 4. Onset of symptoms within 3-9 hours prior to initiation of thrombolysis treatment
- 5. Stroke symptoms present for at least 30 minutes and has not significantly improved before treatment
- 6. The National Institutes of Health Stroke Scale (NIHSS) score greater than or equal to 4
- 7. m-CT screening to be started within 8.5 hrs after stroke onset
- 8. Perfusion abnormality of CT scan >2cm in diameter involving hemisphere
- 9. CT Perfusion/CTA source image mismatch greater than or equal to 20%
- 10. CTA shows occlusion or significant stenosis of large vessels (Thrombolysis In Cerebral Ischemia [TICI] grade is 0 or 1)
- 11. Patients are willing to receive thrombolysis treatment and to give informed consent
- 12. Patients are willing and able to comply with the study protocol

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

80 years

Sex

All

Key exclusion criteria

- 1. Evidence of IntraCranial Haemorrhage (ICH), brain tumors, vascular malformation, aneurysm, SubArachnoid Hemorrhage (SAH)
- 2. Major infarct involving greater than 1/3 of Middle Cerebral Artery (MCA) territory on the CTA-SI
- 3. Presenting obvious neurologic deficits because of past stroke (mRS >2)
- 4. Severe stroke as assessed clinically (e.g. The National Institute of Health Stroke Scale [NIHSS] >25) and/or by appropriate magnetic imaging techniques.
- 5. Seizure at onset of stroke
- 6. Prior stroke within the last 3 months
- 7. Patients with any history of prior stroke and concomitant diabetes
- 8. Administration of heparin within the previous 48 hours and a thromboplastin time exceeding the upper limit of normal for laboratory
- 9. Platelet count of below 100,000/mm3
- 10. Uncompensated hypertension at study entry or hypertension requiring aggressive treatment to reduce blood pressure to nonhypertensive limits. Uncompensated hypertension is defined as systolic blood pressure >185 mm Hg or diastolic blood pressure >=110 mm Hg on 3 repeated measures at least 10 minutes apart
- 11. Blood glucose <50 or >400 mg/dl
- 12. Known haemorrhagic diathesis within the last 6 months
- 13. Patients receiving oral anticoagulants e.g. warfarin sodium, and coagulant response time (INR) >1.5
- 14. Known history of or suspected intracranial haemorrhage including subarachnoid haemorrhage
- 15. Pregnancy or lactation
- 16. Any history of severe central nervous system damage (i.e. neoplasm, aneurysm, intracranial or spinal surgery)
- 17. Haemorrhagic retinopathy,e.g. in diabetes (vision disturbances may indicate haemorrhagic retinopathy)
- 18. Bacterial endocarditis, pericarditis
- 19. Prolonged traumatic external heart massage, or recent (less than 10 days) obstetrical delivery or recent puncture of a non-compressible blood-vessel (e.g. subclavian or jugular vein puncture)
- 20. Acute pancreatitis
- 21. Documented ulcerative gastrointestinal disease during the last 3 months,
- 22. Oesophageal varices, arterial aneurysm, arterial/venous malformation
- 23. Neoplasm with increased bleeding risk
- 24. Severe liver disease including hepatic failure, cirrhosis, portal hypertension, oesophageal varices and active hepatitis
- 25. Major surgery or significant trauma in past 10 days
- 26. Known serious sensitivity to alteplase

Date of first enrolment

01/10/2007

Date of final enrolment 01/06/2010

Locations

Countries of recruitment

China

Study participating centre Neurology Department Beijing China 100050

Sponsor information

Organisation

Beijing Tiantan Hospital (China)

ROR

https://ror.org/003regz62

Funder(s)

Funder type

Government

Funder Name

The Ministry of Science and Technology and the Ministry of Health of the People's Republic of China - the key scientific research program of the 11th National Five-Year Planning of China

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration