

Efficacy of EPs 7630 in acute non-streptococcal tonsillopharyngitis

Submission date 14/06/2002	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 14/06/2002	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 02/10/2007	Condition category Ear, Nose and Throat	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Marianne Heger

Contact details
Director Research Center HomInt
PO Box 41 02 40
Karlsruhe
Germany
76202

Additional identifiers

Protocol serial number
UM002

Study information

Scientific Title

Study objectives

This study was a prospective, monocentre, randomised, open clinical pilot trial, comparing the efficacy of the herbal medicine EPs 7630 versus symptomatic therapy in patients with acute, non-streptococcal tonsillopharyngitis.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Acute, non-streptococcal tonsillopharyngitis

Interventions

60 patients were randomised to receive either:

1. Herbal remedy EPs 7630, 20 drops thrice daily, or
2. Symptomatic therapy (gargling with fruit vinegar and lukewarm water, Priessnitz compresses).

The duration of individual treatment lasted over a maximum of 10 days.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

EPs 7630

Primary outcome(s)

Not provided at time of registration

Key secondary outcome(s))

Not provided at time of registration

Completion date

30/04/1997

Eligibility

Key inclusion criteria

The study took place between March and April 1997. It is terminated. Patients, who met the following inclusion criteria, were suitable for the trial:

1. Age 6-10 years
2. Acute exsudative tonsillopharyngitis
3. Duration of symptoms less than 48 h
4. No Group A Beta Hemolytic Streptococcus (GABHS)-infection
5. Tonsillopharyngitis Severity Score (TSS) 6 or more points, and
6. Informed consent in writing by legal guardians who were able to understand the nature, meaning and consequences of the trial

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Not Specified

Sex

Not Specified

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/03/1997

Date of final enrolment

30/04/1997

Locations**Countries of recruitment**

Germany

Study participating centre

Director Research Center HomInt

Karlsruhe

Germany

76202

Sponsor information

Organisation

ISO Arzneimittel GmbH & Co KG (Germany)

ROR

<https://ror.org/045xrc244>

Funder(s)**Funder type**

Not defined

Funder Name

ISO Arzneimittel GmbH & Co KG

Results and Publications**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration