Dose escalation to intraprostatic tumour nodules in localised prostate cancer

Submission date	Recruitment status No longer recruiting	Prospectively registered	
17/08/2011		Protocol	
Registration date	Overall study status	Statistical analysis plan [Y] Posults [Y]	
	Condition entropy	[] Individual participant data	
29/09/2022	Cancer		

Plain English summary of protocol

http://cancerhelp.cancerresearchuk.org/trials/a-study-looking-at-increasing-dose-of-radiotherapy-to-areas-of-cancer-inside-prostate-gland-delineate

Contact information

Type(s) Scientific

Contact name

Ms Annie Gao

Contact details

Downs Road Sutton United Kingdom SM2 5PT

annie.gao@rmh.nhs.uk

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers 10309

Study information

Scientific Title

Dose EscaLation to Intraprostatic tumour Nodules in localisEd prostATE cancer: A phase II study examining the toxicity and feasibility of a dose escalated boost to a magnetic resonance imaging identified tumour nodule or nodules in localised prostate cancer

Acronym

DELINEATE

Study objectives

Dose Escalation to Intra-prostatic Tumour Nodules in Localised Prostate Cancer To assess the toxicity and feasibility of a dose escalated intensity-modulated radiotherapy boost to tumour nodules within the prostate using anatomical and functional magnetic resonance (MR) imaging to identify tumour. The aim is to maintain current levels of late toxicity.

Ethics approval required

Old ethics approval format

Ethics approval(s) 11/LO/0510

Study design Non-randomised; Interventional; Design type: Process of Care, Treatment

Primary study design Interventional

Secondary study design

Non randomised study

Study setting(s) Hospital

Study type(s) Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Topic: National Cancer Research Network; Subtopic: Prostate Cancer; Disease: Prostate

Interventions

Interventions as of 03/05/2017:

350 patients to be recruited and have MRI scans to diagnose intra-prostatic tumour nodules. Intra-prostatic tumour lesions will be treated with a radiotherapy boost. Radiotherapy boost: A dose escalated external beam radiotherapy boost to intraprostatic tumour nodules within the prostate gland; Follow Up Length: 60 month(s); Study Entry : Registration only

Original interventions:

100 patients to be recruited and have MRI scans to diagnose intra-prostatic tumour nodules. 50% of patients expected to have lesions and so 50 patients to be treated with a radiotherapy boost

Radiotherapy boost: A dose escalated external beam radiotherapy boost to intraprostatic tumour nodules within the prostate gland; Follow Up Length: 60 month(s); Study Entry : Registration only

Intervention Type

Other

Phase Phase II

Primary outcome measure

Late rectal toxicity; Timepoint(s): 12 months

Secondary outcome measures

- 1. Acute genitourinary (Gu) and gastrointestinal (GI) toxicity; Timepoint(s): 18 weeks
- 2. Biochemical Recurrence; Timepoint(s): 24 months
- 3. Late GU and GI toxicity; Timepoint(s): 12 months and 24 months
- 4. Quality of Life Scores; Timepoint(s): 24 months

Overall study start date

13/07/2011

Completion date

31/12/2023

Eligibility

Key inclusion criteria

- 1. Age more than or equal to 18 years
- 2. Histologically confirmed adenocarcinoma of the prostate

3. National Collaborative Cancer Network+ (NCCN) risk groups intermediate or high risk localised prostate cancer

4. Normal blood count [haemoglobin (Hb) > 11g/dl, white blood cell (WBC) > 4000/mm³, platelets > 100,000/mm³]

- 5. World Health Organisation (WHO) performance status 0 or 1
- 6. Life expectancy of 10 years or more
- 7. Written informed consent
- 8. Patients must be prepared to attend follow-up

9. For template biopsy sub-study must be considered fit for general / spinal anaesthetic; Target Gender: Male ; Lower Age Limit 18 no age limit or unit specified

Participant type(s)

Patient

Age group

Adult

Lower age limit 18 Years

Sex

Male

Target number of participants

Planned Sample Size: 350; UK Sample Size: 350

Total final enrolment

265

Key exclusion criteria

- 1. Prior radiotherapy to the prostate or pelvis
- 2. Bilateral hip replacement
- 3. Prior hormone therapy
- 4. Radical prostatectomy
- 5. Lymph Node Risk > 30%
- 6. National Collaborative Cancer Network+ (NCCN) Favourable Risk Group
- 7. Evidence of seminal vesicle invasion, nodal or metastatic disease

8. Any previous invasive cancer in the past 5 years, with the exception of non-melanoma skin cancer

9. Patients with medical contraindication to magnetic resonance imaging (MRI) scanning

Date of first enrolment

13/07/2011

Date of final enrolment 30/10/2018

Locations

Countries of recruitment England

United Kingdom

Study participating centre Downs Road Sutton United Kingdom SM2 5PT

Sponsor information

Organisation Institute for Cancer Research (UK)

Sponsor details Section of Clinical Trials 15 Cotswold Road Sutton United Kingdom SM2 5NG

Sponsor type Research organisation

ROR https://ror.org/043jzw605

Funder(s)

Funder type Charity

Funder Name Cancer Research UK (CRUK) (UK)

Alternative Name(s) CR_UK, Cancer Research UK - London, CRUK

Funding Body Type Private sector organisation

Funding Body Subtype Other non-profit organizations

Location United Kingdom

Results and Publications

Publication and dissemination plan Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		20/09/2022	29/09/2022	Yes	No