Endoscopic versus surgical drainage of the pancreatic duct in chronic pancreatitis: a prospective randomised trial

Submission date	Recruitment status No longer recruiting	Prospectively registered		
20/12/2005		Protocol		
Registration date	Overall study status	Statistical analysis plan		
20/12/2005	Completed	[X] Results		
Last Edited 17/09/2008	Condition category Digestive System	[] Individual participant data		
11/02/2000	DIACOURE DAOLEIII			

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Djuna Cahen

Contact details

Academic Medical Center Meibergdreef 9 Amsterdam Netherlands 1105 AZ +31 (0)20 3474723 Djunacahen@hotmail.com

Additional identifiers

Protocol serial number N/A

Study information

Scientific Title

Acronym

CEPAN

Study objectives

To compare endoscopic and surgical drainage of the pancreatic duct.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received from the local medical ethics committee

Study design

Randomised, active controlled, parallel group trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Chronic pancreatitis (CP)

Interventions

- 1. Surgical drainage: pancreaticojejunostomy
- 2. Endoscopic drainage: extracorporeal shock wave lithotripsy (ESWL) and/or pancreatic stenting

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Mean Izbicki pain score during follow-up

Key secondary outcome(s))

Clinical success:

- 1. Pain relief at end of FU
- 2. Complete (Izbicki pain score less than or equal to 10)
- 3. Partial (greater than 50% decrease, total score greater than 10)
- 4. Morbidity and mortality rate
- 5. Intervention rate
- 6. Hospital stay
- 7. Development of endo- and/or exocrine pancreatic insufficiency

Completion date

01/10/2004

Eligibility

Key inclusion criteria

- 1. A diagnosis of CP, based on clinical symptoms in combination with morphological changes established by imaging studies (calcifications or ductal changes) and/or pancreatic functional insufficiency
- 2. A dominant obstruction of the pancreatic duct, demonstrated by presence of a stenosis and /or intraductal stones on magnetic resonance cholangiopancreatography (MRCP) and abdominal computed tomography (CT) scan, located left from the spine, with pre-stenotic ductal dilatation of at least 5 mm
- 3. Severe recurrent pancreatic pain with insufficient relieve by non-narcotic analgesics or requiring opiates

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Not Specified

Key exclusion criteria

- 1. Age below 18 or over 80 years
- 2. Pancreatic head enlargement greater than 4 cm
- 3. Contra-indication for surgery; American Society of Anaesthesiologists (ASA) class 4, severe portal hypertension
- 4. Contra-indications for endoscopy: gastrectomy with Billroth II reconstruction, other pancreatitis related complications (bile duct stricture, pseudocyst) requiring surgery
- 5. Previous pancreatic surgery
- 6. Suspected pancreatic malignancy
- 7. Limited life expectancy (less than 2 years)
- 8. Pregnancy

Date of first enrolment

01/01/2000

Date of final enrolment

01/10/2004

Locations

Countries of recruitment

Netherlands

Study participating centre Academic Medical Center Amsterdam Netherlands 1105 AZ

Sponsor information

Organisation

Academic Medical Centre (AMC) (Netherlands)

ROR

https://ror.org/03t4gr691

Funder(s)

Funder type

Industry

Funder Name

AstraZeneca (The Netherlands) - unrestricted grant

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	Results	15/02/2007		Yes	No