

Use of [-2]proPSA for the detection of prostate cancer in patients who are candidate for prostatic biopsy

Submission date 10/11/2011	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 19/12/2011	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 03/11/2015	Condition category Cancer	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

The aim of this study is to investigate how well a new biomarker (a substance used as an indicator of the presence of prostate cancer) works to improve the detection of prostate cancer and reduce the number of negative prostate biopsies (where tissue is unnecessarily removed as it is not cancerous).

Who can participate?

Men from the participating institution scheduled for a prostate biopsy (removal of cancerous tissue), based on clinician decision.

What does the study involve?

One single blood sample was taken from those participating (which is already part of the standard care, for patients scheduled for prostate biopsy). All patients will receive the appropriate treatment according to their condition and the study will have no influence of any kind on the choice of treatment offered to the patients.

What are the possible benefits and risks of participating?

Development of the new biomarker could improve detection of prostate cancer and reduce the number of negative prostate biopsies that are common with existing tests. This new biomarker can improve patient care by reducing the number of invasive and costly procedures. There are no known risks associated with this study.

Where is the study run from?

San Raffaele Hospital-Turro , Milan, Italy (Lead center).

Other centers: Hopital Henri Mondor, Paris, France; Fundacio Puigvert, Barcelona, Spain; University Hamburg-Eppendorf, Hamburg, Germany; Lister Hospital, Stevenage, UK

When is study starting and how long is it expected to run for?

Recruitment started in December 2011 and ended in March 2012. The study ran for 9 months.

Who is funding the study?
Beckman Coulter, Switzerland

Who is the main contact?
Ms Amy Sussman
sussman.amy@hsr.it

Contact information

Type(s)
Scientific

Contact name
Prof Gorgio Guazzoni

Contact details
Department of Urology
San Raffaele Turro Hospital
Via Stamira d'Ancona 20
20127 Milan Italy
Milan
Italy
20127
+39 (0)226 436 431
sussman.amy@hsr.it

Additional identifiers

Protocol serial number
2PROPSA

Study information

Scientific Title
Evaluation of the of the sensitivity, specificity and diagnostic accuracy of the [-2]proPSA biomarker and its derivatives for the detection of prostate cancer and determination of the relationship with prostate cancer characteristics at biopsy

Study objectives
The two most commonly used tests for prostate cancer detection are digital rectal examination and serum prostate-specific antigen (PSA) level. In the last two decades, PSA has considerably improved our ability to detect prostate cancer (PCa). However, this marker has some limitations in sensitivity and specificity for prostate cancer detection leading to false negative and false positive results. Thus, in the recent years, several promising biomarkers have been studied in order to improve PSA specificity in the early detection of PCa. Preliminary reports have shown that some PSA precursors such as [-2]proPSA may be more specific than total PSA or free PSA in predicting the presence of PCa especially in the PSA range between 2 and 10 ng/mL. The goal of this trial is to test the ability of [-2]proPSA and its derivatives in discriminating between patients with or without PCa within a prospectively collected, multicentric, European, large and contemporary cohort of candidates for a prostate biopsy.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Medical Ethics Committee, San Raffaele Hospital Milan, Italy, 08/04/2010, ref: 2PROPSA

Study design

Non-randomized observational multicenter trial

Primary study design

Observational

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Prostate cancer

Interventions

For each patients included, as part of the standard care at each institution, blood will be drawn and serum (blood centrifuged and serum collected) will be prepared within 3 hours of the blood draw. The serum samples will be coded so that no patients name or details will be recorded. The serum samples will be then frozen at -20°C or -70°C. The samples will be then sent to Hospital San Raffaele in Milan for the determination of the PSA, fPSA and [-2]proPSA serum concentrations and calculation of the prostate health index (phi). The patients clinical information including diagnostic will be collected and coded at each institution and sent to the data manager at Hospital San Raffaele. The statistical analysis will be then performed at Hospital San Raffaele in Milan on the complete data set including laboratory information (PSA, fPSA, [-2] proPSA, phi) and clinical information. The performance of the new diagnostic test will be assessed in comparison with currently used test such as tPSA or %fPSA using ROC curve analysis, univariate and multivariate logistic regression and determination of gain in clinical specificity at a given sensitivity. Each patients included in the trail will be followed at each institution for a period of 6 months following the initial blood draw.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Evaluate the specificity, sensitivity and diagnostic accuracy of -2proPSA and its derivatives (index text) in determining the presence of PCa at prostate biopsy and to compare them with those of total and free PSA (referenced text). Thus, we will quantify the number of prostate biopsies that can be spared using -2proPSA in the prostate biopsy decision path.

Key secondary outcome(s)

1. Determine a -2proPSA (and its derivatives) cut-off value to be used in clinical practice
2. Determine the relationship between -2proPSA (and its derivatives) and PCa characteristics at biopsy (e.g. Gleason grade, % of cores involved, % of involvement of a single core)

Completion date

31/03/2012

Eligibility

Key inclusion criteria

1. Male patients older than 45 years old
2. Negative/positive digital rectal examination
3. Patients subjected to previous prostate biopsy will be included and analyzed in a nested case control study
4. Patients treated with drugs that may alter serum PSA levels (Finasteride and Dutasteride) will be included and analysed later in a nested case control study

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Male

Key exclusion criteria

1. Patients with bacterial acute prostatitis and with a positive sperm culture in the last three months prior to biopsy
2. Patients subjected to previous endoscopic surgery of the prostate (TransUrethral Resection of the Prostate [TURP] or Holmium-laser Enucleomorcellation of the Prostate [HoLEP])

Date of first enrolment

01/12/2011

Date of final enrolment

31/03/2012

Locations

Countries of recruitment

United Kingdom

France

Germany

Italy

Spain

Study participating centre
San Raffaele Turro Hospital
Milan
Italy
20127

Sponsor information

Organisation
Beckman Coulter Eurocenter (Switzerland)

Funder(s)

Funder type
Industry

Funder Name
Beckman Coulter Eurocenter S.A. (Switzerland)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/11/2014		Yes	No