

Preterm Premature Rupture Of Membranes between 34 and 37 weeks: EXpectant management versus Induction of Labour 2

Submission date 31/01/2010	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 07/06/2010	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 29/12/2020	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Ms Jantien van der Heyden

Contact details
De Run 4600
Veldhoven
Netherlands
5500 MB

-
j.vanderheyden@mmc.nl

Additional identifiers

Protocol serial number
N/A

Study information

Scientific Title

The effectiveness of immediate delivery after preterm prelabour rupture of membranes between 34 and 37 weeks compared to expectant management in a multicentre randomised controlled trial

Acronym

PPROMEXIL 2

Study objectives

Induction of labour in patients with preterm premature rupture of membranes (PPROM) between 34 and 37 weeks' gestation will reduce the incidence of neonatal sepsis. This advantage may outweigh the effects of prematurity (e.g. respiratory distress syndrome and hypoglycaemia)

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics committee of the University Hospital Maastricht

Study design

Multicentre randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Preterm prelabour rupture of membranes

Interventions

Participants will be randomised to induction of labour or expectant management. If randomised for induction: same day or the day after. If randomised for expectant management: induction at 37 weeks' gestation. Duration of treatment: Maximum 3 weeks. The follow-up will be done after 6 weeks, 6 months and 2 years.

The analysis will be done by intention to treat. Relative risks and 95% confidence intervals will be calculated for the relevant outcome measures. The analysis will be stratified for centre and parity. In case of equivalence between outcomes, the analysis will be repeated on a per protocol basis. Quality of life as well as pain scores will be analysed using repeated measures analysis of variance. Also a cost-effectiveness analysis will be done.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Neonatal sepsis, measured immediately on discharge from hospital

Key secondary outcome(s)

1. Respiratory distress syndrome (RDS)
2. Transient tachypnoea of the newborn
3. Asphyxia
4. Pneumothorax/pneumomediastinum
5. Late onset sepsis
6. Hypoglycaemia
7. Meconium aspiration syndrome
8. Necrotising enterocolitis (NEC)
9. Hyperbilirubinaemia
10. Intraventricular haemorrhage
11. Periventricular leucomalacia
12. Convulsions
13. Other neurological abnormalities and congenital abnormalities

Secondary maternal outcome measures:

1. Antepartum haemorrhage
2. Umbilical cord prolapse
3. Signs of chorioamnionitis
4. Maternal sepsis
5. Thrombo-embolic complications
6. Urinary tract infection treated with antibiotics
7. Signs of endometritis
8. Pneumonia
9. Anaphylactic shock
10. Haemolysis, Elevated Liver Enzymes, Low Platelets (HELLP) syndrome
11. Death
12. Incidence of instrumental deliveries
13. Maternal quality of life
14. Maternal intervention reference
15. Costs

Other outcomes:

Direct medical and non-medical costs, generated by maternal and neonatal resource utilisation during admission and post-discharge follow-up until 6 weeks after randomisation. The economic evaluation will integrate the primary clinical outcome and costs in a cost effectiveness analysis.

All outcomes will be measured immediately on discharge from hospital, and also 6 weeks and 6 months post-partum.

Completion date

31/12/2010

Eligibility

Key inclusion criteria

1. Women presenting with preterm prelabour rupture of the foetal membranes between 34+0 and 37+0 weeks' gestation and have not delivered within 24 hours after rupture of the foetal membranes

3. Women presenting with preterm prelabour rupture of foetal membranes after 26+0 weeks gestation who have not delivered at 34+0 weeks of gestation
4. Single and multiple gestations
5. Women with child in breech presentation can also be included

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Total final enrolment

195

Key exclusion criteria

1. Monochorionic multiple pregnancies
2. Abnormal (non-reassuring) cardiotocogram (CTG)
3. Meconium stained amniotic fluid
4. Signs of intrauterine infection
5. Major foetal anomalies
6. Being in labour
7. Hemolytic anaemia, elevated liver enzymes and low platelet count (HELLP) syndrome
8. Severe pre-eclampsia

Date of first enrolment

01/01/2010

Date of final enrolment

31/12/2010

Locations**Countries of recruitment**

Netherlands

Study participating centre

De Run 4600

Veldhoven

Netherlands

5500 MB

Sponsor information

Organisation

Academic Medical Centre (AMC) (Netherlands)

ROR

<https://ror.org/03t4gr691>

Funder(s)

Funder type

Government

Funder Name

The Netherlands Organisation for Health Research and Development (ZonMw) (Netherlands)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/10/2012	29/12/2020	Yes	No
Other publications	secondary analysis	01/09/2014		Yes	No
Other publications	prediction model development	01/05/2014	29/12/2020	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes