

Reducing adverse electroconvulsive treatment effects on memory by magnetic stimulation

Submission date 19/04/2006	Recruitment status Stopped	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 07/06/2006	Overall study status Stopped	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 06/10/2009	Condition category Signs and Symptoms	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Prof Klaus Ebmeier

Contact details
Kennedy Tower
Morningside Park
Edinburgh
United Kingdom
EH10 5HF
+44 (0)131 5376505
k.ebmeier@ed.ac.uk

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
G0401083

Study information

Scientific Title

Study objectives

We propose a two-year pilot randomised controlled trial of electroconvulsive treatment (ECT) versus magnetic seizure therapy (MST) in 80 Edinburgh patients recruited from 100 new treatment courses started per year in Edinburgh (75 after giving informed consent) to examine the following questions:

1. Is MST less liable than ECT to cause anterograde and retrograde memory impairment and what is the likely size of the effect?
2. Is MST equally effective to ECT within the power of a moderately large randomised study with blind evaluation of symptom change?
3. Is MST more user-friendly and user-acceptable than ECT?
4. What is the balance of cost versus benefit comparing ECT and MST in patients referred for ECT?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval not yet received as of 07/06/06

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Not specified

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Memory impairment

Interventions

1. Randomisation to either ECT or MST groups
2. ECT protocol (treatment as usual): ECT will be administered in line with the latest guidelines from the Royal College of Psychiatrists. Seizure threshold will be measured at the outset of treatment, bilateral ECT given with a dose 50% above threshold, and right unilateral ECT given with a dose initially 300% above threshold. Treatment will be given with a modern constant

current ECT machine using doses of 100-400 mC (mode = 150 mC, 800 mA, 20-120 Hz; pulse width 1 ms; MECTA Spectrum™ 5000 M), and monitored by electroencephalogram (EEG). The clinical team responsible for the patient will determine the need for and duration of treatment, usually between 6 and 12 treatments.

3. MST protocol: MST will be administered mirroring the dose titration process above. Prefrontal coil placement will be used with a stimulation frequency between 50 and 100 Hz at various output strengths. Seizures will be monitored using EEG (split electrodes to prevent heating). The clinical team responsible for the patient will determine the need for and duration of treatment, usually between 6 and 12 treatments. The responsible team can request exit from the protocol and transfer to ECT, if there are clinical concerns, such as deterioration or emerging suicidality.

Added 06/10/09: the trial was stopped due to participant recruitment issues.

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

Last follow-up six months after course of treatment of last recruited subject recruited before 31st March 2008 or 80th subject recruited

Secondary outcome measures

Not provided at time of registration

Overall study start date

01/04/2006

Completion date

31/05/2008

Reason abandoned (if study stopped)

Participant recruitment issue

Eligibility

Key inclusion criteria

1. Referred to and accepted by ECT service for treatment of major depressive episode
2. Able to give informed consent to ECT and to trial procedure
3. If patient is detained, ECT would be given with patient's consent (valid T2 form completed)

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

2 x 40

Key exclusion criteria

1. Contraindications for ECT or anaesthesia
2. Unable or unwilling to give informed consent to ECT or to trial procedure
3. Patients with organic diagnoses (e.g. dementia, schizophrenia and substance abuse)
4. Patients with metallic implants or pacemakers
5. Pregnancy
6. Aged <18 years

Date of first enrolment

01/04/2006

Date of final enrolment

31/05/2008

Locations**Countries of recruitment**

Scotland

United Kingdom

Study participating centre

Kennedy Tower

Edinburgh

United Kingdom

EH10 5HF

Sponsor information**Organisation**

University of Edinburgh (UK)

Sponsor details

The University of Edinburgh

Old College

South Bridge

Edinburgh

Scotland

United Kingdom

EH8 9YL

Sponsor type

University/education

Website

<http://www.ed.ac.uk/contact.html>

ROR

<https://ror.org/01nrxf90>

Funder(s)

Funder type

Government

Funder Name

The Medical Research Council (MRC) (UK)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration