Re-stitching of a broken down perineal wound compared to leaving it to heal naturally

Submission date 08/12/2008	Recruitment status No longer recruiting
Registration date 16/01/2009	Overall study status Completed
Last Edited 20/10/2017	Condition category Pregnancy and Childbirth

[X] Prospectively registered

[X] Protocol

[] Statistical analysis plan

[X] Results

[] Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s) Scientific

Contact name Miss Lynn Dudley

Contact details

University Hospital of North Staffordshire NHS Trust Treatment Centre, Children's Services and Obstetrics Academic Unit of Obstetrics and Gynaecology Maternity Building, City General Site Newcastle Road Stoke-on-Trent Staffordshire United Kingdom ST4 6QG +44 (0)1782 552 058/434 lynn.dudley@uhns.nhs.uk

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Study information

Scientific Title

Perineal REpair following Vaginal delivery complicated by an Infected Episiotomy Wound: a feasibility study for a randomised controlled trial

Acronym

PREVIEW

Study objectives

Many women will require suturing to facilitate healing of the trauma site. However, practice varies widely between care given and established professionally agreed standards. There is limited data on the prevalence and consequences of perineal wound infection. In addition, there is only a small amount of information relating to the impact that perineal wound infection has on women's well-being during the immediate and long-term post-natal period. Anecdotal evidence suggests the number of women reporting perineal infections and dehiscence in the community is increasing; however, systems to track these complications following hospital discharge are lacking. Given that postpartum management of perineal trauma is a core component of routine maternity care it is vital that a true estimate of the problem is established using standardised definitions of wound infection and at the same time determine best practice when treating dehisced perineal wounds.

Hypotheses:

1. What is the prevalence of perineal wound infection and dehiscence in the UK?

2. What factors at the time of primary repair are most likely to be associated with perineal wound infection and dehiscence prior to discharge to the community?

3. What factors following discharge home are most likely to be associated with perineal wound infection and dehiscence in the community?

4. What are women's experiences of perineal infection and dehiscence and what types of information and support are most likely to benefit their post-natal recovery?

5. What is the best management for perineal wound infection and wound dehiscence?
6. What is the feasibility of conducting a definitive randomised controlled trial (RCT) comparing re-suturing of dehisced perineal wounds versus healing by secondary intention and what are the implications in terms of health benefits and costs?

Ethics approval required

Old ethics approval format

Ethics approval(s) North Wales Research Ethics Committee, 29/04/2010, ref: 10/WNO03/16

Study design

Computer-randomised controlled feasibility study

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s) Treatment

Participant information sheet

Not available in web format, please use the contact details to request a patient information sheet

Health condition(s) or problem(s) studied

Dehisced perineal wounds

Interventions

The participants will be computer randomised into either immediate resuturing of their dehisced perineal wound in comparison to healing by secondary intention. Both groups will have an independent assessment of their perineal wound at 2 weeks and 6 weeks after trial entry. Both groups of participants will be asked to complete a questionnaire at 6 weeks, 3 months and 6 months after trial entry.

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

Time taken for the dehisced perineal wound to heal, assessed at 2 weeks, 6 weeks, 3 months and 6 months, respectively.

An independent assessment of the primary outcome will be conducted at 2 weeks and 6 weeks using a questionnaire format, the title of the questionnaire being: 'PREVIEW' Independent Perineal Assessment 2 weeks, and 'PREVIEW' Independent Assessment 6 weeks

Mothers will also complete questionnaires at 6 weeks, 3 months and 6 months, respectively, also assessing primary and secondary outcomes, the title of these questionnaires being: 'PREVIEW' Mothers questionnaire 6 weeks 'PREVIEW' Mothers questionnaire 3 months 'PREVIEW' Mothers questionnaire 6 months

Secondary outcome measures

- 1. Woman's satisfaction with aesthetic results of perineal wound at 6 months post-natal
- 2. Pain at 6 weeks, 3 and 6 months post-natal
- 3. Dyspareunia (painful intercourse) at 3 6 months
- 4. Rates of breastfeeding

Mothers will complete questionnaires at 6 weeks, 3 months and 6 months respectively, assessing primary and secondary outcomes, the title of these questionnaires being: 'PREVIEW' Mothers questionnaire 6 weeks 'PREVIEW' Mothers questionnaire 3 months 'PREVIEW' Mothers questionnaire 6 months

Overall study start date

01/04/2009

Completion date

01/04/2011

Eligibility

Key inclusion criteria

1. Women (aged 18 - 40 years) referred to the perineal care clinic at the University Hospital of North Staffordshire

2. Dehisced perineal wound (spontaneous second, third or fourth tear or episiotomy)

3. Occurs within two weeks following childbirth

Participant type(s)

Patient

Age group Adult

Lower age limit

Upper age limit

40 Years

Sex

Female

Target number of participants

Approximately 250 patients

Key exclusion criteria

- 1. Women that have not given their written consent to participate in the study
- 2. Women who have delivered a stillborn infant
- 3. Women under the age of 16 years
- 4. Women who cannot speak English or cannot read or write

Date of first enrolment

01/04/2009

Date of final enrolment 01/04/2011

Locations

Countries of recruitment England

United Kingdom

Study participating centre University Hospital of North Staffordshire NHS Trust Staffordshire United Kingdom ST4 6QG

Sponsor information

Organisation University Hospital of North Staffordshire NHS Trust (UK)

Sponsor details

c/o Darren Clement Research and Development Manager Research and Development Department North Staffordshire Medical Institute Hartshill Road Hartshill, Stoke-on-Trent Staffordshire England United Kingdom ST4 7NY

Sponsor type Hospital/treatment centre

Website http://www.uhns.nhs.uk/

Funder(s)

Funder type Charity

Funder Name

Smith and Nephew Foundation

Alternative Name(s)

Funding Body Type Private sector organisation

Funding Body Subtype Trusts, charities, foundations (both public and private)

Location United Kingdom

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<u>Protocol article</u>	protocol	24/07/2012		Yes	No
<u>Results article</u>	nested qualitative study results	10/02/2017		Yes	No
<u>Results article</u>	results	10/02/2017		Yes	No