

# The Efficacy of Acupuncture in Chronic Rotator Cuff Tendinitis

<b>Submission date</b> 23/11/2009	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 18/03/2010	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
<b>Last Edited</b> 11/08/2011	<b>Condition category</b> Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Victoria Bombin

**Contact details**  
Ancaster Sports Medicine Clinic  
1015 Golflinks Road  
Ancaster  
Canada  
L9K 1L6

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**  
0.01

## Study information

**Scientific Title**

# The Efficacy of Acupuncture in Chronic Rotator Cuff Tendinitis: A Pilot Randomised Controlled Trial

## Study objectives

Is acupuncture is an effective method of treating chronic rotator cuff tendinitis?

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

1. Hamilton Health Science Research Ethics Board, McMaster University, Canada approved on the 26th of February 2010 (ref: 09-607)
2. Approval pending from the Institutional Review Board, Mayo Clinic, USA

## Study design

Multicentre 3 arm randomised active and placebo controlled double blinded study

## Primary study design

Interventional

## Secondary study design

Randomised controlled trial

## Study setting(s)

Hospital

## Study type(s)

Treatment

## Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

## Health condition(s) or problem(s) studied

Chronic rotator cuff tendinitis

## Interventions

After informed consent, patients will be randomised to one of three groups.

1. Traditional needle acupuncture
2. Laser acupuncture
3. Sham laser acupuncture

On the same day, these patients will have a history taken and a physical examination performed to exclude other causes of shoulder pain. Each patient will be examined by two independent physicians. Patients who meet the diagnostic criteria of rotator cuff tendinitis as assessed by the two physicians will be selected to enter the study. Disagreements between these two physicians will be resolved at the committee meeting. If the disagreement is not resolved, a third opinion by a qualified physician will be obtained. Other patients who do not meet the criteria will be excluded and referred to their family physician.

All subjects are then given education about Codman pendulum shoulder exercises to prevent potential shoulder stiffness. Patients are strongly encouraged to continue to take ibuprofen 200

to 600 mg every 6 hours or acetaminophen 320 to 500 mg every 4 hours on an as needed basis. These instructions will be given by one of the evaluating physicians.

The patients in all three groups will have ten sessions of metal needle, laser, or sham laser acupuncture treatments over a five week period by the same treating physician as described in the blinding section. Ten treatments are used because of previous systematic review suggests that this is the minimum number of treatments needed to produce sustained clinical effects.

#### **Traditional Acupuncture Group:**

Patients in this group will have needle insertion to a depth of 10-35 mm. The needle is then manually twirled for 30 seconds or until patients feel a dull pressure and warmth feeling around the needle (deqi). The treatment sessions will last 20 minutes with needles manipulated at the 10 minute mark.

#### **Laser groups:**

A research assistant not involved in the treatment or data analysis will enter the room prior to each laser intervention and flip the switch to either real treatment or placebo according to the group the patient has been randomly assigned to. The patient and the treating physician will be blinded to this procedure and will enter the room only after the research assistant exits the area. The laser leads will then be applied to the same points used for the metal needle acupuncture group. The patient will remain hooked up to the machine for twenty minutes to produce 20 joules energy per point at 810 nm infrared laser.

The Shoulder Pain and Disability Index and the Oxford Analgesic Chart will be given before each treatment by the blinded research assistant. The assistants will also contact the patients at three and six months after completing their course of treatments. Each patient will keep a log of the amount of ibuprofen and acetaminophen used using the Oxford analgesic chart, which will be confirmed by the research assistant weekly. The patient will also fill out a form at the end of the treatment to document any adverse effects noted during treatment.

### **Intervention Type**

Other

### **Phase**

Not Applicable

### **Primary outcome measure**

Physical function and pain status, assessed by the Shoulder Pain and Disability Index (SPADI) at baseline, 10 weeks (end of intervention) and 1 month post-intervention.

### **Secondary outcome measures**

Amount of analgesics consumed, assessed by Oxford analgesic chart at baseline, 10 weeks (end of intervention) and 1 month post-intervention.

### **Overall study start date**

02/01/2010

### **Completion date**

30/06/2010

## **Eligibility**

### **Key inclusion criteria**

1. Age 18-70
2. Shoulder pain with abduction and aggravated by resistance testing (abduction, internal rotation, or external rotation)
3. Duration of pain  $\geq$  3 months
4. Willingness to travel to treatment centres and to be available for repeated follow-up phone calls

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

18 Years

**Upper age limit**

70 Years

**Sex**

Both

**Target number of participants**

30 at each site: 60 total

**Key exclusion criteria**

1. Receiving other modality of treatment aside from analgesics or non-steroidal anti-inflammatory drugs (NSAIDs)
2. Patients who have contraindication to NSAIDs such as ulcers or gastroesophageal bleeding
3. Unsettling legal or worker compensation claims

**Date of first enrolment**

02/01/2010

**Date of final enrolment**

30/06/2010

**Locations****Countries of recruitment**

Canada

United States of America

**Study participating centre**

**Ancaster Sports Medicine Clinic**  
Ancaster  
Canada  
L9K 1L6

## **Sponsor information**

### **Organisation**

Ancaster Sports Medicine Clinic (Canada)

### **Sponsor details**

c/o Dr. Victoria Bombin  
1015 Golflinks Road  
Ontario  
Ancaster  
Canada  
L9K 1L6

### **Sponsor type**

Hospital/treatment centre

## **Funder(s)**

### **Funder type**

Industry

### **Funder Name**

Magna Corporation (Canada)

## **Results and Publications**

### **Publication and dissemination plan**

Not provided at time of registration

### **Intention to publish date**

### **Individual participant data (IPD) sharing plan**

### **IPD sharing plan summary**

Not provided at time of registration