# Targeting synovitis in knee osteoarthritis

Submission date	Recruitment status  No longer recruiting	Prospectively registered			
21/05/2010		☐ Protocol			
Registration date	Overall study status	Statistical analysis plan			
21/05/2010	Completed	[X] Results			
<b>Last Edited</b> 10/05/2017	Condition category  Musculoskeletal Diseases	[] Individual participant data			

#### Plain English summary of protocol

Background and study aims

Osteoarthritis (OA) is the most common form of arthritis and remains one of the few chronic diseases of ageing for which there is little, if any, effective treatment. Symptomatic knee osteoarthritis affects roughly 12% of people aged 60 and over and, despite medical advances, remains a major source of pain and functional limitation. While the main feature of OA is loss of cartilage (the protective surface that allows your joints to move smoothly), all parts of the joint are affected. Treatment development has focused on protecting the cartilage from damage. Another important potential source of OA pain is the lining of the joint (the synovium). Inflammation of the synovium is seen in about 50% of the knees of patients with painful OA. Since corticosteroid drugs reduce inflammation, injection of corticosteroids into the knee might be effective for the treatment of OA pain. The aim of this study is to determine whether patients with evidence of synovial inflammation in OA respond better to corticosteroids than those who do not, and whether treatment response is associated with change in synovial inflammation.

Who can participate?

Patients aged 40 - 79 years with symptomatic knee osteoarthritis (OA).

What does the study involve?

Participants have their knee aspirated (where a sterile needle and syringe are used to drain fluid) and injected with the corticosteroid drug methylprednisolone acetate (Depo-Medrone). We measure the changes in knee pain and inflammation of the synovium following the injection.

What are the possible benefits and risks of participating? Not provided at time of registration

Where is the study run from? University of Manchester (UK)

When is the study starting and how long is it expected to run for? May 2010 to May 2012

Who is funding the study?
Arthritis Research Campaign (ARC) (UK)

Who is the main contact?
Helen Williams
Helen.Williams@manchester.ac.uk

## Contact information

### Type(s)

Scientific

#### Contact name

Mrs Helen Elizabeth Williams

#### Contact details

University of Manchester ARC Epidemiology Unit Stopford Building Oxford Road Manchester United Kingdom M13 9PT

\_

Helen.Williams@manchester.ac.uk

## Additional identifiers

Clinical Trials Information System (CTIS) 2009-015849-22

Protocol serial number

8358

# Study information

#### Scientific Title

An open-label study of intra-articular steroid injection in the management of symptomatic knee osteoarthritis

#### Acronym

TASK v1.2

### Study objectives

Osteoarthritis (OA) of the knee is a common disorder and at present there are few effective therapies. Intra-articular (IA) steroids have been used in the management of knee OA, however, the results of these studies have been variable. The aim of this study is to determine factors which predict the improvement in pain in patients treated with IA steroids. In total 120 patients with knee OA will be studied in an open label study of IA steroids.

The primary outcome measure will be the change in knee pain following steroid injection. We plan to determine whether features of synovitis at baseline assessed both clinically and on

magnetic resonance imaging (MRI) imaging predict response to steroids and whether change in features of synovitis, assessed using serial MRI scanning are associated with change in pain status. We will also determine the predictive ability of synovial fluid parameters and other factors including quality of life predict the response to steroids.

### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Leicester 2 REC, 27/01/2010, ref: 09/H0402/107

#### Study design

Single-centre non-randomised interventional treatment trial

#### Primary study design

Interventional

#### Study type(s)

Treatment

#### Health condition(s) or problem(s) studied

Topic: Musculoskeletal, Primary Care Research Network for England; Subtopic: Not Assigned, Musculoskeletal (all Subtopics); Disease: Musculoskeletal, All Diseases

#### **Interventions**

Participants will have their index knee aspirated and injected with 80 mg methylprednisolone acetate (Depo-Medrone). Subjects will have one injection only during the course of the study.

Follow up length: 6 months Study entry: registration only

### Intervention Type

Drug

#### Phase

Phase IV

## Drug/device/biological/vaccine name(s)

Depomedrone

## Primary outcome(s)

Change in knee pain, analysis will focus on whether change in pain level from baseline to one week after the injection is given

### Key secondary outcome(s))

- 1. Visual Analogue Scale (VAS) global knee pain
- 2. VAS pain on nominated activity
- 3. VAS wellness
- 4. Knee injury and Osteoarthritis Outcome Score (KOOS) subscales

Data collected at each study visit, baseline, week 1 and 1 x follow-up.

#### Completion date

15/12/2014

# **Eligibility**

### Key inclusion criteria

- 1. Age 40 79 years
- 2. Male or female
- 3. Ambulatory (not wheelchair bound)
- 4. Able and willing to attend or comply with intervention and follow up
- 5. Within the last 24 months:
- 5.1. Radiological (X-ray) evidence of grade 2 or more OA
- 5.2. Evidence of significant OA on MRI scan
- 5.3. Documented evidence of at least grade 2 arthritis on arthroscopy
- 6. Moderate knee pain lasting 48 hours in the past 2 weeks
- 7. Presence of clinically apparent knee effusion
- 8. Written informed consent
- 9. Glomerular filtration rate (GFR) greater than 44 ml/min

#### Participant type(s)

**Patient** 

### Healthy volunteers allowed

No

#### Age group

Adult

#### Sex

ΔII

#### Key exclusion criteria

- 1. Secondary OA septic arthritis, gout
- 2. History of inflammatory arthritis
- 3. Previous intra-articular Ostenil or steroid injection within 6 months
- 4. Previous knee surgery (including cartilage surgery) or arthroscopy within 6 months
- 5. Inability to understand the procedures
- 6. Pregnancy
- 7. Chronic kidney disease with estimated glomerular filtration rate (eGFR) greater than or equal to 44 ml/min
- 8. Concurrent life threatening illness
- 9. Implants which prohibit safe use of MRI scan including cochlear implants/metal objects in the body including joint prosthesis, cardiac or neural pacemakers, hydrocephalus shunts, intrauterine device or coil
- 10. Known hypersensitivity to Depomedrone or any components of its excipients
- 11. Systemic infection (unless specific anti infective therapy is employed)

#### Date of first enrolment

04/05/2010

### Date of final enrolment

10/06/2014

## Locations

#### Countries of recruitment

United Kingdom

England

## Study participating centre University of Manchester

Manchester United Kingdom M13 9PT

## Study participating centre Salford Royal NHS Foundation Trust

Stott Lane Salford United Kingdom M6 8HD

# Sponsor information

### Organisation

Salford Royal NHS Foundation Trust (UK)

#### **ROR**

https://ror.org/019j78370

# Funder(s)

# Funder type

Charity

#### **Funder Name**

Arthritis Research Campaign (ARC) (UK)

# **Results and Publications**

# Individual participant data (IPD) sharing plan

# IPD sharing plan summary

Available on request

## **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/01/2016		Yes	No
Results article	results	08/05/2017		Yes	No
HRA research summary			28/06/2023		No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes