The effect of continuous positive airway pressure (CPAP) and non-invasive positive pressure ventilation (NIPPV) in acute cardiogenic pulmonary oedema (ACPO)

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
25/04/2003		Protocol		
Registration date 25/04/2003	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited 02/08/2010	Condition category Circulatory System	[] Individual participant data		

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Alasdair James Gray

Contact details

Emergency Department
Royal Infirmary of Edinburgh
51 Little France Crescent
Old Dalkeith Road
School of Health and Related Research
Edinburgh
United Kingdom
EH16 4SA
+44 (0)131 242 1334
alasdair.gray@luht.scot.nhs.uk

Additional identifiers

Protocol serial number

HTA 01/43/01

Study information

Scientific Title

Acronym

3CPO

Study objectives

To assess the clinical effectiveness of non-invasive ventilation against standard therapy alone

More details can be found at http://www.hta.ac.uk/project/1338 Protocol can be found at http://www.hta.ac.uk/protocols/200100430001.pdf

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not requested at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Cardiovascular diseases: Heart disease

Interventions

Continuous Positive Airway Pressure (CPAP) versus Non-Invasive Positive Pressure Ventilation (NIPPV)

Please note that, as of 28 January 2008, the anticipated end date of this trial was updated from 31 December 2007 to 30 April 2008.

Please note that as of 12th June 2007, the anticipated end date of this trial was extended to 31st December 2007 (Anticipated end date provided at time of registration: 30 June 2006). Recruitment to this trial was closed on 30 April 2007.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Added on the 12th June 2007:

- 1. NIV versus standard therapy: seven-day mortality
- 2. CPAP versus NIPPV: seven-day mortality or intubation

Key secondary outcome(s))

Added on the 12th June 2007:

- 1. Physiology
- 2. Side-effects including Myocardial Infacrtion (MI)
- 3. Cost-effectiveness

Completion date

30/04/2008

Eligibility

Key inclusion criteria

- 1. More than 16 years
- 2. Acute Cardiogenic Pulmonary Oedema (ACPO) principal clinical complaint
- 3. Confirmed diagnosis of ACPO

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Not Specified

Key exclusion criteria

- 1. Patients needing immediate life-saving interventions
- 2. Inability to provide informed consent

Date of first enrolment

01/06/2003

Date of final enrolment

30/04/2008

Locations

Countries of recruitment

United Kingdom

Scotland

Study participating centre Emergency Department Edinburgh United Kingdom EH16 4SA

Sponsor information

Organisation

Department of Health (UK)

ROR

https://ror.org/03sbpja79

Funder(s)

Funder type

Government

Funder Name

NIHR Health Technology Assessment Programme - HTA (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	10/07/2008		Yes	No
Results article	results	01/07/2009		Yes	No
Results article	results	01/08/2010		Yes	No