

Comparison of laparoscopic versus open radical gastrectomy for advanced gastric cancer

Submission date 22/01/2010	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 18/03/2010	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 11/04/2019	Condition category Cancer	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Prof Yu Pei Wu

Contact details
Department of General Surgery and Center of Microinvasive Gastrointestinal Surgery
Southwest Hospital
Chongqing
China
400038

Additional identifiers

ClinicalTrials.gov (NCT)
NCT01043835

Protocol serial number
N/A

Study information

Scientific Title
Comparison of laparoscopic versus open radical gastrectomy for advanced gastric cancer: a prospective randomised controlled trial

Study objectives

The use of laparoscopic surgery in the management of advanced gastric cancer (AGC) has not yet met with widespread acceptance and remains limited to only a few centres. The purpose of this study is to compare the short- and long-term results between the laparoscopy-assisted gastrectomy and the open gastrectomy.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committee of Southwest Hospital, Third Military Medical University, approved on the 24th September 2009 (ref: KY200908)

Study design

Prospective randomised trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Advanced gastric cancer

Interventions

Laparoscopy-assisted gastrectomy:

One initial 10-mm trocar for a laparoscope was inserted below the umbilicus. Another 10-mm trocar was introduced in the left preaxillary line 2 cm below the costal margin as a major hand port. A 5-mm trocar then was inserted in the left midclavicular line 2 cm above the umbilicus as an accessory port, and a 15-mm trocar (also as an accessory port) was placed at the contralateral site, through which a linear cutter was inserted. A 5-mm trocar was inserted in the right preaxillary line 2 cm below the costal margin for traction and exposure of the liver. The operator stood on the left side of the patient. Subtotal or total gastrectomy and D2 lymph node dissection will be performed basically. As a general rule, Billroth I, Billroth II or Roux-Y method was used for gastric reconstruction for all cases. Dissected stomach and lymph node are collected through additional 3 - 5 cm incision at a median superior abdominal incision.

Open gastrectomy:

Approximately 15 - 20 cm length incision is made from falciform process to periumbilical area. Subtotal or total gastrectomy and D2 lymph node dissection will be performed basically. As a general rule, Billroth I, Billroth II or Roux-Y method was used for gastric reconstruction for all cases.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Disease free survival at 3 years

Key secondary outcome(s)

Complications, recurrence, quality of life measured by EORTC QLQ-C30 V 3.0 and EORTC QLQ-STO22 at 3 years

Completion date

31/01/2015

Eligibility

Key inclusion criteria

1. Pathologically proven gastric adenocarcinoma
2. Aged older than 18 years old, younger than 80 years old, either sex
3. Pre-operative stage (computed tomography [CT], gastrofiberscopy [GFS] stage): cT2N0M0, cT2N1M0, cT2N2M0, cT3N0M0, cT3N1M0, cT3N2M0
4. American Society of Anaesthesiologists (ASA) score: less than or equal to 3
5. No invasion of the gastric serosa exceeding 10 cm² according to ultrasound examination or examination during surgery
6. No history of other cancer
7. No history of chemotherapy or radiotherapy
8. Written informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Concurrent cancer patients or patient who was treated due to other types of cancer before the patient was diagnosed as a gastric cancer patient
2. Patient who was treated by other types of treatment methods, such as chemotherapy, immunotherapy, or radiotherapy
3. Patient who was received upper abdominal surgery (except laparoscopic cholecystectomy)
4. ASA score: greater than 3
5. Contraindication of laparoscopy: severe cardiac disease, abdominal wall hernias, diaphragmatic hernias, uncorrected coagulopathies, portal hypertension, pregnancy
6. Complicated case needed to get emergency operation
7. Any accompanying surgical condition needed to be performed in same time

Date of first enrolment

01/02/2010

Date of final enrolment

31/01/2015

Locations

Countries of recruitment

China

Study participating centre

Department of General Surgery and Center of Microinvasive Gastrointestinal Surgery

Chongqing

China

400038

Sponsor information

Organisation

Southwest Hospital (China)

ROR

<https://ror.org/02jn36537>

Funder(s)

Funder type

Government

Funder Name

Chongqing Municipal Government (China) - Science and Technology Research

Results and Publications

Individual participant data (IPD) sharing plan**IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/05/2018	11/04/2019	Yes	No
Results article	results	01/06/2019	11/04/2019	Yes	No