Prediction of Preterm delivery by the Electrohysterogram

Submission date	Recruitment status No longer recruiting	Prospectively registered	
05/12/2011		[X] Protocol	
Registration date	Overall study status	Statistical analysis plan	
31/07/2012	Completed	Results	
Last Edited	Condition category Pregnancy and Childbirth	Individual participant data	
24/02/2015		Record updated in last yea	

Plain English summary of protocol

Background and study aims

The electrohysterogram (EHG) is a new way of measuring contractions of the uterus. In this study the EHG will be used to monitor patients with premature contractions. The aim is to investigate whether the EHG can predict when delivery will occur.

Who can participate?

You can participate if you are admitted the hospital for imminent premature delivery between 24 and 34 weeks of gestation.

What does the study involve?

Besides the usual tests, you will have an additional test: the electrohysterogram (EHG). This means a measurement using a patch that is connected to the skin of your abdomen. The result will only be used for research and not for your treatment.

What are the possible benefits and risks of participating?

The measurement is not dangerous for you or your baby and has no side effects.

Where is the study run from?

The study will be run from the Maxima Medical Center in Veldhoven but four other hospitals in the Netherlands will participate as well.

When is the study starting and how long is it expected to run for? The study started in March 2012 and will end in June 2014.

Who is funding the study?

This research is paid for by the Maxima Medical Center Veldhoven.

Who is the main contact? Hinke de Lau h.delau@mmc.nl

Contact information

Type(s)

Scientific

Contact name

Mr Hinke Lau

Contact details

Tannhauserdreef 272 Utrecht Netherlands 3561HR

Additional identifiers

Protocol serial number

10620

Study information

Scientific Title

Prediction of Preterm delivery by the Electrohysterogram: an observational cohort study

Acronym

PoPE

Study objectives

The electrohysterogram is better predictor of preterm delivery than current diagnostics.

On 30/01/2014 the overall trial end date was changed from 01/02/2013 to 01/06/2014.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Medical Ethical Committee, Maxima Medical Center Veldhoven, 06/12/2011

Study design

Observational multicenter cross-sectional cohort study

Primary study design

Observational

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Preterm delivery, threatening preterm labor

Interventions

Electrohysterogram: power spectral density peak frequency, conduction velocity

Current diagnostics: digital examination, cervical length, fetal fibronectin

All diagnostics will be performed at admission: a 4-channel electrohysterogram recording of 30 minutes using a fixed electrode configuration, a digital examination, transvaginal cervical length measurement and fetal fibronectin testing. Electrohysterogram information will be blinded to the attending physician, analysis will be done afterwards. The peak frequency of the power density spectrum (by Fourier Transform) will be identified. For conduction velocity the corresponding action potentials will be visually identified in the different channels. The velocity (vector) will be calculated from these intervals from which the speed (scalar) will be used as conduction velocity.

Two cohorts will formed based on delivery within or after 7 days after measurement.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Sensitivity, specificity, positive and negative predicting value for predicting preterm delivery within 7 days from measurement

Key secondary outcome(s))

Area Under the Curve of the Receiver Operating Characteristic (ROC) curve for predicting preterm delivery within 7 days from measurement

Completion date

01/06/2014

Eligibility

Key inclusion criteria

- 1. Patients admitted for threatened preterm labor
- 2. Gestational age between 23+5 and 34+0 weeks
- 3. Clinically evaluated symptoms of preterm labor: at least 6 contractions in 60 minutes based on the (cardio) tocogram and/or maternal perception

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Key exclusion criteria

Patients in active labor: cervical dilatation ≥3cm

Date of first enrolment

01/02/2012

Date of final enrolment

01/06/2014

Locations

Countries of recruitment

Netherlands

Study participating centre Tannhauserdreef 272

Utrecht Netherlands 3561HR

Sponsor information

Organisation

Maxima Medical Center (Netherlands)

ROR

https://ror.org/02x6rcb77

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Maxima Medical Center (Netherlands)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summaryNot provided at time of registration

Study outputs

Output type	Details	Date created Date added	Peer reviewed?	Patient-facing?
Protocol article	protocol	05/06/2014	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	No	Yes