

External peripheral neuromodulation for the relief of chronic pain

Submission date 05/08/2009	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 18/02/2010	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 18/07/2016	Condition category Nervous System Diseases	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Teodor Goroszeniuk

Contact details
Pain Management & Neuromodulation Centre
St Thomas' Hospital
Westminster Bridge Road
London
United Kingdom
SE1 7EH

Additional identifiers

Protocol serial number
RJ1 09/N084

Study information

Scientific Title
A randomised controlled feasibility trial of external peripheral neuromodulation in the treatment of intractable localised chronic neuropathic pain

Study objectives

The study hypothesis is that, when given over three treatment sessions at weekly intervals, external peripheral neuromodulation will reduce pain severity at 7 days after the final treatment in patients with intractable localised chronic pain with predominantly neuropathic features.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Guy's Research Ethics Committee, 22/04/2009, ref: 09-H0804-46

Study design

Single centre double-blind randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Chronic neuropathic pain

Interventions

Three 5-minute treatment sessions of external peripheral neuromodulation at 2 Hz (or sham stimulation) at weekly intervals. Follow-up to 7 days after the last session, i.e. 21 days.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Visual Analogue Score (VAS) of pain severity one week following third (final) neuromodulation session.

Key secondary outcome(s)

1. VAS of pain severity immediately after treatment and at 7 day intervals to end of study (i.e., days 7, 14 and 21)
2. VAS of pain severity at shorter intervals of less than a week
3. Numerical rating scores (NRS) of:
 - 3.1. Sleep quality
 - 3.2. Functional impact
 - 3.3. Emotional impactMeasured at 7 days following third (final) neuromodulation session.
4. Numerical rating scores on Pain Self-Efficacy Questionnaire (PSEQ) at 7 days following third (final) neuromodulation session
5. Changes in medication usage
6. Change in size of painful area on pain/paraesthesiae map

Completion date

16/08/2010

Eligibility

Key inclusion criteria

1. Patient age is 18 years or older, either sex
2. Patient has chronic neuropathic (or predominantly neuropathic) intractable pain
3. Patient has an area of pain with a typical dermatomal distribution that can be expected to be covered with a single episode of local stimulation
5. In the opinion of the Investigator and the patient's consultant, all standard medical options have been tried without sufficient improvement in pain control (including opioids, nerve blocks, etc.)
6. Patient has a Visual Analogue Scale (VAS) pain score of 5 cm (or greater) on a 10 cm line

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Patient has a history of substance abuse or substance dependency in the past 6 months prior to baseline data collection
2. Patient is currently participating in another clinical study
3. Patient lacks capacity for informed consent to the trial in the view of person taking consent and/or investigators
4. Pregnancy - if the painful area lies within 50 cm of the gravid uterus
5. Patient has difficulties in adequate understanding of English for consent, clinical review and self-completion questionnaires
6. Patient does not permit notification to General Practitioner of enrolment in the study
7. Patient has previous experience of peripheral neuromodulation. Note that prior transcutaneous electrical nerve stimulation (TENS) experience is permitted.

Date of first enrolment

17/08/2009

Date of final enrolment

16/08/2010

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Pain Management & Neuromodulation Centre

London

United Kingdom

SE1 7EH

Sponsor information

Organisation

Guy's and St Thomas' NHS Foundation Trust (UK)

ROR

<https://ror.org/00j161312>

Funder(s)

Funder type

Government

Funder Name

Guy's and St Thomas' NHS Foundation Trust (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration