

Effectiveness of anti-inflammatory treatment versus antibiotic therapy and placebo in patients with acute bronchitis and purulent expectoration

Submission date 02/03/2010	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 04/03/2011	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 28/10/2013	Condition category Respiratory	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Ana Moragas

Contact details
Av. Horta de Santa Maria, 4, 3r 4a
Cambrils
Spain
43850
+34 (0)60 007 21 70
amoragasm@meditex.es

Additional identifiers

Clinical Trials Information System (CTIS)
2007-006727-12

Protocol serial number
E07/90333

Study information

Scientific Title

Effectiveness of anti-inflammatory treatment versus antibiotic therapy and placebo in patients with acute bronchitis and purulent sputum: a double-blind randomised controlled clinical trial

Acronym

BAAP Study

Study objectives

The number of days of frequent cough (defined by the symptom diary with a score of 1 or more) among patients taking the anti-inflammatory drug will be fewer than among patients assigned to the other two arms.

Ethics approval required

Old ethics approval format

Ethics approval(s)

1. The Ethical Committee of Investigation in Primary Care (Fundació d'Investigació en Atenció Primària) approved on the 2nd of December 2007 (ref: P07/15)
2. Spanish Agency of Medicines and Health Products (Agencia Española de Medicamentos y Productos Sanitarios [AEMPS]) and the EudraCT approved (ref: 2007-006727-12)

Study design

Double blind randomised controlled clinical trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Acute bronchitis and purulent sputum

Interventions

The patients will be randomised with simple random numbers to one of the three following treatment arms:

1. Ibuprofen 600 mg/8 hours, during 10 days, taken after meals
2. Amoxicillin plus clavulanic acid 500-125 mg/8 hours, during 10 days, taken after meals
3. Placebo, one tablet every 8 hours, during 10 days, taken after meals

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Number of days with frequent cough

Key secondary outcome(s)

Secondary result variables considered will be the efficacy of the treatment, the time of symptom resolution, the reduction in the total daily score of the symptom diary and the secondary effects and adverse reactions which may present with the three treatment arms. With regard to efficacy, three outcomes will be distinguished: cure, defined as the disappearance of the acute signs and symptoms related to the infection (complete return to the previous situation of stability), improvement, defined as the non complete resolution of the symptoms and failure, with an insufficient reduction in the signs and symptoms of infection.

Completion date

31/03/2011

Eligibility

Key inclusion criteria

Patients from 18 to 70 years of age without associated respiratory comorbidity or immunosuppression will be recruited from seven healthcare centers in Catalonia.

1. Presenting respiratory infection of at least one week of evolution
2. Cough as the predominant symptom
3. Presence of purulent expectoration of at least one week of duration
4. At least one other respiratory tract symptom such as dyspnoea, wheezing, chest discomfort or pain, with no alternative explanation such as pneumonic condensation

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

70 years

Sex

All

Key exclusion criteria

1. Patients less than 18 and older than 70 years of age
2. Presence of radiological signs of pneumonia
3. Signs of severe infection
 - 3.1. confusion
 - 3.2. tachypnoea > 25 respirations per minute of tachycardia > 120 beats per minute
4. History of digestive haemorrhage or intolerance to anti-inflammatory treatment
5. Hypersensitivity to β -lactam or intolerance to clavulanic acid or lactose
6. Pregnancy, lactation and women of fertile age not using contraceptive measures

7. Antibiotic, anti-inflammatory or corticoid use in the previous two weeks
8. Associated comorbidity
 - 8.1. bronchial asthma
 - 8.2. chronic obstructive pulmonary disease
 - 8.3. moderate-severe heart failure
 - 8.4. dementia
 - 8.5. stroke
 - 8.6. immunosuppression or the use of immunosuppressive drugs
9. Emergency situation
10. Institutionalisation in a residence
11. And/or subjects unable to personally provide informed consent

Date of first enrolment

01/04/2010

Date of final enrolment

31/03/2011

Locations

Countries of recruitment

Spain

Study participating centre

Av. Horta de Santa Maria, 4, 3r 4a
Cambrils
Spain
43850

Sponsor information

Organisation

Primary Care Research Institute Jordi Gol i Gurina (Institut d'Investigació en Atenció Primària Jordi Gol i Gurina) (Spain)

ROR

<https://ror.org/0370bpp07>

Funder(s)

Funder type

Government

Funder Name

Health Research Fund of the Ministry of Health and Consumption (Fondo de Investigación Sanitaria de Ministerio de Sanidad y Consumo) (Spain) (EC07/90333)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	04/10/2013		Yes	No
Protocol article	protocol	21/06/2011		Yes	No