

Effect of prophylactic antibiotics on post flexible cystoscopy infection rate

Submission date 28/09/2007	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 28/09/2007	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 15/03/2016	Condition category Urological and Genital Diseases	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Conal S L Quah

Contact details
115B Loftus House
Marton Road
Middlesbrough
United Kingdom
TS4 3TQ
+44 (0)7738005994
conalquah@yahoo.com

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
N0227189911

Study information

Scientific Title

Effect of prophylactic antibiotics on post flexible cystoscopy infection rate

Study objectives

Are patients less susceptible to urinary tract infection post cystoscopy following administration of single-dose antibiotic prophylaxis?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Newcastle and North Tyneside 1 Ethics Committee (UK), 06/03/2007, ref: REC 07/Q0905/3

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Urological and Genital Diseases: Urinary tract infection (UTI)

Interventions

Null hypothesis: The use of a single dose oral antibiotic prophylaxis will not influence the rate of urinary tract infection post flexible cystoscopy.

Purpose: The aim of this study is to determine if patients are less susceptible to urinary tract infection post cystoscopy following administration of single-dose antibiotic prophylaxis.

Design: This study is a prospective randomised clinical trial. Every patient listed for flexible cystoscopy will be randomised appropriately, dividing them into 2 groups. One group will receive antibiotics and one group will not.

Methodology: Patients listed for flexible cystoscopy from Nov 06-March 07 will be recruited for this study. Clinicians and staff as well as the patients' GPs will be informed about this research

study. Those recruited will be given a patient information sheet as well as informed consent obtained. Pre and post procedure urine samples will be obtained to objectively evaluate the presence of infection. Cystoscopies will be carried out in the surgical day unit (James Cook University Hospital) by either a specialist trainee or a consultant. External genitalia will be cleaned with topical chlorhexidine and plain lignocaine jelly 2% will be used for local anaesthetic. The cystoscope will be disinfected with Transept. 5-7 days post procedure, they would be required to return the urine samples to their respective GP. The results will be analyzed to determine the rate of positive cultures in both groups.

Predictable risks and inconvenience weighted against the anticipated benefits for participants include a small possibility of a side effect from the antibiotic. However, the antibiotic administration might prevent an infection post procedure hence the interest in this research study. The results of this study will then help develop a local protocol which will standardize the current practice. Patients may experience minor inconvenience as they have to hand in the urine sample themselves to the GP 5-7 days following the procedure.

Interpreting and analysing findings, preparing the final report will be done for around a month after the proposed recruitment date and a paper on the subject will be written and submitted to a national/international journal for peer review.

Intervention Type

Other

Phase

Not Specified

Primary outcome measure

The primary outcome measure is to identify the incidence of urinary tract infection post flexible cystoscopy following prophylaxis antibiotic administration.

Secondary outcome measures

No secondary outcome measures

Overall study start date

15/11/2006

Completion date

30/09/2009

Eligibility

Key inclusion criteria

1. Male and female patients of all ages
2. Patients undergoing check or flexible cystoscopy

Participant type(s)

Patient

Age group

Not Specified

Sex

Both

Target number of participants

328, 80 patients recruited as of August 2008.

Key exclusion criteria

1. High susceptibility patients who would have definitely received antibiotics - diabetics, immuno compromised patients, recurrent UTIs
2. Patients who underwent biopsy / procedure ie stent removal etc during cystoscopy
3. Patients with indwelling catheters
4. Patients who are allergic to Penicillin

Date of first enrolment

15/11/2006

Date of final enrolment

30/09/2009

Locations**Countries of recruitment**

England

United Kingdom

Study participating centre

115B Loftus House

Middlesbrough

United Kingdom

TS4 3TQ

Sponsor information**Organisation**

Record Provided by the NHSTCT Register - 2007 Update - Department of Health

Sponsor details

The Department of Health, Richmond House, 79 Whitehall

London

United Kingdom

SW1A 2NL

+44 (0)20 7307 2622

dhmail@doh.gsi.org.uk

Sponsor type

Government

Website

<http://www.dh.gov.uk/Home/fs/en>

Funder(s)**Funder type**

Government

Funder Name

South Tees Hospitals NHS Trust (UK)

Results and Publications**Publication and dissemination plan**

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration