A pilot randomised controlled study of oxygen delivery via Vapotherm® in infants with severe acute bronchiolitis

Submission date Recruitment status Prospectively registered 28/09/2007 No longer recruiting [] Protocol Statistical analysis plan Registration date Overall study status 28/09/2007 Completed [X] Results [] Individual participant data Condition category Last Edited 08/05/2013 Respiratory

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Tom Hilliard

Contact details

C/O Research and Effectiveness Department Level 1, Old Building Bristol Royal Infirmary Marborough Street Bristol United Kingdom BS2 8HW +44 0117 928 3473

Additional identifiers

Protocol serial number

N0264191693

Study information

Scientific Title

Study objectives

We wish to study the safety and efficacy of the Vapotherm oxygen delivery device in infants with severe acute bronchiolitis in an initial pilot trial.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised open pilot study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Respiratory: Severe acute bronchiolitis

Interventions

Randomised open pilot study of oxygen delivery via Vapotherm compared with conventional therapy (optimum oxygen delivery via a headbox and appropriate intravenous fluids. Parents have up to 6 hours to decide whether to take part.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Vapotherm®

Primary outcome(s)

Oxygen saturation (SpO2) 8 hours post randomisation to either Vapotherm or continuing conventional therapy.

Key secondary outcome(s))

- 1. SpO2, heart rate, respiratory rate, FiO2 (oxygen concentration), blood pressure and combined bronchiolitis
- 2. Severity score at 4 hours, 8 hours, 12 hours, 24 hours, 36 hours and 48 hours after point of randomisation
- 3. Length of stay in hospital; length of time receiving oxygen therapy after randomisation; length of time before enteral feeds re-started after randomisation
- 4. Proportion of infants in each group that require further respiratory support, with either CPAP or intubation and mechanical ventilation

Completion date

15/05/2007

Eligibility

Key inclusion criteria

Subjects will be infants less than 12 months of age admitted to hospital with a clinical diagnosis of

- 1. Bronchiolitis (clinical picture of cough, tachypnoea, chest retraction and crackles on auscultation) and
- 2. Features of severe disease:
- 2.1 Head–box oxygen requirement greater than 50% to maintain SpO2 at least 92%
- 2.2 Feeds have been discontinued, and intravenous fluid started
- 2.3 Moderate to severe tachypnoea and increased work of breathing
- 2.4 Triggering of the Paediatric Early Warning Tool

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Neonate

Sex

Not Specified

Key exclusion criteria

- 1. Congenital cyanotic heart disease
- 2. Repeated severe apnoeas
- 3. Requirement for resuscitation with bag-mask ventilation
- 4. Severe hypercapnia (increased carbon dioxide) with acidosis on blood gas analysis of pH less than 7.2
- 5. Parental refusal or inability to give consent

Date of first enrolment

15/01/2007

Date of final enrolment

15/05/2007

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
C/O Research and Effectiveness Department
Bristol
United Kingdom
BS2 8HW

Sponsor information

Organisation

Record Provided by the NHSTCT Register - 2007 Update - Department of Health

Funder(s)

Funder type

Government

Funder Name

United Bristol Healthcare NHS Trust

Funder Name

NHS R&D Support Funding

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/02/2012		Yes	No