Radon Balneology and Physical Activity for Osteoporosis Prevention

Submission date	Recruitment status No longer recruiting	Prospectively registered		
14/01/2014		∐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
10/02/2014	Completed	[X] Results		
Last Edited 02/11/2015	Condition category Musculoskeletal Diseases	Individual participant data		

Plain English summary of protocol

Background and study aims

Osteoporosis is a common condition characterized by decreased bone mass leading to increased fracture risk. The economic burden was estimated at 799 million in Austria for 2010 and could increase by 28 % to 1,025 million in 2025. Mountain hiking is a popular form of exercise in the alpine region and a cost effective treatment. Numerous studies have confirmed the effects of exercise programs on bone metabolism and inflammatory markers in postmenopausal women (a high risk group for osteoporosis). Studies have also shown that radon balneotherapy can help reduce pain for patients with degenerative spinal disease, rheumatoid arthritis and ankylosing spondylitis. The aim of the study is to assess whether a combination of exercise and radon balneotherapy works better than exercise only.

Who can participate?

Females and males aged between 50 and 65 who are fit enough to complete a hiking program and can complete German questionnaires.

What does the study involve?

Participants are randomly allocated to one of two groups: exercise and radon balneotherapy or exercise and non radon (dummy) balneotherapy.

Participants have a week long treatment, then 6 weeks off, then another 3 day treatment, and a follow up period of 240 days.

The first treatment includes five 3-4 hours guided GPS monitored mountain hiking tours with a minimum of 300 meters altitude difference per day, and five radon or placebo baths.

The second treatment includes two hiking tours plus two radon or placebo baths.

Participants have blood tests at the start and end of each treatment weeks and 6 months after the last treatment. They also fill in questionnaires each time.

What are the possible benefits and risks of participating?

Benefits are 9 days of free stay, meals and guided hiking tours in an Austrian resort area. The risks are comparable to that of a standard touristic stay in the alpine regions including spa visits with low dose radon hyperthermia.

Where is the study run from?
Gastein resort area in Salzburg, Austria

When is the study starting and how long is it expected to run for? The study started in Autumn 2009 and is expected to run until Autumn 2015

Who is funding the study?
Gastein Research Institute, Paracelsus Medical University Salzburg (Austria).

Who is the main contact? Dr. Arnulf Hartl arnulf.hartl@pmu.ac.at

Contact information

Type(s)

Scientific

Contact name

Dr Arnulf Hartl

Contact details

Institute of Physiology and Pathophysiology Paracelsus Medical University Salzburg Salzburg Austria 5020

Additional identifiers

Protocol serial number

E1216/8-2010

Study information

Scientific Title

Gastein Radon Balneology and Physical Activity for Osteoporosis Prevention: a randomized, placebo controlled intervention study

Study objectives

Does a combined exercise, hyperthermia and radon balneology treatment have any impact on central biomarkers for bone remodeling in a study population at the risk age for developing osteoporosis?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Salzburg Ethics Committee (Ethikkommission für das Bundesland Salzburg], 26 May 2010, ref: E1216/8-2010

Study design

Double blind randomized controlled clinical trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Osteporosis prevention

Interventions

Exercise +/- Radon balneotherapy

Study subjects are allocated by random into a radon group and placebo group Randomization will be computed in blocks of four with an equal treatment allocation ratio. Radon- or placebo thermal-water bath filling is blinded with a personalized digital ID card that provides tub filling with either radon thermal water or radon degassed thermal water (placebo) in an automatic process. In this double blind setting neither the participants nor the bath attendant have knowledge of the composition of the water.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Plasma levels of OPG for bone formation, RANKL for bone reabsorption and their ratio (OPG /RANKL) as a measure for the balance of bone metabolism

Forearm venous blood is collected from all participants at baseline (day 0, timepoint 1; T1), end of first treatment week (day 6, timepoint 2; T2), start and end of the brushup (day 60 and 63, timepoint 3; T3 and timepoint 4; T4) and 6 months after the last treatment (followup, day 240, timepoint 5; T5).

Questionnaires assessing somatic complaints (Beschwerdenliste, Zerssen, D. v., 1976. Die Beschwerden- Liste - Manual. Weinheim: Beltz Test GmbH) and Quality of Life (Quality of life questionnaire of the European foundation for osteoporosis 41; QUALEFFO 41) are handed out for completion on T 1, T2, T3, T4, T5 and days T1, T3, T5, respectively.

Key secondary outcome(s))

- 1. Plasma levels of Osteocalcin (OC), Osteopontin (OPN), Leptin, Parathyroid Hormone (PTH) and Adrenocorticotropic Hormone (ACTH)
- 2. Somatic complaints and self-reported health. Questionnaires assessing somatic complaints (Beschwerdenliste, Zerssen, D. v., 1976. Die Beschwerden- Liste Manual. Weinheim: Beltz Test GmbH) and Quality of Life (Quality of life questionnaire of the European foundation for osteoporosis 41; QUALEFFO 41) are handed out for completion on T 1, T2, T3, T4, T5 and days T1, T3, T5, respectively.

Completion date

01/09/2014

Eligibility

Key inclusion criteria

- 1. Eligible participants are working females and males aged between 50 and 65.
- 2. Physical ability to meet the demands of the exercise program (i.e. at least 300 m of altitude difference per day) and sufficient knowledge of German to complete the questionnaires.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Senior

Sex

All

Key exclusion criteria

- 1. Acute or chronic disturbances of the immune system
- 2. Hyperthyreosis
- 3. Cardiac arrhythmia
- 4. Renal insufficiency
- 5. Severe cardiovascular diseases
- 6. Acute infections or fever
- 7. Iritis
- 8. An acute attack of polyarthritis
- 9. Participants should not have used hormone replacement therapy or any other therapy affecting the bone metabolism during the last 12 months before enrollment.

Date of first enrolment

01/09/2010

Date of final enrolment

01/09/2014

Locations

Countries of recruitment

Austria

Study participating centre Institute of Physiology and PathophysiologySalzburg

Sponsor information

Organisation

Paracelsus Medical University Salzburg (Austria)

ROR

https://ror.org/03z3mg085

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Forschungsfond des Forschungsinstituts Gastein (FOI-FFF) of the Gastein Research Institute, Paracelsus Medical University Salzburg (Austria)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/03/2015		Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes