# Comparing two different loading times for dental implants

Submission date	Recruitment status No longer recruiting	Prospectively registered	
12/03/2019		☐ Protocol	
Registration date	Overall study status	Statistical analysis plan	
08/04/2019	Completed	[X] Results	
<b>Last Edited</b> 26/03/2019	Condition category Oral Health	[] Individual participant data	

## Plain English summary of protocol

Background and study aims

In the past, dental implant screws were placed in the jaw bone without teeth being added for 3-8 months to allow the bone to heal around the screw. More recently, immediate- and early-loaded implants are commonly used. Immediate-loaded implants have teeth attached to the screws within 72 hours from implant placement. Early-loaded implants are loaded approximately 6 weeks after implant placement. The aim of this 10-year study is to compare implant failures, complications, patient satisfaction and bone level of immediate and early loading of dental implants in people with some tooth loss.

#### Who can participate?

Adults with some remaining natural teeth requiring dental implants.

#### What does the study involve?

The participants were randomly allocated to the immediate or early loading group. Both groups had the dental implants inserted as usual. In the immediate loading group the implants were loaded with teeth within 72 hours from implant placement. In the early loading group, the implants were loaded with teeth approximately 6 weeks after implant placement. The implants were monitored for 10 years to measure the implant failure rate, implant complications, patient satisfaction with the function and appearance of the implants and bone loss measured by X-ray.

What are the possible benefits and risks of participating?

The potential benefit was that the immediate-loading group would not have to wait for 6 weeks to have teeth loaded onto their implants. The implant procedure has some risks including bleeding during and after implant placement, pain and swelling around the implant site, fracture of the implant or fracture of the artificial tooth.

Where is the study run from? Clinica Merli, Rimini (Italy)

When is the study starting and how long is it expected to run for? February 2005 to November 2018

Who is funding the study? This study was investigator funded.

Who is the main contact?
Mauro Merli, mauromerli@gmail.com

# **Contact information**

## Type(s)

Public

#### Contact name

Dr Mauro Merli

#### Contact details

Clinica Merli Viale Settembrini 17/O Rimini Italy 47923 +39-0541-52025 mauromerli@gmail.com

## Type(s)

Scientific

#### Contact name

Dr Michele Nieri

#### **ORCID ID**

https://orcid.org/0000-0001-8770-4622

#### Contact details

Viale largo Brambilla 3 Firenze Italy 50134 +39055416434 michelenieri@gmail.com

# Additional identifiers

**Protocol serial number** RCT7

# Study information

Scientific Title

Immediate versus early non-occlusal loading of dental implants placed flapless in partially edentulous patients: A 10-year randomized clinical trial

#### **Acronym**

Immediate versus early loading

#### Study objectives

To compare immediate versus early non-occlusal loading of dental implants placed flapless

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

This study started in early 2005 and at the time in Italy there was no a clear regulation for studies on a medical device.

# Study design

Single-centre assessor-blind parallel randomized clinical trial

## Primary study design

Interventional

## Study type(s)

**Treatment** 

# Health condition(s) or problem(s) studied

Dental implantation in partially edentulous patients

#### **Interventions**

Patients were randomized to receive implants for fixed partial dentures. The test group was represented by immediate non-occlusal implant loading, whereas the control group was represented by early non-occlusal implant loading. An investigator, not involved in the selection and treatment of the patients, randomly assigned participants following simple randomization procedures (computerized random numbers) to one of two treatment groups. The randomized codes were enclosed in sequentially numbered, identical, opaque, sealed envelopes. Envelopes were opened sequentially only after the implants to be included in the trial were inserted, and therefore treatment allocation was concealed to the investigator in charge of enrolling and treating the patients included in the trial.

All provisional restorations of the immediate loaded group were placed within 72 h of implant placement. The occlusal surface of the provisional restoration was ground to avoid any occlusal contact with the opposite dentition in static and dynamic analyses.

For patients in the early loading group, impressions were taken approximately 6 weeks after implant placement. They received non-occlusally provisional fixed restorations, identical to those of the immediately loaded group.

## Intervention Type

Procedure/Surgery

## Primary outcome(s)

Implant failure (the presence of any mobility of the implant and/or any situation dictating implant removal) after 10 years

#### Key secondary outcome(s))

- 1. Biological or prosthetic complications defined as unexpected deviations from the normal treatment outcome: examples of biological complications are hemorrhaging during and after implant placement and/or peri-implantitis. Prosthetic complications included fracture of the implant or fracture of the prosthesis.
- 2. Patient's aesthetic and functional satisfaction were assessed by asking the patient for satisfaction on a scale from 0 to 10 where 0 meant completely dissatisfied and 10 meant completely satisfied
- 3. Peri-implant marginal bone level assessed using periapical intra-oral radiographs taken with the parallel technique at 10 years of follow-up

#### Completion date

05/11/2018

# Eligibility

#### Key inclusion criteria

- 1. Aged 18 years or over
- 2. Partially dentate and requiring dental implants
- 3. Implant site allows for the placement of at least one 9.5 mm long implant
- 4. Bone thickness at the implant site at least 5.5 mm
- 5. For patients with multiple areas to be restored, the operator was free at the screening visit to select one area to be included in the trial. In this area, multiple neighboring implants could be placed.

#### Participant type(s)

Patient

# Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Sex

Αll

#### Total final enrolment

60

#### Key exclusion criteria

- 1. General contraindications to implant surgery
- 2. Patients irradiated in the head and neck area within a year prior to surgery

- 3. Patients with poor oral hygiene (full-mouth plaque score ≥30) and lack of motivation
- 4. Uncontrolled diabetes (diabetes that is not being treated at all, or is not being adequately treated)
- 5. Pregnant or lactating
- 6. Substance abusers
- 7. Psychiatric problems
- 8. Lack of opposing occluding dentition in the area intended for implant placement
- 9. Severe bruxism or clenching
- 10. Active infection or severe inflammation in the area intended for implant placement
- 11. Presence of ≤4 mm of keratinized mucosa and/or the need for bone augmentation procedures, with the exception of post-extractive sites treated with Bio-Oss® granules

# Date of first enrolment

06/07/2005

Date of final enrolment 25/07/2007

# Locations

#### Countries of recruitment

Italy

# Study participating centre

Clinica Merli

Viale Settembrini 17/O Rimini Italy 47923

# Sponsor information

# Organisation

Clinica Merli

# Funder(s)

## Funder type

Other

#### Funder Name

Investigator funded

# **Results and Publications**

# Individual participant data (IPD) sharing plan

The datasets generated during the current study will be available upon request from Mauro Merli (mauromerli@gmail.com) after the publication of the trial at 10-year follow-up. Fields related to patient demographics will be deleted.

# IPD sharing plan summary

Available on request

## **Study outputs**

Output type	Details	Date created Date added	Peer reviewed?	Patient-facing?
Results article	1-year follow-up results	01/09/2008	Yes	No
Results article	3-year follow-up results	01/02/2012	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	No	Yes