# CO2 gap and weaning outcome from mechanical ventilation

Submission date	Recruitment status	Prospectively registered
27/08/2019	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
06/09/2019	Completed	Results
Last Edited	Condition category	Individual participant data
10/09/2019	Respiratory	[] Record updated in last year

## Plain English summary of protocol

Background and study aims

After recovery from acute events that have led to intubation and ventilation, weaning from mechanical ventilation is the essential and usual component in the management of critically ill patients. Weaning failure, which occurs in approximately 25 to 42% of patients meeting the readiness criteria, is associated with prolonged mechanical ventilation, increased morbidity, and mortality. Thus, early identification of patients that are going to fail the liberation from mechanical ventilation is of paramount importance to improve the outcome of these critically ill patients.

Discontinuation of ventilatory support can be very challenging for clinicians mainly because the pathophysiology of weaning failure is complex and not fully understood. This study aims to investigate the central venous-to-arterial difference in carbon dioxide in patients who fail a weaning trial.

Who can participate?

Patients aged 18 years or older under mechanical ventilation for at least 48 hours who fulfil the inclusion criteria

What does the study involve?

Treatment will be continued as normal. Data will be collected on the success or failure of weaning from mechanical ventilation

What are the possible benefits and risks of participating?

The possible benefits are: It is well known that critically ill patients under mechanical ventilation who fail the extubation are at high risk of getting aspiration pneumonia, increased morbidity, and mortality. Weaning failure still occurs in as high as 40% of patients who already met the standard criteria for weaning as established by different societies of respiratory and critical care medicine. Therefore, looking for new indices that help to predict the weaning outcome accurately is of utmost importance to reduce the incidence of weaning failure, which is associated with high morbidity and mortality. We found that the combination of changes in CO2gap (difference between venous-to arterial PCO2 difference) and ScvO2 (central venous oxygen saturation) during a spontaneous breathing reliably predict the weaning outcome and can assist the Physician in his/her decision to extubate the patient in minimalization the risk of

failure and its associated elevated morbidity and mortality.

Theoretically, there is no risk in participating in this study as the included patients have already venous central and arterial catheters inserted before being included in the study. Also, the decision of extubation was left at the discretion of the attending Physician who was following the standard international criteria of readiness for extubation.

Where is the study run from?

- 1. Centre Hospitalier de Lens
- 2. Centre Hospitalier d'Arras
- 3. Centre Hospitalier de Cambrai

When is the study starting and how long is it expected to run for? December 2016 to February 2018

Who is funding the study? Investigator initiated and funded

Who is the main contact? Dr Jihad Mallat mallatjihad@gmail.com

## Contact information

## Type(s)

Scientific

#### Contact name

Dr Jihad Mallat

#### **ORCID ID**

https://orcid.org/0000-0001-6372-7360

#### Contact details

99 Route de la bassee Lens France 62300 +33321691088 jmallat@ch-lens.fr

## Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

161205

# Study information

#### Scientific Title

Combination of central venous-to-arterial PCO2 difference and central venous oxygen saturation accurately predict weaning outcome from mechanical ventilation

## **Study objectives**

PCO2 gap is able to accurately predict the weaning outcome from mechanical ventilation in critically ill patients

### Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approved 01/12/2016, Centre Hospitalier de Lens Comité d'éthique (99 route de la Bassée, 62307, Lens Cedex, France; +33 321691234; bdelepine@ch-lens.fr), ref: 160402

### Study design

Prospective multi-center study

#### Primary study design

Observational

### Study type(s)

Diagnostic

## Health condition(s) or problem(s) studied

Mechanical ventilation

#### Interventions

A patient was included when readiness to wean criteria were met (see inclusion criteria)

A spontaneous breathing trial (SBT) was then performed, in semi-recumbent patients, on a T-piece with supplementary humidified oxygen to achieve arterial oxygen saturation >90% as measured by pulse oximetry. The duration of the SBT was 60 minutes. Criteria for poor SBT tolerance were:

- 1. Diaphoresis
- 2. Use of accessory respiratory muscles
- 3. RR > 35/min
- 4. Oxygen saturation by pulse oximetry <90% with FiO2 ≥50%
- 5. HR >140/min or greater than a 20% increase from baseline
- 6. Systolic blood pressure >180 mmHg or <90 mmHg
- 7. Development of cardiac arrhythmias; and/or low level of consciousness (Glasgow Coma Scale score <13).

The decision to stop SBT was made by physicians. Patients who did not tolerate the SBT were reconnected to a ventilator.

Patients who successfully completed the SBT were extubated and followed up for 72 hours. Weaning failure was diagnosed if SBT was aborted because of clinical intolerance or if the

patient was extubated but required mechanical ventilation (invasive or non-invasive), because of respiratory failure, within the following 48 hours. Respiratory failure after extubation was defined as the development of at least one of the following:

- 1. Respiratory acidosis with pH <7.32 and arterial CO2 pressure (PaCO2) >45 mmHg
- 2. Arterial oxygen saturation <90% with FiO2 >0.5
- 3. RR>35/min
- 4. Clinical signs of respiratory fatigue

The management of post-extubation respiratory failure was not protocolized and left to the physician's discretion

#### Intervention Type

Other

## Primary outcome(s)

Weaning failure rate from mechanical ventilation defined as the development of at least one of the following:

- 1. Respiratory acidosis with pH <7.32 and arterial CO2 pressure (PaCO2) >45 mmHg
- 2. Arterial oxygen saturation <90% with FiO2 >0.5
- 3. RR>35/min
- 4. Clinical signs of respiratory fatigue

## Key secondary outcome(s))

None

### Completion date

20/02/2018

# Eligibility

## Key inclusion criteria

- 1. Aged 18 years or older
- 2. Under mechanical ventilation for at least 48 hours
- 3. Satisfied the weaning criteria. The readiness to wean criteria were:
- 3.1 The resolution or improvement of the underlying cause of respiratory failure for which the patient was intubated
- 3.2 Hemodynamic stability, defined as heart rate (HR) < 140/min and systolic blood pressure between 90 and 160 mmHg with no or minimal doses of vasopressors
- 3.3 Stable respiratory status, defined as oxygen saturation >90% with fraction of inspired oxygen (FiO2) ≤0.4 and positive end expiratory-pressure (PEEP) ≤8 cmH2O, respiratory rate (RR) ≤35/min, spontaneous tidal volume (Vt) >5 mL/kg, ratio of RR/Vt <105/min per liter, and no significant respiratory acidosis
- 3.4 Adequate mental status (Glasgow Coma Scale score >13)
- 3.5 Adequate cough.
- 4. Patients also had to have an arterial catheter and a central line with the tip confirmed by x-ray to be in the superior vena cava near at the entrance or in the right atrium.

## Participant type(s)

**Patient** 

## Healthy volunteers allowed

## Age group

Adult

## Lower age limit

18 years

#### Sex

All

## Key exclusion criteria

- 1. Tracheotomized patients
- 2. Do-not-reintubate orders
- 3. Pregnancy
- 4. No informed consent
- 5. Poor cardiac echogenicity

#### Date of first enrolment

28/12/2016

#### Date of final enrolment

19/02/2018

## Locations

#### Countries of recruitment

France

## Study participating centre Centre Hospitalier de Lens

99 Route de la bassee

Lens

France

62300

## Study participating centre Centre Hospitalier d'Arras

Boulevard Georges Besnier

Arras

France

62022

## Study participating centre

### Centre Hospitalier de Cambrai

Avenue de Paris Cambrai France 59400

# Sponsor information

## Organisation

Centre Hospitalier de Lens

#### **ROR**

https://ror.org/01fbc7819

# Funder(s)

## Funder type

Other

#### **Funder Name**

Investigator initiated and funded

## **Results and Publications**

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request

## IPD sharing plan summary

Available on request

### **Study outputs**

Output type **Details** Date created Date added Peer reviewed? Patient-facing? Participant information sheet 11/11/2025 11/11/2025 No

Participant information sheet

Yes