Aerobic versus anaerobic exercise and oral nutritional supplementation in nutritional status and physical function of adults hemodialysis patients

Recruitment status	Prospectively registered
No longer recruiting	☐ Protocol
Overall study status	Statistical analysis plan
Completed	[X] Results
Condition category	Individual participant data
	No longer recruiting Overall study status Completed

Plain English summary of protocol

Background and study aims

Hemodialysis is a process of purifying the blood of a person whose kidneys are not working normally. A condition called protein-energy wasting, where the body has decreased stores of protein and fat, is common in patients undergoing hemodialysis. Studies have looked at the benefits of oral nutritional supplements, resistance exercise (e.g., weight training) and aerobic exercise (e.g., cycling) on nutrition and physical function, but it is not known what kind of treatment is best for patients. The aim of this study is to compare the effect of oral nutritional supplementation plus resistance exercise, oral nutritional supplementation plus aerobic exercise, and oral nutritional supplementation alone.

Who can participate?

Patients aged 18 to 45 undergoing hemodialysis

What does the study involve?

Participants are randomly allocated to receive one of three treatments two times per week for 12 weeks: either oral nutritional supplement plus resistance exercise, oral nutritional supplement plus aerobic exercise, or oral nutritional supplement alone. At the end of the study participants' body measurements, blood test results, physical function and quality of life are assessed.

What are the possible benefits and risks of participating?

It has been shown that exercise and oral nutritional supplements are safe and beneficial. Low blood pressure and symptoms such as diarrhoea and nausea have been associated with oral nutritional supplements, but there is a lack of evidence supporting this.

Where is the study run from? Hospital General de México, Dr Eduardo Liceaga (Mexico) When is the study starting and how long is it expected to run for? January 2016 to September 2017

Who is funding the study? Hospital General de México, Dr Eduardo Liceaga (Mexico)

Who is the main contact?

- 1. Dr Lucía Monserrat Pérez
- 2. Miss Geovana Martin

Contact information

Type(s)

Scientific

Contact name

Dr Lucía Monserrat Pérez

ORCID ID

https://orcid.org/0000-0002-9281-9202

Contact details

Dr. Balmis 148 Cuauhtémoc Doctores México City Mexico 06726

Type(s)

Public

Contact name

Miss Geovana Martin

ORCID ID

https://orcid.org/0000-0003-3583-6126

Contact details

Dr. Balmis 148 Cuauhtémoc Doctores México City Mexico 06726

Additional identifiers

Protocol serial number

DI/16/105-B-04-128

Study information

Scientific Title

A prospective randomized trial, Aerobic Versus ANaerobic exercise and oral nuTritional supplEmentation in nutritional status and physical function of adults HEMOdialysis Patients: the AVANTE-HEMO study

Acronym

AVANTE-HEMO

Study objectives

Hemodialysis (HD) patients who perform resistance exercise (RE) + receive oral nutritional supplementation (ONS) will have better parameters of nutritional status and physical function than patients of the aerobic plus oral nutritional supplementation group and oral nutritional supplementation group.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Hospital General de México, Dr. Eduardo Liceaga ethics committee, 28/09/2016, ref: DI/16/105-B-04-128

Study design

Parallel single-centre randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Hemodialysis

Interventions

Method of randomization: A blocked design will be used, using an online statistical computing web programming (www.randomizer.org) to generate the randomization schedule, 14 sets of 3 unique numbers per set range from 1 to 3 (representing the three treatment arms).

Control arm

During the HD sessions, patients receive a can of a specialized oral nutritional supplement for maintenance dialysis patients that consists of 480 kcal, 20 g protein and 20 g lipids (Enterex RNL, Pisa Laboratories). Enterex RNL contains omega-3 and omega-6 fatty acids, maltodextrin and sucralose.

Aerobic exercise arm

Patients of this group receive the same specialized oral nutritional supplement during the HD session, half of the a can is administered during the first hour of the HD session and the other half after the AE routine. Aerobic exercise consists of pedaling a stationary bike for the first two

hours of the HD session, the optimal exercise time is established during the first session, then is gradually increased to reach 20-30 minutes. The scale of perceived exertion Borg (6-20) is used to establish the intensity of the exercise (moderate intensity, 12-13).

Resistance exercise arm

Patients of this group receive the same specialized oral nutritional supplement during the HD session, half of the a can is administered during the first hour of the HD session and the other half after the RE routine. Patients are trained for three months according to an adaptation to the program "Exercise: A Guide for People on Dialysis". Four types of resistance exercise are performed during the second hour of the HD session (four sets of 20 repetitions for 40 min). TheraBand Latex Resistance Bands are used to individualize the exercise. The scale of perceived exertion Borg (6-20) is used to establish the intensity of the exercise (moderate intensity, 12-13)

Frequency and duration of treatment: Two times per week for 12 weeks (40 minutes of resistance exercise and 30 minutes of aerobic exercise)
Follow-up: 12 weeks

Intervention Type

Mixed

Primary outcome(s)

Measured at baseline and 3 months:

- 1. Weight: The weight reached by the patient at the end of a session which has removed the maximum fluid without inducing hypotension (seca 676 wheelchair scale, wireless)
- 2. Body mass index: body mass divided by the square of the body height
- 3. Anthropometric measurements: skinfold thickness (biceps, triceps, subescapular, supra-iliac) using a Lange skin fold caliper to estimate:
- 3.1. Fat mass percentage: calculated using Siri's (1956) equation = (4.95/density)-4.50*100
- 3.2. Mid-upper arm circumference: measured at the mid-point between the tip of the shoulder and the tip of the elbow (olecranon process and the acromium)
- 3.3. Mid arm muscle circumference = mid arm circumference (π x triceps skinfold thickness)
- 3.4. Bone-free arms muscle area: males = $[(midarm\ circumference\ (cm) \pi\ x\ triceps\ (cm)]2/4\ \pi)$ 10; females = $[(midarm\ circumference\ (cm) \pi\ x\ triceps\ (cm)]2/4\ \pi)$ 6.5
- 4. Handgrip strength, measured using hand dynamometry (Smedley III, Takei Scientific Instruments, Niigata City, Japan)
- 5. Serum albumin, measured using the bromocresol purple albumin assay
- 6. Physical function, measured using the six-minute walk test according to the American Thoracic Society Guidelines, Time Up and Go Test (Podsiadlo, D et al. J Am Geriatr Soc, 1191;39:142-148), five times sit to stand test
- 7. Bioelectrical impedance analysis (resistance, reactance and phase angle), measured using conventional impedance (single frequency; Quantum System, RJL Systems, Clinton Township, MI, USA)

Key secondary outcome(s))

Measured at baseline and 3 months:

- 1. Quality of life, measured using the Kidney Disease Quality of Life Short Form (KDQOF-SF)
- 2. C Reactive Protein, measured using the nephelometric assay

Completion date

01/09/2017

Eligibility

Key inclusion criteria

- 1. Regular HD two or more times a week
- 2. Signed informed consent
- 3. Any gender
- 4. Age >18 to <45 years

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

45

Key exclusion criteria

- 1. Patients with an amputation
- 2. Hospitalization in the last 3 months
- 3. Unsatisfactory attendance at HD sessions
- 4. Pregnancy
- 5. Excessive pallor
- 6. Severe dyspnea
- 7. Femoral fistula
- 8. Arrhythmias
- 9. Precordial pain
- 10. Orthopedic or neurological compromises or cognitive alterations affecting their participation in the study

Date of first enrolment

01/11/2016

Date of final enrolment

31/01/2017

Locations

Countries of recruitment

Mexico

Study participating centre Hospital General de México, Dr Eduardo Liceaga

Dr. Balmis 148 Cuauhtémoc Doctores Ciudad de México, D.F. Mexico 06726

Sponsor information

Organisation

Hospital General de México, Dr Eduardo Liceaga

ROR

https://ror.org/01php1d31

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Hospital General de México, Dr Eduardo Liceaga

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Geovana Martin Alemañy. The trialists can share data (variables) represented in the subsequent results publication and if someone needs information for a systematic review with meta-analysis, the data will be available before the publication of the original article. The trialists are going to share anonymous information and the datasets generated and/or analysed during the current study during this study will be included in the subsequent results publication.

IPD sharing plan summary

Available on request

Study outputs

Output type

Details results

Date created Date added Peer reviewed? Patient-facing?

Results article		01/03/2020	15/10/2019 Yes	No
Basic results			12/12/2018 No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025 No	Yes