

# Use of preoperative desmopressin in preventing bleeding in patients treated with selective serotonin reuptake inhibitors (SSRIs)

<b>Submission date</b> 20/12/2005	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 20/12/2005	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 02/09/2008	<b>Condition category</b> Surgery	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr S.C. Marczinski

**Contact details**  
Willy Brandtlaan 10  
Ede  
Netherlands  
6716 RP  
+31 (0)318 435546  
Marczinskis@zgv.nl

## Additional identifiers

**Protocol serial number**  
UMCU 04-298; NTR175

## Study information

**Scientific Title**

**Study objectives**

Not provided at time of registration

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Not provided at time of registration

**Study design**

Randomised, double-blinded, placebo controlled, parallel group trial

**Primary study design**

Interventional

**Study type(s)**

Not Specified

**Health condition(s) or problem(s) studied**

Primary haemostasis disorder

**Interventions**

Pre-operative infusion placebo/desmopressin:

1. Less than 50 kg: 15 µg
2. 50 - 100 kg: 30 µg
3. Greater than 100 kg: 45 µg

**Intervention Type**

Drug

**Phase**

Not Specified

**Drug/device/biological/vaccine name(s)**

Desmopressin

**Primary outcome(s)**

Pre- and post-operative blood loss.

**Key secondary outcome(s)**

1. Peri- and post-operative haemoglobin
2. Number of peri-operative blood transfusion
3. Peri-operative fluid infusion
4. Post-operative drainage

**Completion date**

01/06/2007

**Eligibility**

**Key inclusion criteria**

1. Patients aged over 18 who receive a serotonergic antidepressant (fluvoxamine, fluoxetine, paroxetine, sertraline, venlafaxine, clomipramine, citalopram) at least started two weeks before the surgery
2. Surgery: orthopedic, abdominal, breast

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Sex**

Not Specified

**Key exclusion criteria**

1. No informed consent
2. Disorder in primary haemostasis
3. Hyponatraemia (sodium [serum] less than 130 mmol/l)
4. Laparoscopic surgery
5. Use of vitamin K antagonists, aspirin, iron supplements, methotrexate, heparin
6. Acute coronary syndrome (unstable angina and myocardial infarction)
7. Spinal anaesthesia during surgery

**Date of first enrolment**

01/01/2005

**Date of final enrolment**

01/06/2007

**Locations****Countries of recruitment**

Netherlands

**Study participating centre**

Willy Brandtlaan 10

Ede

Netherlands

6716 RP

# Sponsor information

## Organisation

Hospital Gelderse Vallei (The Netherlands)

## ROR

<https://ror.org/03862t386>

# Funder(s)

## Funder type

Hospital/treatment centre

## Funder Name

Hospital Gelderse Vallei (The Netherlands)

# Results and Publications

Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Not provided at time of registration