

# Direct pulp capping in primary molars: a feasibility study

<b>Submission date</b> 11/10/2018	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 25/10/2018	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 17/11/2023	<b>Condition category</b> Oral Health	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Exposures of the dental pulp (the centre of the tooth) are common in dental practice and can be challenging especially in primary (baby) teeth. Direct pulp capping (DPC) is a technique used to cover the exposed pulp. It is still controversial as a treatment. However, calcium hydroxide (CH) is considered the gold standard in dental pulp treatments. The introduction of a combination of antibiotics could be promising in pulp treatments. However, improving DPC outcomes would provide an easier and less invasive treatment option for pulp exposures in these teeth. The aim of this study is to assess the feasibility of conducting a future trial on DPC using 3Mix-MP (triple antibiotic paste) compared with calcium hydroxide, and to estimate the number of participants needed for the main trial on the basis of the success rate.

### Who can participate?

Children who have primary molars (back teeth) with large carious lesions (cavities) that could be restored with composite

### What does the study involve?

The participants' teeth are randomly allocated into two groups to be treated with DPC with either 3Mix-MP or calcium hydroxide (CH). All teeth are restored with composite resin and assessed at 3 and 6 months after treatment.

### What are the possible benefits and risks of participating?

Treating these teeth could prevent inflammation/abscesses and the associated signs like pain and swelling. No harms are encountered since all failed cases are treated with another method (non-instrumental endodontic treatment or traditional pulpectomy).

### Where is the study run from?

Tishreen University (Syria)

### When is the study starting and how long is it expected to run for?

August 2016 to June 2017

Who is funding the study?  
Tishreen University (Syria)

Who is the main contact?

1. Dr Nabih Raslan
2. Dr Hasan Ali

## Contact information

### Type(s)

Scientific

### Contact name

Dr Nabih Raslan

### ORCID ID

<https://orcid.org/0000-0001-9967-9575>

### Contact details

Department of paediatric Dentistry  
Faculty of Medical Dentistry, Tishreen University  
Lattakia  
Syria  
00963

### Type(s)

Scientific

### Contact name

Dr Hasan Ali

### ORCID ID

<https://orcid.org/0000-0001-6959-7552>

### Contact details

Department of paediatric Dentistry  
Faculty of Medical Dentistry, Tishreen University  
Lattakia  
Syria  
00963

## Additional identifiers

### Protocol serial number

U1111-1222-2654

## Study information

Scientific Title

Direct pulp capping (DPC) in primary molars using three antibiotics combination 3mix-mp: a randomized controlled pilot trial

### **Study objectives**

Is conducting a large randomized clinical trial feasible and what is the proper sample size?

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

The institutional review board of Tishreen University, 02/08/2016, Approval No. 3179

### **Study design**

Pilot randomized double-blind parallel two-arm control trial

### **Primary study design**

Interventional

### **Study type(s)**

Treatment

### **Health condition(s) or problem(s) studied**

Management of large carious lesions in primary teeth

### **Interventions**

Twenty molars in eight cooperative healthy children were randomly allocated into two groups using a simple randomization process. In both groups, teeth were anesthetized and isolated with a rubber dam. All carious tissues were removed using a small round bur mounted on a high-speed handpiece and copious water supply.

After observing an exposure site, a cotton pellet soaked with sodium hypochlorite was applied for five minutes. The dressing agent was chosen randomly and the teeth were divided into two groups: 3Mix-MP (Group A) and calcium hydroxide powder mixed with saline CH (Group B). The dressing agent was prepared in creamy form and was applied to the exposure site before it was gently pressed into a thin layer on the cavity floor using a cotton pellet. The agent was covered with a thin layer of light cured calcium hydroxide. Composite resin restorations were placed, and then were finished after removing the rubber dam with diamond finishing burs; consequently, polishing was performed using silicone polishers.

The participants were asked to attend follow-up examination, and were clinically evaluated at 3 and 6 months post-treatment and were evaluated radiographically at 6 months.

### **Intervention Type**

Procedure/Surgery

### **Primary outcome(s)**

Feasibility, estimated using the following measurements:

1. Proportion of eligible children among those examined, calculated after completion of the assigned number of participants
2. Proportion of children who completed the 6-month follow-up among those randomized, calculated after completion of the assigned period

**Key secondary outcome(s))**

Success rates of the materials, which will be used to calculate the sample size for the definitive randomized controlled trial. Criteria used: absence/presence of clinical and radiographic signs, proportions calculated using simple formulas. Measured at the end of the treatment session as a baseline; the participants were clinically evaluated at 3 and 6 months post-treatment and evaluated radiographically after 6 months

**Completion date**

23/06/2017

**Eligibility****Key inclusion criteria**

1. Cooperative healthy children
2. Vital primary molars with large carious lesions that could be restored with composite
3. No history of spontaneous pain, pathologic mobility, redness or swelling of the vestibule, draining sinus tracts, or sensitivity to vestibular palpation
4. Absence of internal or external root resorption or inter-radicular/apical radiolucency
5. Both children and their parents had to be able to attend a 6-month follow-up procedure

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Child

**Sex**

All

**Total final enrolment**

17

**Key exclusion criteria**

Any clinical or radiographic signs indicating non-vital teeth

**Date of first enrolment**

20/08/2016

**Date of final enrolment**

16/12/2016

**Locations****Countries of recruitment**

Syria

**Study participating centre**  
**Tishreen University**  
Department of Pediatric Dentistry  
Lattakia  
Syria  
00963

## Sponsor information

**Organisation**  
Tishreen University

**ROR**  
<https://ror.org/04nqts970>

## Funder(s)

**Funder type**  
University/education

**Funder Name**  
Tishreen University

## Results and Publications

### Individual participant data (IPD) sharing plan

The datasets generated and/or analysed during this study will be included in the subsequent results publication. Individual participant data that underlie the results, after deidentification (text, tables, figures, and appendices), would be available immediately after publication from [raslan.nabih@tishreen.edu.sy](mailto:raslan.nabih@tishreen.edu.sy).

### IPD sharing plan summary

Other

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>		04/02/2021	27/09/2021	Yes	No
<a href="#">Abstract results</a>	P026	01/09/2019	17/11/2023	No	No