# Calabrian functional tomato sauce for the treatment of non-alcoholic fatty liver disease

Submission date	Recruitment status  No longer recruiting	Prospectively registered		
27/05/2025		Protocol		
Registration date	Overall study status Completed  Condition category Digestive System	Statistical analysis plan		
04/06/2025		Results		
Last Edited		Individual participant data		
03/06/2025		[X] Record updated in last year		

## Plain English summary of protocol

Background and study aims

Non-alcoholic fatty liver disease (liver steatosis) is the most common cause of liver-related illness and death worldwide. However, no approved drug treatments currently exist for this condition. Lycopene, a powerful antioxidant found in tomatoes, has shown potential benefits for liver health. OsteoCol® is a lycopene-enriched tomato sauce obtained through on-vine ripening, which increases its natural lycopene content. This study aims to evaluate the effect of daily consumption of OsteoCol® tomato sauce as a functional food for the treatment of patients with liver steatosis.

## Who can participate?

Adult patients adults of both genders, aged 30-75 years, with a diagnosis of NAFLD.

#### What does the study involve?

Participants are allocated with a simple randomization using computer-generated random numbers to the intervention or the control group. Ninety-eight adults were randomly assigned to two groups for 12 weeks, as follows:

Intervention group: Participants received a functional tomato sauce (OsteoCol®), naturally enriched with lycopene. Twenty-five participants consumed 80 g/day, while twenty-four participants consumed 160 g every other day.

Control group: Participants received a commercial tomato sauce. Twenty-five participants consumed 80 g/day, while twenty-four participants consumed 160 g every other day. The primary outcome measured was a change in the amount of intrahepatic fat, assessed by CAP score, after a 12-week intervention between groups.

What are the possible benefits and risks of participating? The potential benefit is a reduction in the amount of liver fat and an improvement in the severity of NAFLD. There are no risks to participants.

Where is the study run from? University Magna Grecia (Italy)

When is the study starting and how long is it expected to run for? From April 2022 to August 2023.

Who is funding the study?

This study was supported by a Grant from Calabria Region, Italy (Bando INGEGNO - POR CALABRIA FESR-FSE 2014/2020-SALNAF "Functional Calabrian Tomato Sauce Naturally Rich in Lycopene for the Treatment of Non-Alcoholic Fatty Liver Disease").

Who is the main contact? Prof. Arturo Pujia, pujia@unicz.it

## Contact information

## Type(s)

Public, Scientific, Principal Investigator

#### Contact name

Prof Arturo Pujia

#### **ORCID ID**

https://orcid.org/0000-0003-0763-4930

#### Contact details

Viale Europa, Campus Universitario S. Venuta Catanzaro Italy 88100 + 39 0961 3697080 pujia@unicz.it

## Additional identifiers

## **EudraCT/CTIS** number

Nil known

**IRAS** number

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

124/2022/CE

# Study information

#### Scientific Title

SALNAF – Calabrian functional tomato sauce naturally rich in lycopene for the treatment of non-alcoholic fatty liver disease

## **Acronym**

## **Study objectives**

Non-alcoholic fatty liver disease (liver steatosis) is the most common cause of liver-related illness and death worldwide. However, no approved drug treatments currently exist for this condition. Lycopene, a powerful antioxidant found in tomatoes, has shown potential benefits for liver health. OsteoCol® is a lycopene-enriched tomato sauce obtained through on-vine ripening, which increases its natural lycopene content. This study aims to evaluate the effect of daily consumption of OsteoCol® tomato sauce as a functional food for the treatment of patients with liver steatosis.

## Ethics approval required

Ethics approval required

## Ethics approval(s)

Approved 21/04/2022, Comitato Etico Regione Calabria Sezione Area Centro (A.O.U. Mater Domini in Via Tommaso Campanella, 115, Catanzaro, 88100, Italy; +39 (0)961 712 111; comitatoetico@hotmail.it), ref: 124/2022/CE

## Study design

Interventional randomized controlled trial

## Primary study design

Interventional

## Secondary study design

Randomised parallel trial

## Study setting(s)

Community

## Study type(s)

Treatment

## Participant information sheet

See study outputs table

## Health condition(s) or problem(s) studied

Treatment of hepatic steatosis in patients with NAFLD

#### **Interventions**

Participants were enrolled after providing written informed consent and were randomly assigned (by computer-generated random numbers) in a 1:1 ratio to one of two treatment groups:

Intervention group: Participants received a functional tomato sauce (OsteoCol®), naturally enriched with lycopene. Twenty-five participants consumed 80 g/day, while twenty-four participants consumed 160 g every other day.

Control group: Participants received a commercial tomato sauce. Twenty-five participants consumed 80 g/day, while twenty-four participants consumed 160 g every other day

## Intervention Type

#### Other

## Primary outcome measure

Amount of intrahepatic fat (i.e. Controlled Attenuation Parameter value, CAP score), assessed by Transient Vibration-Controlled Elastography (Fibroscan), at baseline and 12 weeks

## Secondary outcome measures

At baseline and 12 weeks:

- 1. Severity of NAFLD, assessed by FibroScan
- 2. LDL-cholesterol, measured using biochemical assays, after 12 weeks of treatment
- 3. Inflammation markers, measured using biochemical assays, after 12 weeks of treatment
- 4. Lipid accumulation (i.e., Lipid Accumulation Product, LAP) and cardiovascular risk markers (i.e., Atherogenic Index of Plasma, AIP)
- 5. Antioxidant status parameters, evaluated using ELISA kits

## Overall study start date

21/04/2022

## Completion date

29/08/2023

## **Eligibility**

## Key inclusion criteria

- 1. NAFLD defined by a CAP score value ≥248 dB/m
- 2. Age 30 75 years

## Participant type(s)

Patient

## Age group

Adult

## Lower age limit

30 Years

## Upper age limit

75 Years

#### Sex

Both

## Target number of participants

98

#### Total final enrolment

98

### Key exclusion criteria

- 1. Alcohol intake >30 g/day for men or >20 g/day for women
- 2. BMI  $\geq$  35 kg/m<sup>2</sup>
- 3. Allergy to tomatoes and nickel
- 4. HBV, HCV, HIV infection
- 5. Pregnancy
- 6. Type 2 Diabetes Mellitus (T2DM)
- 7. Triglycerides > 250 mg/dL
- 8. Gastroesophageal reflux disease
- 9. Hepatic and renal failure
- 10. Cirrhosis of the liver and other chronic liver diseases
- 11. Gallstones
- 12. Chronic intestinal diseases
- 13. Chronic debilitating diseases
- 14. Parenteral nutrition
- 15. Use of specific drugs (amiodarone, antiretrovirals, corticosteroids, methotrexate, tamoxifen, valproate, or lipid-lowering drugs for less than 6 weeks)
- 16. Use of food supplements or functional foods aimed at reducing hepatic steatosis

#### Date of first enrolment

27/03/2023

## Date of final enrolment

31/05/2023

## Locations

#### Countries of recruitment

Italy

88100

## Study participating centre

Nutrition Unit of the "R. Dulbecco" Azienda University Hospital

Viale Europa - Campus Universitario S. Venuta Catanzaro Italy

# Sponsor information

## Organisation

Regione Calabria

#### Sponsor details

Viale Europa - Zona Tramontana - Cittadella Regionale Catanzaro Italy 88100 +39 0961 858524 programmazione@regione.calabria.it

## Sponsor type

Government

#### Website

https://www.regione.calabria.it

#### **ROR**

https://ror.org/03swz0133

# Funder(s)

## Funder type

Government

#### **Funder Name**

Regione Calabria - SALNAF - CUP: J39J22005540005

## **Results and Publications**

## Publication and dissemination plan

Publication and dissemination plan: Publication in journal in English with impact factor; mass media.

## Intention to publish date

01/06/2025

## Individual participant data (IPD) sharing plan

Individual participant data (IPD) that underlie the results reported in the main publication (including baseline data, primary outcomes, and secondary outcomes) will be made available after de-identification. The data will be accessible beginning 6 months after publication and for up to 5 years, to researchers who provide a methodologically sound proposal. Data will be shared in accordance with participant consent and applicable ethical guidelines. A data-sharing agreement may be required. Requests should be directed to pujia@unicz.it.

## IPD sharing plan summary

Available on request

## **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet			03/06/2025	No	Yes