# Online social support for early psychosis - A pilot feasibility trial of the PSYCHE-MOST intervention.

Submission date	<b>Recruitment status</b> No longer recruiting	<ul><li>Prospectively registered</li></ul>		
09/08/2024		☐ Protocol		
Registration date	Overall study status Ongoing Condition category Mental and Behavioural Disorders	Statistical analysis plan		
05/12/2024		☐ Results		
Last Edited		Individual participant data		
05/12/2024		[X] Record updated in last year		

#### Plain English summary of protocol

Background and study aims

With a noticeable increase in individuals seeking help for mental health concerns in recent times, digital health interventions—mental health support delivered through web-based or mobilebased platforms—hold significant promise in enhancing outcomes, broadening access, and addressing the growing demand for mental health services. One such approach aimed at enhancing mental health recovery among young adults is Moderated Online Social Therapy (MOST). Initially developed as a digital mental health platform available via both a web interface and a dedicated app, MOST offers a low-intensity, cost-effective, and engaging method to sustain or augment the benefits of specialized Early Intervention for Psychosis (EIP) services. Studies have shown its effectiveness in facilitating return to education and employment while reducing the need for emergency care. It has also undergone a trial involving young individuals aged 12 to 25 years old who sought help for mental health issues. MOST comprises evidencebased online therapy content supplemented by therapist interaction, alongside a community feature resembling a Facebook-style platform, supported and moderated by peer support workers. The objective of the current study is twofold: firstly, to assess the feasibility and acceptability of MOST among individuals who are currently attending an EIP service, and secondly, to provide preliminary data on the effectiveness of MOST, laying the groundwork for a definitive randomized controlled trial.

#### Who can participate?

Patients aged between 18 and 35 years old who are clinically stable, able to give consent and self-reporting mental health difficulties of longer than one year in duration

#### What does the study involve?

MOST (the intervention being assessed) consists of

1. Interactive online therapy modules based on third-wave cognitive behavioural therapy and primarily targeting social functioning by e.g., fostering self-efficacy (identifying personal strengths based on the strengths-based framework), positive emotions and subjective well-being (e.g., practicing mindfulness and self-compassion), or positive connections with others (e.g., focusing on empathy skills). Completing this therapy journey is supported by clinicians

available by telephone/video chat on a fortnightly basis.

2. An online social network or 'Café' to foster social support. Participants are encouraged to communicate with one another and with peer and expert moderators. This is moderated by clinicians and led by 'peer-support workers' with lived experience and informed by the evidence-based problem-solving framework.

A further feature of MOST is an online group function to enable users to nominate issues (e.g., 'how to break through shyness and make new friends?'), which are discussed in moderated groups through structured phases (e.g., brainstorming, pros and cons and wrap-up). The intervention is 26 weeks in duration with follow-ups at 3 and 6 months.

What are the possible benefits and risks of participating?

Possible benefits include (1) a reduction in clinical symptoms of anxiety and low mood, (2) opportunities to receive support from an online therapist while completing a CBT-informed online therapy intervention, and (3) opportunities to receive peer support from a peer support worker.

Possible, risks include (1) distress caused by participating in the assessment, which asks questions about early childhood adversity and clinical symptom severity, and (2) low-level distress caused by participating in an online intervention that may not meet the psychological needs of the individual.

Where is the study run from? University of Galway (Ireland)

When is the study starting and how long is it expected to run for? September 2019 to December 2025

Who is funding the study? Health Research Board (Ireland)

Who is the main contact? Dr Gary Donohoe, gary.donohoe@universityofgalway.ie

### Contact information

#### Type(s)

Public, Scientific, Principal investigator

#### Contact name

Prof Gary Donohoe

#### **ORCID ID**

https://orcid.org/0000-0003-3037-7426

#### Contact details

School of Psychology, University of Galway, University road Galway Ireland H91Tk33 +3530866008405 gary.donohoe@nuigalway.ie

## Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

RL-2020-007

# Study information

#### Scientific Title

Improving Psychosocial Supports in youth mental health (PSYCHE) - A pilot feasibility randomised controlled trial of Moderated Online Social Therapy (MOST) for improving social function in early psychosis

#### Acronym

**PSYCHE-MOST-EIP** 

#### Study objectives

The trial explores the feasibility and acceptability of delivering MOST to individuals who are currently attending early intervention for psychosis services in Ireland. The main study hypothesis is that (a) it will be possible to recruit n=30 and retain a majority of these at follow up.

#### Ethics approval required

Ethics approval required

#### Ethics approval(s)

approved 14/10/2020, Galway University Hospital Clinical Research Ethics Committee (Room 2, 2nd Floor, HR building, Merlin Park, Galway, H91N973, Ireland; +353-91-731990; colette. collins@hse.ie), ref: CA2468

#### Study design

Interventional pilot feasibility multi-center randomized controlled study

#### Primary study design

Interventional

#### Study type(s)

Treatment

#### Health condition(s) or problem(s) studied

**Psychosis** 

#### **Interventions**

This trial is a pilot randomised controlled study recruiting patients attending an early intervention for psychosis service in Ireland. The trial explores the feasibility, acceptability, and effectiveness of delivering MOST to participants who attended the EIP service.

MOST (The intervention being assessed) consists of:

- 1. Interactive online therapy modules based on third-wave cognitive behavioural therapy and primarily targeting social functioning by e.g., fostering self-efficacy (identifying personal strengths based on the strengths-based framework), positive emotions and subjective well-being (e.g., practicing mindfulness and self-compassion), or positive connections with others (e.g., focusing on empathy skills). Completing this therapy journey is supported by clinicians available by telephone/video chat on a fortnightly basis.
- 2. An online social network or 'Café' to foster social support. Participants are encouraged to communicate with one another and with peer and expert moderators. This is moderated by clinicians and led by 'peer-support workers' with lived experience and informed by the evidence-based problem-solving framework.

A further feature of MOST is an online group function to enable users to nominate issues (e.g., 'how to break through shyness and make new friends?'), which are discussed in moderated groups through structured phases (e.g., brainstorming, pros and cons and wrap-up).

The control group receives treatment as usual, defined as continuing to receive any care that they were receiving prior to the intervention.

The intervention is 26 weeks duration. The follow-up periods are 3 and 6 months post baseline. Randomisation is based on the sealed envelope online tool.

#### Intervention Type

Behavioural

#### Primary outcome(s)

Feasibility outcomes: Rates of recruitment and retention measured using data recorded in patient records at baseline, 3 months (12 weeks) and 6 months (26 weeks)

#### Key secondary outcome(s))

The following secondary outcome measures will be measured during interviews with the participant at baseline, 3 months (12 weeks) and 6 months (26 weeks):

Demographic/Clinical measures:

- 1. Data on age, sex, living situation, educational experience, and employment history measured using and assessment Interview
- 2. Drug use measured using the Drug Use Disorders Identification Test (DUDIT)
- 3. Alcohol use measured using the Alcohol Use Disorders Identification Test (AUDIT)
- 4. Severity of positive and negative symptoms in psychosis measured using the Scale for the Assessment of Positive Symptoms (SAPS) and the Scale for the Assessment of Negative Symptoms (SANS)
- 5. Anxiety symptoms measured using the Generalized Anxiety Disorder Assessment (GAD-7)
- 6. Depressive symptoms measured using the Patient Health Questionnaire (PHQ-9)
- 7. Self-care capability measured using the Independent Living Scale
- 8. Time spent on various daily activities measured using the Time Use Survey
- 9. Social and occupational functions measured using the Social and Occupational Functioning

#### Assessment Scale (SOFAS)

- 10. Social and occupational function measured using the Global Assessment of Functioning (GAF)
- 11. Occupational functioning, social functioning, and symptom severity measured using the Mental Illness Research, Education, and the Clinical Center version of the Global Assessment of Functioning (MIRECC GAF)
- 12. Loneliness measured using the UCLA Loneliness Scale

#### Cognition measures:

- 1. Theory of mind measured using Reading the Eyes in the Mind
- 2. Emotion recognition measured using the Cambridge Neuropsychological Test Automated Battery Emotional Recognition Task (CANTAB ERT)
- 3. General cognitive ability prior to illness measured using the Test of Premorbid Functioning (TOPF)
- 4. Verbal episodic memory measured using the Weschler Memory Scale (WMS) Logical Memory Test
- 5. Visuospatial abstract reasoning measured using the Weschler Matrix Reasoning
- 6. Verbal abstract reasoning measured using the Weschler Similarities
- 7. Visuospatial working memory measured using the Cambridge Neuropsychological Test Automated Battery Spatial Working Memory Test (CANTAB SWM)

#### Completion date

30/12/2025

# Eligibility

#### Key inclusion criteria

- 1. Aged between 18 and 35 years old
- 2. Clinically stable and having the ability to give consent
- 3. Self-reporting mental health difficulties of longer than one year in duration

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Upper age limit

35 years

#### Sex

Αll

#### Key exclusion criteria

- 1. History of organic impairment (including IQ <70), or history of a head injury with loss of consciousness > 5-minute duration
- 2. Drug or alcohol dependent

# Date of first enrolment 01/04/2021

Date of final enrolment 31/12/2024

#### Locations

# **Countries of recruitment**Ireland

Study participating centre Sligo/Leitrim EIP center Dublin road Sligo Ireland F91 X92V

Study participating centre
Galway Adult Mental Health Service
University Hospital Galway
Galway
Ireland
H91YR71

# Sponsor information

#### Organisation

Ollscoil na Gaillimhe – University of Galway

#### **ROR**

https://ror.org/03bea9k73

# Funder(s)

Funder type

#### Government

#### **Funder Name**

Health Research Board

#### Alternative Name(s)

Health Research Board, Ireland, An Bord Taighde Sláinte, HRB

#### **Funding Body Type**

Government organisation

#### **Funding Body Subtype**

National government

#### Location

Ireland

## **Results and Publications**

#### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from the study PI, Prof. Gary Donohoe, gary.donohoe@universityofgalway.ie

#### IPD sharing plan summary

Available on request

#### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes