# Exploring the use of a stroke patient concerns inventory during stroke follow-up appointments

Submission date	Recruitment status  No longer recruiting	[X] Prospectively registered		
07/11/2024		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
22/11/2024	Ongoing	Results		
<b>Last Edited</b> 21/11/2024	<b>Condition category</b> Nervous System Diseases	Individual participant data		
		[X] Record updated in last year		

#### Plain English summary of protocol

Background and study aims

Stroke survivors experience ongoing physical, psychological and social concerns affecting quality-of-life. There is a lack of systems and tools that allow healthcare staff to identify stroke survivors' concerns and so they are often not adequately addressed. Stroke outpatient follow-up appointments are time-limited, reducing opportunity for patients to discuss, and staff to identify, concerns.

The Patient Concerns Inventory (PCI), is used in cancer-care appointments. The PCI lists potential concerns for patients to select from before their appointment which they would like to discuss. Research shows that using the PCI increased satisfaction with appointments and improved patient quality-of-life, without lengthening appointment time. The PCI has been adapted for stroke (Stroke Patient Concerns Inventory (sPCI)) and a large study is needed to see if it is also effective. Firstly, the study design and the sPCI need to be tested in a smaller study. This study, funded by the NIHR, aims to determine if the sPCI can be used in stroke outpatient services by both staff and patients.

#### Who can participate?

Four stroke services performing follow-up appointments will recruit a total of 96 participants, who are: stroke survivors within 7-months of stroke, aged 18 years and over, and attending their first follow-up appointment.

#### What does the study involve?

Two sites will receive training in the sPCI (using it during appointments to discuss concerns and manage them appropriately), and provide their participants the sPCI to complete before their appointment (in addition to their usual practice). The remaining two sites will continue their usual practice. Data collection will include: participants' demographic and stroke information, and quality-of-life at baseline; completed sPCI and participants' feedback on its use immediately after the appointment; participants' satisfaction and empowerment within one-week of the appointment; and participants' quality-of-life and use of health services three-months later. A selection of participants and staff will be interviewed about their experiences of the study.

What are the possible benefits and risks of participating?

Benefits: Not provided at time of registration

Risks: The main disadvantages are that this will take some of your time and you may find thinking or talking about some of the issues is upsetting. If this happens, the research practitioner or university researcher can provide suggestions of who you can talk to, such as helplines or your GP.

Where is the study run from? University of Central Lancashire (UK)

When is the study starting and how long is it expected to run for? May 2024 to October 2025

Who is funding the study? National Institute for Health and Care Research (NIHR) (UK).

Who is the main contact? esaviciute@uclan.ac.uk

# Contact information

#### Type(s)

Principal Investigator

#### Contact name

Miss Kulsum Patel

#### **ORCID ID**

http://orcid.org/0000-0002-0865-3941

#### Contact details

Brook Building 247, University of Central Lancashire Preston United Kingdom PR1 2HE +44 1772 893635 Kpatel@UCLan.ac.uk

# Additional identifiers

# **EudraCT/CTIS** number

Nil known

#### IRAS number

344756

#### ClinicalTrials.gov number

Nil known

#### Secondary identifying numbers

# Study information

#### Scientific Title

Use of the Stroke Patient Concerns Inventory in Stroke Outpatients Clinics: a feasibility cluster randomised controlled trial

#### Acronym

**SPaCIS** 

#### **Study objectives**

To determine whether a future, larger study can be conducted to explore how effective the Stroke Patient Concerns Inventory (sPCI) is when used in stroke outpatient clinics.

#### Ethics approval required

Ethics approval required

#### Ethics approval(s)

Approved 04/10/2024, Wales REC 3 (Health and Care Research Wales Support and Delivery Centre, Castlebridge 4, 15-19 Cowbridge Road East, Cardiff, CF11 9AB, United Kingdom; +44 2922 940963; Wales.REC3@wales.nhs.uk), ref: 24/WA/0259

#### Study design

Interventional non-randomized

#### Primary study design

Interventional

# Secondary study design

Non randomised study

#### Study setting(s)

Hospital, Medical and other records

# Study type(s)

Screening

#### Participant information sheet

See outputs table

#### Health condition(s) or problem(s) studied

Stroke

#### **Interventions**

Four stroke services performing follow-up appointments will recruit a total of 96 participants, who are: stroke survivors within 7-months of stroke, aged 18 years and over, and attending their first follow-up appointment. Two sites will receive training in the sPCI (using it during

appointments to discuss concerns and manage them appropriately), and provide their participants the sPCI to complete before their appointment (in addition to their usual practice). The remaining two sites will continue their usual practice.

Data collection will include: participants' demographic and stroke information, and quality-of-life at baseline; completed sPCI and participants' feedback on its use immediately after the appointment; participants' satisfaction and empowerment within one-week of the appointment; and participants' quality-of-life and use of health services three-months later. A selection of participants and staff will be interviewed about their experiences of the study.

#### Intervention Type

Behavioural

#### Primary outcome measure

- 1. The rate of completed sPCIs is measured using the number of sPCIs completed at the clinical follow-up appointment
- 2. sPCI use in consultation is measured using a participant's immediate feedback question on whether they recall the use of the sPCI in their consultation and audio recordings of consultations at clinical follow-up appointments and immediate feedback
- 3. Participant feedback on the use of sPCI is measured using an immediate feedback question on whether they recall the use of the sPCI in their consultation at immediate feedback
- 4. Patients completing sPCI is measured using the number of sPCIs completed at clinical followup appointments and process evaluation interviews
- 5. sPCI use during the consultation is measured using a participant's immediate feedback question on whether they recall the use of the sPCI in their consultation and audio recordings of consultations at clinical follow-up appointments and process evaluation interviews
- 6. Staff trained in sPCI use is measured using the number of staff trained during the training period and process evaluation interviews
- 7. sPCIs provided before the appointment are measured using the number of sPCIs sent to patients at the clinical follow-up appointment
- 8. Completed sPCIs available to staff conducting consultations are measured using data on consultation CRF about whether sPCI was available at the clinical follow-up appointment
- 9. Sites recruited are measured using the number of sites in the study at the end of the recruitment period
- 10. Participants recruited are measured using the number of participants recruited to the study at the end of the recruitment period
- 11. Eligibility rate is measured using screening logs at the end of the recruitment period
- 12. The recruitment rate is measured using screening logs and the number of participants recruited to the study at the end of the recruitment period
- 13. Retention/completion rates are measured using the number of completed 3-month questionnaires at the end of the data collection period
- 14. Questionnaire completion rates are measured using the number of completed 1-week and 3-month questionnaires at the end of the data collection period and process evaluation interviews
- 15. Patient satisfaction, empowerment, and quality of life are measured using the Patient Satisfaction Survey and Southampton Stroke Self-Management Scale at 1-week, and EQ-5D-5L at 3-months of data collection and process evaluation interviews
- 16. Case report form completion rates are measured using the number of completed 1-week and 3-month questionnaires at the end of the data collection period and process evaluation interviews
- 17. Stop/Go criteria is measured using pre-defined criteria and process evaluation at the end of the study

#### Secondary outcome measures

There are no secondary outcome measures

#### Overall study start date

01/05/2024

#### Completion date

31/10/2025

# **Eligibility**

#### Key inclusion criteria

- 1. >= 18 vears.
- 2. Experienced acute stroke within previous 7 months.
- 3. Attending first routine post-discharge stroke clinic appointment.
- 4. Able to complete sPCI, either themselves, with assistance for those physically unable to complete for themselves (e.g. have visual or motor problems), or by proxy for those without capacity to consent (i.e. relative/carer completes with their opinion of what the participant's concerns are).
- 5. Able to communicate in English, or have a relative/carer willing to translate trial communications and materials for the participant, or able to use site-specific translation services, if available.

#### Participant type(s)

Patient

#### Age group

Adult

#### Lower age limit

18 Years

#### Sex

Both

#### Target number of participants

Planned Sample Size: 96; UK Sample Size: 96

#### Kev exclusion criteria

1. Lack of capacity to consent themselves and no consultee assent available.

#### Date of first enrolment

25/11/2024

#### Date of final enrolment

01/03/2025

# Locations

#### Countries of recruitment

#### England

#### **United Kingdom**

# Study participating centre Lancashire Teaching Hospitals NHS Foundation Trust

Royal Preston Hospital Sharoe Green Lane Fulwood Preston United Kingdom PR2 9HT

# Study participating centre Royal Cornwall Hospitals NHS Trust

Royal Cornwall Hospital Treliske Truro United Kingdom TR1 3LJ

# Study participating centre The Royal Wolverhampton NHS Trust

New Cross Hospital
Wolverhampton Road
Heath Town
Wolverhampton
United Kingdom
WV10 0QP

# Study participating centre East Lancashire Hospitals NHS Trust

Royal Blackburn Hospital Haslingden Road Blackburn United Kingdom BB2 3HH

# Sponsor information

#### Organisation

University of Central Lancashire

#### Sponsor details

Clayton's Gate
Preston
England
United Kingdom
PR1 2HE
+44 1772895583
irassponsor@uclan.ac.uk

#### Sponsor type

University/education

#### Website

http://www.uclan.ac.uk/

#### **ROR**

https://ror.org/010jbqd54

# Funder(s)

#### Funder type

Government

#### **Funder Name**

NIHR Central Commissioning Facility (CCF)

# **Results and Publications**

#### Publication and dissemination plan

Planned publication in a peer-reviewed journal

#### Intention to publish date

31/10/2026

#### Individual participant data (IPD) sharing plan

The current data sharing plans for this study are unknown and will be available at a later date

#### IPD sharing plan summary

Data sharing statement to be made available at a later date

#### **Study outputs**

Participant information sheet	Control group version 1.1	27/09/2024	18/11/2024	No	Yes
Participant information sheet	Intervention group version 1.1	27/09/2024	18/11/2024	No	Yes