

Efforts to overcome orthostatic hypotension in schizophrenia patients

Submission date 11/10/2025	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 16/10/2025	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 16/10/2025	Condition category Circulatory System	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Orthostatic hypotension (OH) is a common adverse effect of antipsychotic medication that increases the risk of falls and injury among psychiatric patients. Due to limited options for pharmacological adjustment, non-pharmacological interventions are needed. This study aims to evaluate the effectiveness of leg exercise and water consumption interventions in preventing or reducing the incidence of orthostatic hypotension among schizophrenia patients on antipsychotic therapy.

Who can participate?

Participants diagnosed with schizophrenia and identified as experiencing OH in several psychiatric hospitals in Indonesia.

What does the study involve?

Quasi-experimental study with four groups:

- Group A: Control (no intervention)
- Group B: Leg Exercise + Water
- Group C: Leg Exercise only
- Group D: Water only

Each group consisted of 45 participants.

Leg Exercise: Performed in a supine position using a 3 kg foot band. Included hip, knee, and ankle flexions held for 25 seconds each, performed alternately on the right and left leg.

Water Consumption: Adequate fluid intake monitored over 24 hours using labeled water bottles with volume indicators.

Outcome Measurement

Blood pressure was measured in a supine position after 5 minutes of rest and during standing at minutes 1, 3, 5, and 10. Measured daily from Day 1 to Day 5.

Data Analysis was used to assess within-group and between-group differences.

Study Timeline

- Day 0: Screening and group allocation
- Days 1–5: Intervention and daily evaluation
- Day 5: Final assessment

What are the possible benefits and risks of participating?

The benefit of this intervention is to help address recurrent orthostatic hypotension through leg exercises and hydration according to protocol. There are no known downsides or negative effects from this intervention.

Where is the study run from?

Binawan University, Indonesia.

When is the study starting and how long is it expected to run for?

June 2023 to April 2024

Who is funding the study?

Investigator-initiated and funded

Who is the main contact?

Mrs Sari Binawan, sari@binawan.ac.id

Contact information

Type(s)

Principal investigator

Contact name

Mrs Sari Narulita

ORCID ID

<https://orcid.org/0000-0001-9888-4024>

Contact details

Jl. Batu Ampa V No.45A RT 08 RW 05

Kota Jakarta Timur

Indonesia

13520

+62-8128160314

snarulita1706@gmail.com

Type(s)

Scientific

Contact name

Dr Doreen Sumpat

ORCID ID

<https://orcid.org/0000-0001-5507-4275>

Contact details

Jalan UMS
kota kinabalu
Malaysia
88400
+60168112893
doreenes2@ums.edu.my

Type(s)

Public

Contact name

Mrs Sari Narulita

Contact details

Jalab Batu Ampar V No 45A RT 08 RW 05
KOTA JAKARTA TIMUR
Indonesia
13520
+62 8128160314
sari@binawan.ac.id

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Nil known

Study information

Scientific Title

Effectiveness of leg exercise and hydration interventions on orthostatic hypotension among schizophrenia patients in mental hospital

Study objectives

Leg exercise and water consumption are effective in reducing the incidence of orthostatic hypotension

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 17/06/2023, Health Research Ethics Committee of Institute of Research and Community Service University of Catholic Atmajaya (Jalan Jenderal Sudirman 51, Jakarta, 12930, Indonesia; +62-21-5703306; lppm@atmajaya.ac.id), ref: 0006P/III/PPPE.PM.10.05/06/2023

Study design

Quasi-experimental design with four groups

Primary study design

Interventional

Study type(s)

Treatment, Safety

Health condition(s) or problem(s) studied

Patient with schizophrenia with experiment orthostatic hypotension

Interventions

Orthostatic hypotension is defined as a significant drop in blood pressure upon standing, specifically a decrease in systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg within three minutes of standing.

A quasi-experimental study with four groups was conducted in a psychiatric hospital. A total of 180 patients diagnosed with schizophrenia were enrolled and assigned into four equal groups (n = 45): control, leg exercise, water consumption, and combined intervention. Interventions were conducted over five consecutive days. Blood pressure was measured at supine and at 1, 3, 5, and 10 minutes after standing.

Intervention Procedures

Group A (Control): No intervention, follow the routine activities and management nursing care in the wards.

Group B (Leg exercise + water consumption)

Participants received a combination of leg exercise and water consumption interventions:

Water Consumption: Each participant was provided with a water bottle labeled with their name and water volume indicators. Water intake was monitored over 24 hours to ensure adequate hydration. Once adequate hydration was confirmed, the participant proceeded to the leg exercise intervention.

Leg Exercise: The exercise was conducted in a supine position using 3 kg ankle weights (1.5 kg per foot). The exercise protocol involved sequential lower limb movements: hip flexion to 60 degrees, knee flexion to 90 degrees, and ankle dorsiflexion to 90 degrees. Each movement was held for 25 seconds and performed alternately on the right and left legs, completing 10 repetitions per leg.

Before starting the exercise, participants were instructed to lie in a supine position for 5 minutes, and their blood pressure was measured and recorded in that position. Immediately after completing the exercises, participants were instructed to stand up, and their blood pressure was measured at minutes 1, 3, 5, and 10.

Group C (Leg exercise only): Participants received a leg exercise intervention only.

Before starting the exercise, participants were instructed to lie in a supine position for 5 minutes, and their blood pressure was measured and recorded in that position. The exercise was conducted in the supine position using a 3 kg per pair foot band. The exercise protocol included: hip flexion to a 60-degree angle, knee flexion to a 90-degree angle, ankle flexion to a 90-degree angle, done simultaneously and held for 25 seconds. The exercise is performed alternately on the right and left legs. Immediately after completing 10 repetitions in each leg, the participant stood up, and their blood pressure was measured at minutes 1, 3, 5, and 10.

Group D (Water consumption only): Each participant was provided with a labeled water bottle indicating their name and volume markers. Water intake was monitored over 24 hours to ensure adequate hydration, and the results were then documented using a standardized form. Once adequate hydration was confirmed, participants were instructed to lie in a supine position for five minutes. Blood pressure was then measured and recorded in the supine position. Afterwards, participants were instructed to stand up, and their blood pressure was measured at 1, 3, 5, and 10 minutes post-standing.

Inferential statistics were applied to test the effectiveness of the interventions. A repeated measures analysis of variance (RM-ANOVA) was used to evaluate changes in systolic and diastolic blood pressure over five days within and between the intervention and control groups. This approach allowed the detection of time effects, group effects, and interaction effects (time-group). When a significant difference was found, post hoc tests, Bonferroni, were performed to determine which groups differed significantly from one another.

The research was a quasi-experimental (non-randomized) clinical study. Several wards were designated as representatives of the intervention groups. For example, Ward A was for the control group; Wards B and C were for the combination intervention group; Wards D and E were for the leg exercise intervention group; and Wards E and F were for the water consumption group.

Intervention Type

Behavioural

Primary outcome(s)

Orthostatic hypotension: blood pressure was measured using a sphygmomanometer at 1, 3, 5, and 10 minutes post-standing, daily from day 1 to day 5

Key secondary outcome(s)

There are no secondary outcome measures

Completion date

06/04/2024

Eligibility

Key inclusion criteria

1. Diagnosed with schizophrenia
2. Undergoing antipsychotic treatment
3. Identified as having orthostatic hypotension based on blood pressure measurements
4. In a calm, non-aggressive state

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

59 years

Sex

All

Total final enrolment

198

Key exclusion criteria

1. Haemoglobin levels <10 g/d l
2. Estimated glomerular filtration rate (e gfr) <30%
- 3) New York Heart Association (NYHA) class iii or iv heart failure
4. Undergoing dialysis
5. Violent behavior
6. Recent lower limb orthopedics surgery

Date of first enrolment

01/11/2023

Date of final enrolment

30/04/2024

Locations**Countries of recruitment**

Indonesia

Study participating centre**Marzoeki Mahdi Hospital**

Jl. DR. Sumeru No.114, RT.02/RW.01

Bogor

Indonesia

16111

Study participating centre**Soeharto Heerdjan**

Jl. Prof. Dr. Latumeten No.1 RT.1/RW.4

Jakarta

Indonesia

11460

Sponsor information

Organisation

Binawan University

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be stored in a publicly available repository (<https://repository.binawan.ac.id/>)

IPD sharing plan summary

Stored in publicly available repository