

The effect of EMG biofeedback: a randomized controlled clinical trial in patients with shoulder pain

Submission date 03/11/2017	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 24/11/2017	Overall study status Completed	<input checked="" type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 23/04/2021	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

The shoulder injury known commonly as shoulder impingement is the most common disorder of the shoulder and accounts for the major part of neck-shoulder pain in north and mid-western Europe. It is the most common cause of work-related sickness absence, followed by back pain. Neither surgery nor physical treatment has yet found a solution to this problem, as two-thirds of patients are still looking for one or more subsequent treatments. Electromyography (EMG) biofeedback involves using sensors to provide feedback on muscle activity. The aim of this study is to find out whether EMG biofeedback-guided training reduces shoulder pain and improves shoulder function.

Who can participate?

Patients aged 18 to 65 with shoulder impingement

What does the study involve?

Participants are randomly allocated to either the intervention group or the control group. All participants are instructed to perform the same exercises with 2x10 repetitions once a day over 8 weeks. The intervention group receive biofeedback on a computer screen from the muscles the exercises are focused on, while the control group do not receive this biofeedback. Shoulder pain and function are assessed using questionnaires at the start and end of the study.

What are the possible benefits and risks of participating?

The benefits of participation are access to free supervised shoulder muscle exercise training as well as a clinical examination of the shoulders. Participation does not involve any risks for the patients.

Where is the study run from?

University of Southern Denmark (Denmark)

When is the study starting and how long is it expected to run for?

April 2009 to July 2012

Who is funding the study?
University of Southern Denmark (Denmark)

Who is the main contact?
1. Prof. Karen Søgaard
2. Prof. Birgit Juul-Kristensen

Contact information

Type(s)
Scientific

Contact name
Prof Karen Søgaard

ORCID ID
<https://orcid.org/0000-0003-3968-6364>

Contact details
Dep of Sport Sciences and Clinical Biomechanics.
Dep of Clinical Research
University of Southern Denmark, Odense
Campusvej 55
Odense
Denmark
5230

Type(s)
Scientific

Contact name
Prof Birgit Juul-Kristensen

Contact details
Dep of Sport Sciences and Clinical Biomechanics.
University of Southern Denmark, Odense
Campusvej 55
Odense
Denmark
5230

Additional identifiers

Protocol serial number
Danish Association of Rheumatism R62 A985

Study information

Scientific Title

The effect of eight weeks of training with and without EMG biofeedback, on pain and muscle function in patients with shoulder impingement: a randomized controlled clinical trial

Acronym

BIONEX

Study objectives

In impingement patients, an eight-week biofeedback-guided intervention exercise group with EMG feedback will be compared to a control group receiving the same exercises but without EMG biofeedback:

1. Decrease more in shoulder pain measured via self-reported pain (NRS)
2. Increase level of shoulder function (measured via self-reported level of function scores, DASH and OSS)
3. Increase muscular activity in lower trapezius (LT) and serratus anterior (SA), as well as decrease muscle activity in upper trapezius (UT), thereby also decreasing the ratios UT/LT & UT/SA

Ethics approval required

Old ethics approval format

Ethics approval(s)

Regional Ethical Committee of Southern Denmark, 09/11/2009, Project ID: S-20090090

Study design

Single-centre randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Subacromial impingement syndrome (SIS)

Interventions

The intervention program was divided into two periods. The first period was a three-week phase with instruction in different exercises with focus on LT and SA activation. The purpose was to (re) activate the muscles and thereby increase the subjects' awareness of those muscles' position and function. The purpose of the second period (a five-week phase) was to transfer the awareness gained through the first phase into the performance of more functionally complicated exercises. Here, coactivation of the LT and SA in dynamic movements was emphasised in order to master exercises and succeed to the next. Standardised progression regimes according to the subjects' individual pain levels were developed and followed in both first and second period. The subjects were supervised once a week by the same physiotherapist and instructed how to progress and to do the exercises with 2x10 repetitions once a day during the eight weeks of intervention. Also, stretching exercises and ergonomic instructions were given.

All participants in this study received the same exercises. The only difference between the interventions applied to the intervention and control group was the provision of online visual

feedback of muscle activity, shown horizontally (from left to right) on a monitor (biofeedback) visible for both the subject and instructing physiotherapist in the intervention group. The no-EMG control group only received instructions from the physiotherapist based on quality of exercise and if needed, manual/tactile corrections.

Intervention Type

Other

Primary outcome(s)

Self-reported pain assessed using a numeric rating scale (NRS) as part of a questionnaire filled out before commencing physical testing at baseline and follow-up (week 0 and 8):

1. "Pain now" measured on NRS (0-10) at baseline and follow-up testing
 2. "Pain within the last 24 hours" measured on NRS (0-9) at baseline and follow-up testing
 3. "Pain within the last 7 days" measured on NRS (0-9) at baseline and follow-up testing
- Furthermore, the subjects were asked to register their daily pain level in a diary throughout the eight weeks of home exercise. This diary was used to measure the daily pain development (NRS 0-10) and then to measure compliance, as the subjects would register the execution of their home exercises.

Key secondary outcome(s)

1. Self-reported shoulder-scores obtained from questionnaires at baseline and follow-up:
 - 1.1. Disability of the Arm, Shoulder and Hand questionnaire (DASH)
 - 1.2. The Oxford Shoulder Score (OSS)
2. Muscle activity measured by surface electromyography (sEMG) signals from three muscles: upper trapezius (UT), lower trapezius (LT) and serratus anterior (SA). The measurements were carried out according to a standardized experimental procedure, containing a voluntary movement task (described in experimental procedure). During arm elevation, mean relative activity (percentage of maximal voluntary electric activity, %MVE) was measured and muscle activation ratios between the muscles (UT/LT and UT/SA) were calculated. Measured at baseline and follow-up.

Completion date

11/07/2012

Eligibility

Key inclusion criteria

1. At least 30 days with pain/discomfort in the shoulder/neck region within the last year
2. At least two or more positive impingement tests based on the Jobe, Neer; Hawkins and Apprehensions tests
3. 18-65 years old

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

65 years

Sex

All

Total final enrolment

49

Key exclusion criteria

1. Equal or more than 8 in pain/discomfort - measured with Numeric Rating Scale from 0-10 (NRS) - throughout the last 24 hours (on test day)
2. Had more than three regions with pain, for at least 30 days during the last 12 months
3. Had a history of severe shoulder-neck pathology/trauma and/or orthopaedic surgery and/or received anti-inflammatory injections within the last 3 months
4. If they were pregnant (EMG precaution)
5. Any documented life threatening diseases
6. Cardiovascular diseases
7. Rheumatoid arthritis
8. Generalized pain
9. Adverse psychosocial conditions
10. Signs for cervical radiculopathy, i.e. Spurling A test, Involved Cervical Rotation test (less than 60), Neck Distraction test

Date of first enrolment

01/09/2010

Date of final enrolment

03/05/2012

Locations**Countries of recruitment**

Denmark

Study participating centre

University of Southern Denmark

Dept of Sport Sciences and Clinical Biomechanics

Campusvej 55

Odense

Denmark

5230

Sponsor information

Organisation

Forsknings- og Innovationsstyrelsen (National Research Council)

Organisation

Gigtforeningen (Danish Association of Rheumatism)

Organisation

Region Syddanmark

Organisation

Danish Ministry of Higher Education and Science

ROR

<https://ror.org/03ge1nb22>

Funder(s)**Funder type**

University/education

Funder Name

Syddansk Universitet

Alternative Name(s)

University of Southern Denmark, SDU

Funding Body Type

Private sector organisation

Funding Body Subtype

Universities (academic only)

Location

Denmark

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	Participant information sheet	01/10/2019	23/04/2021	Yes	No
Participant information sheet		11/11/2025	11/11/2025	No	Yes
Statistical Analysis Plan				No	No