## The Operative Rib Fixation (ORiF) study

Submission date	Recruitment status  No longer recruiting	Prospectively registered		
29/05/2019		[X] Protocol		
Registration date	Overall study status Completed Condition category	[X] Statistical analysis plan		
30/05/2019		☐ Results		
Last Edited		Individual participant data		
27/02/2025	Injury, Occupational Diseases, Poisoning	[X] Record updated in last year		

### Plain English summary of protocol

Background and study aims

Rib fractures are a common injury. They usually occur as a result of a serious injury, such as those suffered in a road traffic accident or falling from a height. They can also occur in less traumatic accidents, often in elderly people who have fragile bones. Rib fractures are painful and can cause problems with breathing. Lung tissue sits just underneath the ribs and when a fracture occurs, the lungs are also often injured. As a result, rib fractures can lead to problems such as pneumonia, pulmonary effusions (the build-up of fluid in the lungs due to swelling), and some patients can even die as a result of the injury. Most rib fractures are treated without the need for an operation. Doctors use supportive treatments such as pain relief and physiotherapy to help patients recover. Fractures in other bones are usually fixed with an operation that secures the broken bones using metal plates and screws. Recently surgeons have found that some rib fractures can also be fixed in this way. They have also found patients are recovering better with an improvement to their quality of life. However, surgery always carries some risk, especially in patients who have had major injuries. It is not known whether surgical treatment, and its risks, is better than the current non-operative/supportive treatments. The aim of this study is to compare rib fixation with plates and screws to the supportive treatments currently available in the NHS.

## Who can participate?

Patients with three or more rib fractures suitable for surgical repair and one or more of the following: clinical flail chest, breathing difficulty requiring support, and uncontrollable pain using standard modalities

#### What does the study involve?

Participants are randomly allocated to receive either supportive treatments, as is the standard care, or to also undergo an operation to stabilise their rib fractures. Patient outcomes (survival after the injury and quality of life, among other things) and also the cost of treatment to the NHS are measured up until study completion at 12 months. Most of the information is routinely collected in hospitals through an existing system, but some other details are collected directly from patients by specialist research nurses at routine visits to the clinic.

What are the possible benefits and risks of participating?

Both supportive management and operative rib fixation are already routinely carried out in the NHS. However, surgery always carries some risk, especially in patients who have experienced

major injuries. Although there is some evidence to suggest a reduced length of hospital stay and improved quality of life in some patients who undergo operative rib fixation, a benefit cannot be guaranteed to those who take part in this study. The results from the study are likely to benefit future patients with multiple rib fractures, as it is not known whether the surgery, and its risks, are better than the current non-operative, supportive treatments.

Where is the study run from? 15 trauma centres across the UK

When is the study starting and how long is it expected to run for? June 2018 to December 2024

Who is funding the study?
NIHR Health Technology Assessment Programme (UK)

Who is the main contact?

- 1. Prof Benjamin Ollivere, Benjamin.Ollivere@nottingham.ac.uk
- 2. Mr Nicholas Beale, orif@ndorms.ox.ac.uk

## Contact information

#### Type(s)

Scientific

#### Contact name

Prof Benjamin Ollivere

#### **ORCID ID**

https://orcid.org/0000-0002-1410-1756

#### Contact details

School of Medicine
Faculty of Medicine and Health Sciences
Queen's Medical Centre
Derby Road
Nottingham
United Kingdom
NG7 2UH
+44 (0)1158 231115
Benjamin.Ollivere@nottingham.ac.uk

## Type(s)

Public

#### Contact name

Mr Nicholas Beale

#### **ORCID ID**

https://orcid.org/0000-0003-4593-706X

#### Contact details

Surgical Intervention Trials Unit Botnar Research Centre University of Oxford Windmill Road Oxford United Kingdom OX3 7LD +44 (0)1865 613756 orif@ndorms.ox.ac.uk

## Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Protocol serial number

18OR001, HTA 16/61/10

## Study information

#### Scientific Title

A multicentre randomised controlled trial assessing the mortality, quality of life, and costeffectiveness of operative rib fixation plus supportive management versus supportive management alone for patients with multiple rib fractures

#### Acronym

**ORiF** 

#### Study objectives

Operative rib fixation with supportive management is better than supportive management alone.

### Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approved 28/01/2019, South Central – Berkshire Research Ethics Committee (Easthampstead Baptist Church, South Hill Road, Bracknell, RG12 7NS; Tel: +44 (0)0207 104 8360, +44 (0) 2071048046; Email: nrescommittee.southcentral-berkshire@nhs.net), REC ref: 18/SC/0666

### Study design

Pragmatic interventional multi-centre two-arm parallel group randomized controlled trial nested within a population registry

## Primary study design

Interventional

## Study type(s)

Quality of life

#### Health condition(s) or problem(s) studied

Rib fractures

#### **Interventions**

- 1:1 randomisation where patients will receive either:
- 1. Operative rib fixation plus supportive management (intervention)
- 2. Supportive management alone (control)

Minimisation factors: age; polytrauma; mechanical ventilation and study site

#### Intervention Type

Procedure/Surgery

#### Primary outcome(s)

Current primary outcome measure as of 11/12/2023:

All-cause mortality at 12 months

Previous primary outcome measure:

- 1. All-cause mortality at 12 months
- 2. Quality of life measured using the EQ-5D-5L questionnaire at baseline, 30 days, 3 months, 6 months and 12 months

#### Key secondary outcome(s))

Current secondary outcome measures as of 17/03/2023:

- 1. Quality of life measured using the EQ-5D-5L questionnaire at baseline, 30 days, 3 months, 6 months and 12 months
- 2. Patient-reported pain and function measured using the pain Visual Analogue Scale (VAS) and function-related patient questionnaire at 3 months, 6 months and 12 months
- 3. Need for further intervention assessed using patient medical records and Trauma Audit and Research Network (TARN) data, specifically: operative and supportive management details (until discharge), complications (over 12 months), further intervention (over 12 months), ventilator days (until discharge), CT imaging (at baseline) and x-ray imaging (at 6 8 weeks post-discharge for the operative group only)
- 4. Length of stay (LOS) until discharge, assessed using patient medical records and Trauma Audit & Research Network (TARN) data
- 5. Cost-effectiveness measured using a Health Resource Use questionnaire at 6 months and 12 months
- 6. Generalisability of the findings from the randomised trial against the population registry data over 12 months, using a recent statistical approach, using TARN data for both randomised and non-randomised patients

Previous secondary outcome measures:

- 1. Patient-reported pain and function measured using the pain Visual Analogue Scale (VAS) and function-related patient questionnaire at 3 months, 6 months and 12 months
- 2. Need for further intervention assessed using patient medical records and Trauma Audit and Research Network (TARN) data, specifically: operative and supportive management details (until discharge), complications (over 12 months), further intervention (over 12 months), ventilator days (until discharge), CT imaging (at baseline) and x-ray imaging (at 6 8 weeks post-discharge

for the operative group only)

- 3. Length of stay (LOS) until discharge, assessed using patient medical records and Trauma Audit & Research Network (TARN) data
- 4. Cost-effectiveness measured using a Health Resource Use questionnaire at 6 months and 12 months
- 5. Generalisability of the findings from the randomised trial against the population registry data over 12 months, using a recent statistical approach, using TARN data for both randomised and non-randomised patients

### Completion date

31/12/2024

## **Eligibility**

## Key inclusion criteria

Patients who present with multiple (three or more) rib fractures suitable for surgical repair and one or more of the following:

- 1. Clinical flail chest
- 2. Respiratory difficulty requiring respiratory support
- 3. Uncontrollable pain using standard modalities

## Participant type(s)

**Patient** 

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

16 years

#### Sex

All

#### Total final enrolment

467

#### Key exclusion criteria

- 1. Aged under 16 years
- 2. Thoracic injury requiring emergent operative or interventional radiology
- 3. Cannot be operated on within 72 hours as deemed unfit for surgery

#### Added 06/10/2021:

- 4. Unwilling or unable to comply with protocol follow up requirements
- 5. Any other significant disease or condition which, in the opinion of the local research team, may influence the results of the trial or the patient's ability to participate in the trial

#### Date of first enrolment

01/04/2019

## Date of final enrolment 31/07/2024

## Locations

#### Countries of recruitment

**United Kingdom** 

England

Scotland

Wales

## Study participating centre Queen's Medical Centre

Nottingham University Hospitals NHS Trust Derby Road Nottingham United Kingdom NG7 2UH

## Study participating centre The James Cook University Hospital

South Tees Hospitals NHS Foundation Trust Marton Road Middlesbrough United Kingdom TS4 3BW

## Study participating centre Morriston Hospital

Swansea Bay University Health Board Morriston Swansea United Kingdom SA6 6NL

# Study participating centre John Radcliffe Hospital

Oxford University Hospitals NHS Foundation Trust

Headley Way Oxford United Kingdom OX3 9DU

# Study participating centre Royal London Hospital

Barts Health NHS Trust Whitechapel Road London United Kingdom E1 1FR

## Study participating centre Queen Elizabeth Hospital

University Hospitals Birmingham NHS Foundation Trust Mindelsohn Way Birmingham United Kingdom B15 2GW

## Study participating centre Derriford Hospital

Derriford Road Derriford Plymouth United Kingdom PL6 8DH

## Study participating centre Kings College Hospital

Denmark Hill London United Kingdom SE5 9RS

## Study participating centre Aintree University Hospital

Liverpool University Hospitals NHS Foundation Trust Lower Lane Liverpool United Kingdom L9 7AL

## Study participating centre Bristol Royal Infirmary

Marlborough Street Bristol United Kingdom BS2 8HW

# Study participating centre Southmead Hospital

Southmead Road Westbury-on-trym Bristol United Kingdom BS10 5NB

# Study participating centre Hull Royal Infirmary

Anlaby Road Hull United Kingdom HU3 2JZ

## Study participating centre Glenfield Hospital

Groby Road Leicester United Kingdom LE3 9QP

## Study participating centre St. Mary's Hospital

Imperial College Healthcare NHS Trust Praed Street London United Kingdom W2 1NY

## Study participating centre Manchester Royal Royal Infirmary

Cobbett House Oxford Road Manchester United Kingdom M13 9WL

# Study participating centre University Hospital of Wales

Heath Park Cardiff United Kingdom CF14 4XW

## Study participating centre Aberdeen Royal Infirmary

Grampian Health Board Summerfield House 2 Eday Road Aberdeen United Kingdom AB15 6RE

## Sponsor information

### Organisation

Nottingham University Hospitals NHS Trust

#### **ROR**

https://ror.org/05y3qh794

## Funder(s)

## Funder type

Government

#### **Funder Name**

Health Technology Assessment Programme

#### Alternative Name(s)

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

### Funding Body Type

Government organisation

### **Funding Body Subtype**

National government

#### Location

United Kingdom

## **Results and Publications**

### Individual participant data (IPD) sharing plan

Current IPD sharing statement as of 17/03/2023:

On completion of the study, and with appropriate participant consent, fully anonymised data may be shared with other organisations at the behest of the funder. All requests for the use of the data from the ORiF study will be approved by the CI, TMG and where necessary the TSC. A data request form should be completed detailing the decision as to whether the request is accepted. In cases where individual site data is requested, only summary data would be provided with caveats for dissemination, to emphasise that trial data should be interpreted as a whole.

### Previous IPD sharing statement:

The data sharing plans for the current study are unknown and will be made available at a later date.

### IPD sharing plan summary

Available on request

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023		No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Protocol file	version 6.0	10/07/2023	16/12/2024	No	No
Statistical Analysis Plan	version 1.0	20/12/2024	27/02/2025	No	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes