

# Characterization of problems associated with vision impairment and utility of the current interventions

<b>Submission date</b> 18/12/2018	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 21/01/2019	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 14/11/2022	<b>Condition category</b> Eye Diseases	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

The burden caused by vision loss can be reduced using targeted vision rehabilitation. Vision rehabilitation is intended to promote the best possible functional ability and the individual's ability to maintain their skills. However, there is still a lack of knowledge regarding the "value-for-money" of vision rehabilitation not only for people with impaired vision but also for the society. The aim of this study is to estimate the cost-effectiveness (value-for-money) of vision rehabilitation by measuring the consequences of rehabilitation for visual ability (visual ability is defined as the overall ability to perform activities that depends on vision).

### Who can participate?

Adults diagnosed with diabetic retinopathy (DR) or age-related macular degeneration (AMD) with visual acuity between 0.4 - 0.1 (decimal scale) in the better-seeing eye. Participants must live in the community (not in nursing homes or other types of institution).

### What does the study involve?

The study intervention consists of dispensing new glasses and magnifiers, defined as "vision aids" with minimal training. Instructions are provided in clinical settings with emphasis on reading tasks. Participants randomly allocated to the immediate intervention group receive the device in the first visit. Participants in the delayed intervention group receive the device after 3 months, in the second visit. All participants have a follow-up assessment at 3 month and at 9 months after receiving vision aids. During visits participants also need to response to questionnaires and to perform vision tests (none of them are invasive).

### What are the possible benefits and risks of participating?

This study brings benefits to all participants because they will receive a basic vision rehabilitation intervention which includes new glasses and/or magnifiers with minimal training and instructions provided in clinical settings with emphasis on reading tasks. With the intervention procedures the patients won't be exposed to any risk.

Where is the study run from?

The study is being run by the University of Minho and takes place in a single centre, a public hospital in Barcelos, Portugal

When is the study starting and how long is it expected to run for?

September 2016 to December 2021

Who is funding the study?

The ophthalmic lenses and part of the magnifiers are supported by Essilor Portugal

Who is the main contact?

1. Antonio Filipe Macedo

a.macedo@ucl.ac.uk; antonio.macedo@lnu.se

2. Laura Moreno

PhD student in University of Minho, Braga, Portugal

## Contact information

### Type(s)

Scientific

### Contact name

Dr Antonio Filipe Macedo

### ORCID ID

<https://orcid.org/0000-0003-3436-2010>

### Contact details

Linnaeus University

Department of Medicine and Optometry

Smålandsgatan 26 A

Kalmar

Sweden

39234

+351 (0)768805972

a.macedo@ucl.ac.uk

### Type(s)

Scientific

### Contact name

Miss Laura Moreno

### ORCID ID

<https://orcid.org/0000-0001-6623-9151>

### Contact details

University of Minho

Departamento de Fisica

Campus de Gualtar

Braga

Portugal  
4710 - 057

## **Additional identifiers**

**Protocol serial number**  
N/A

## **Study information**

### **Scientific Title**

Cost-effectiveness of basic vision rehabilitation services (the basic VRS-effect study): a cross-over randomised controlled trial

### **Acronym**

The basic VRS-effect study

### **Study objectives**

The aim of the trial is to compare the effects and costs of “usual low vision care” with a “basic vision rehabilitation service (basic VRS)” on self-reported visual ability.

Hypothesis: A basic low vision intervention, consisting of prescription of new glasses, magnifiers and instructions, improves visual ability more than usual care.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Life Sciences and Health of the University of Minho, 02/05/2017, ref: SECVS 147/2016

CEUM/SECVS – Secretariado  
Att: Isabel Monteiro  
Universidade do Minho – Reitoria  
Campus de Gualtar, CII, Piso 3  
4710-057 Braga  
Tel: +351 (0)253 601 700  
Email: secvs@reitoria.uminho.pt

Registered in the Portuguese data protection authority, ref: 7012/ 2017

### **Study design**

Randomised cross-over controlled trial

### **Primary study design**

Interventional

### **Study type(s)**

Other

### **Health condition(s) or problem(s) studied**

Vision impairment caused by diabetic retinopathy or age-related macular degeneration

## **Interventions**

To perform a cost-effectiveness analysis (CEA) of basic VRS the trialists designed a randomised, cross-over controlled trial (AB/BA structure), comparing the effects and costs of "usual low vision care" (which is not yet systematised in Portugal) with a "basic-VRS intervention" on self-reported visual ability.

Participants will be allocated to the immediate intervention group (IMI), which is represented by group A or to a delayed intervention group (DEI) represented by group B. Participants are allocated to the IMI or DEI to create the best possible match between the groups according to age and visual acuity. The sequence of randomization is done blocks of 4 to guarantee two participants for each group using four similar envelopes. Each envelope will contain the code of the group, two envelopes for IMI and two for DEI group. The study takes place in a single-centre.

The intervention consists of dispensing new glasses and magnifiers, defined as "vision aids", with minimal training and instructions provided in clinical settings with emphasis on reading tasks. Participants allocated to the immediate intervention group will receive the device in the first visit. Participants in the delayed intervention group will receive the device after 3 months, in the second visit. All participants will have a follow-up assessment at 3 months and at 9 months after receiving vision aids.

## **Intervention Type**

Device

## **Primary outcome(s)**

Measured at baseline, 12 weeks and 36 weeks:

1. Visual ability, measured using Massof Activity Inventory (AI)
2. Health-related quality of life, measured using EQ-5D-5L

## **Key secondary outcome(s)**

Measured at baseline, 12 weeks and 36 weeks:

1. Reading ability, measured using Portuguese version of Minnesota Low-Vision Reading Test (MNread test)
2. Depression and anxiety, measured using Hospital Anxiety and Depression Scale (HADS)

## **Completion date**

31/12/2021

## **Eligibility**

### **Key inclusion criteria**

1. Visual acuity between 0.4 - 1.0 logMAR (0.4 - 0.1 decimal) in the better-seeing eye
2. Primary diagnosis (cause of vision loss) diabetic retinopathy or age-related macular degeneration
3. 18 years of age or older
4. Resident in the community (not in nursing homes or other types of institution)

### **Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Sex**

All

**Total final enrolment**

47

**Key exclusion criteria**

1. Cognitive impairment based on scores on mini-mental examination test
2. Communication problems due to, for example, hearing impairment, or unable to speak Portuguese
3. Unable to read in Portuguese due to low level of education
4. Unable to attend the requested appointments at the study setting

**Date of first enrolment**

03/03/2017

**Date of final enrolment**

30/03/2020

**Locations****Countries of recruitment**

Portugal

**Study participating centre**

University of Minho

Campus de Gualtar

Braga

Portugal

4710-057

**Sponsor information****Organisation**

University of Minho

ROR

<https://ror.org/037wpkx04>

## Funder(s)

### Funder type

Other

### Funder Name

Investigator initiated and funded

## Results and Publications

### Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date.

### IPD sharing plan summary

Data sharing statement to be made available at a later date

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>		11/11/2022	14/11/2022	Yes	No
<a href="#">Protocol article</a>	protocol	01/05/2020	14/02/2020	Yes	No
<a href="#">Interim results article</a>	Association between social support and depression and anxiety	03/03/2021	07/09/2021	Yes	No
<a href="#">Interim results article</a>	Association between depression and anxiety and health-related quality of life	10/02/2022	25/07/2022	Yes	No