

Bone proteins in the treatment of periodontal diseases

Submission date 28/10/2022	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 06/11/2022	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 29/12/2023	Condition category Oral Health	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

The advent of recently developed techniques and the availability of biomaterials, such as the use of enamel matrix protein derivatives (EMD), recombinant human platelet-derived growth factor (rhPDGF), and hyaluronic acid (HA), have provided both clinicians and researchers with sophisticated therapeutic options for the management of Grade II furcation-type defects (bone loss, usually as the result of periodontal disease, affecting the base of the root trunk of a tooth where several roots meet). The use of these biomaterials in combination with either bone graft substitutes or membranes aimed at guided tissue regeneration has been considered an effective modality when treating infrabony defects (at the base of the tooth pocket relative to the bone crest) and Grade-II furcation-type lesions. The use of biodegradable membranes to promote periodontal regeneration has been reported. The properties of the biodegradable membrane should permit selective repopulation of periodontal ligament (PDL) cells onto the root surfaces, followed by degradation only when this process is complete. Polylactic acid/polyglycolic acid (PLA/PGA) is a synthetic bioabsorbable membrane made from a co-polymer of lactide and glycolide. The time of degradation of PLA/PGA membrane (Polyglactin-910) was found to be 30-50 days and has been reported to facilitate periodontal regeneration. Hence, the current study protocol was designed to assess the performance of rhBMP-2 with PLA/PGA membrane through the reduction of horizontal and vertical defect depth with simultaneous gain in clinical attachment level to achieve periodontal tissue regeneration in Grade-II furcation-type defects.

Who can participate?

Patients with Grade II furcation defects

What does the study involve?

All patients undergoing treatment for Grade-II furcation-type defects will receive bioabsorbable PLA/PGA membrane to achieve periodontal tissue regeneration. Teeth randomly assigned to the test group will also receive an application of rhBMP-2. Periodontal regeneration will be assessed in both groups using dental probes at baseline, 3 months and 6 months postoperatively.

What are the possible benefits and risks of participating?

Possible benefits of participating include:

1. Active participation in their periodontal health care

2. Gaining access to new treatments before they are widely available
3. A team of dentists who will provide them with regular and attentive dental therapy
4. Contribution to dental research to help others

Possible risks of participating include:

1. A small chance of minor discomfort that will only last a short time
2. The study may require more time and attention than standard treatment would, including visits to the study site

Where is the study run from?

Sharad Pawar Dental College & Hospital (India)

When is the study starting and how long is it expected to run for?

March 2016 to September 2019

Who is funding the study?

Sharad Pawar Dental College & Hospital (India)

Who is the main contact?

Dr Kiran Kumar Ganji

dr.kiran.ganji@jodent.org

Contact information

Type(s)

Principal Investigator

Contact name

Dr Kiran Kumar Ganji

ORCID ID

<https://orcid.org/0000-0002-3178-9513>

Contact details

Department of Periodontics & Implantology

Sharad Pawar Dental College & Hospital

Sawangi Meghe

Wardha

Nagpur

India

442001

+919225755282

dr.kiran.ganji@jodent.org

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

Nil known

Study information

Scientific Title

Evaluation of the effectiveness of recombinant human bone morphogenic protein 2 (rhBMP-2) in combination with bioresorbable membrane (PLA/PGA) for periodontal regeneration in grade II furcation defects: a clinical study

Study objectives

Recombinant human bone morphogenic protein 2 (rhBMP-2) along with a PLA/PGA membrane, does not contribute to gain in clinical attachment levels for Grade-II furcation type defects (if compared to the sole use of a PLA/PGA membrane), and it does not lead to differing results regarding the vertical and horizontal defect depths, or the extent of gingival recessions.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 02/09/2016, Institutional Ethics Committee (Datta Meghe Institute of Medical Sciences, Nagpur, India; +917152 287701; medical_ivda@sancharnet.in), ref: DMIMS(DU)/IEC /1527

Study design

Single-centre longitudinal case-control 6-month study

Primary study design

Observational

Secondary study design

Case-control study

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use contact details to request a participant information sheet

Health condition(s) or problem(s) studied

Regenerative periodontal therapy in the management of furcation defects caused due to periodontal diseases

Interventions

After achieving adequate anesthesia, intra-crevicular incisions were given on the buccal surfaces of teeth, including one tooth adjacent (mesial and distal) to access the furcation defect for treatment. The incisions were made as inter-proximally as possible, for the preservation of the inter-dental papilla, to obtain primary closure of the wound. The releasing incisions were given when needed with the aim to achieve adequate accessibility, and also gain coronal displacement of the periodontal flap. A mucoperiosteal flap was then reflected using a periosteal elevator. The pocket epithelium was curetted out carefully. The furcation defect was debrided, and the roof of the defect and the exposed surfaces of the root were scaled and planed using hand instruments as well as ultrasonic instruments.

Following these intraoperative measurements, the confirmed defects were randomly assigned to test (n = 12) and control groups (n = 12) by using a computer-generated blocked randomization list (Randlist, DatInf), thus referring to the pre-sealed envelopes. In the test group, the flap was pre-sutured without placing a knot in order to allow for rapid wound closure following the placement of the PLA/PGA membrane with rhBMP-2. A bioabsorbable PLA/PGA membrane (BioMesh-s, Samyang Biopharma) was trimmed in order to conceal the lesion, extending approximately 2 mm beyond the edges of the furcation defect. In the control group (PLA/PGA alone), the membrane was trimmed and placed over the defect in a manner similar to the test group, except without the application of rhBMP-2.

Intervention Type

Procedure/Surgery

Primary outcome measure

1. Relative clinical attachment level measured using Williams graduated periodontal probe and numerical values were recorded into a Microsoft Excel spreadsheet at baseline, 3 months and 6 months postoperatively
2. Vertical probing depth measured using Williams graduated periodontal probe and numerical values were recorded into a Microsoft Excel spreadsheet at baseline, 3 months and 6 months postoperatively
3. Horizontal probing depth measured using Nabers graduated furcation probe and numerical values were recorded into a Microsoft Excel spreadsheet at baseline, 3 months and 6 months postoperatively
4. Relative gingival margin level measured using Williams graduated periodontal probe and numerical values were recorded in a Microsoft Excel spreadsheet at baseline, 3 months and 6 months postoperatively

Secondary outcome measures

Periodontal regeneration was assessed by clinical attachment gain, vertical probing depth reduction and horizontal probing depth reduction which were measured using Williams graduated periodontal probe and numerical values were recorded into a Microsoft Excel spreadsheet at baseline, 3 months and 6 months postoperatively

Overall study start date

01/03/2016

Completion date

26/09/2019

Eligibility

Key inclusion criteria

1. Grade II furcation defects
2. Furcation defects present in mandibular 1st molars
3. Systemically healthy
4. Furcation defects present on the buccal surface

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

24

Total final enrolment

24

Key exclusion criteria

1. Grade II furcation defects in second mandibular molars
2. Patients with past history of periodontal therapy
3. Former and current smokers

Date of first enrolment

26/01/2019

Date of final enrolment

24/02/2019

Locations**Countries of recruitment**

India

Study participating centre

Sharad Pawar Dental College & Hospital

Sawangi Meghe

Wardha

Nagpur

India

442001

Sponsor information

Organisation

Sharad Pawar Dental College & Hospital

Sponsor details

Sawangi Meghe

Wardha

India

442001

+917152287714

medical_ivda@sancharnet.in

Sponsor type

University/education

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Sharad Pawar Dental College & Hospital

Results and Publications

Publication and dissemination plan

Planned publication in an high-impact peer-reviewed journal

Intention to publish date

15/06/2023

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Dr Kiran Kumar Ganji, dr.kiran.ganji@jodent.org. Qualitative and quantitative data will be shared. Written consent was obtained from the participants. Personal data has been encrypted by the identifiers to maintain privacy and also to remove the personally identifiable information of the participants.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		10/02/2023	29/12/2023	Yes	No