

INFORM - surgery for hip infection project

Submission date 28/01/2015	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 29/01/2015	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 13/02/2024	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Hip replacement surgery is common, with over 86,000 cases in England and Wales in 2012. After surgery about 1% patients develop a deep infection in their artificial hip, called a prosthetic joint infection (PJI). If left untreated PJI can result in severe pain, disability and death. When an infection is found, there are two types of surgical treatment: a 1-stage revision involves removing the joint, thoroughly cleaning the infected area and implanting a new joint immediately; a 2-stage revision involves removing the joint, and delaying re-implantation for 3 to 6 months whilst treating with antibiotics. Both treatments are widely used but we don't know which has the best long-term outcomes for patients. The aim of this study is to find out whether there is any difference in patient-reported outcomes (pain, stiffness or physical function) after 1-stage or 2-stage revision hip surgery for PJI.

Who can participate?

Patients that have a prosthetic joint infection that needs surgery.

What does the study involve?

Participants are randomly allocated into one of two groups. Those in group 1 have 1-stage revision surgery. Those in group 2 have 2-stage revision surgery. All participants are then followed up for the next 18 months. They are asked to complete questionnaires every 3 months, to assess their hip pain and function, quality of life and any costs incurred during their treatment. They attend two appointments with a research nurse to complete questionnaires and a walk test. A small number of participants and surgeons are also interviewed about their experiences of treatment and participation in the trial. Participants who decide against the randomised trial are asked to join a monitoring group. This group have treatment-as-usual, and then complete questionnaires at home. The monitoring group provide important information on the wider population of patients with hip PJI, including patient reported outcomes, control of infection and adverse events.

What are the possible benefits and risks of participating?

Not provided at time of registration

Where is the study run from?

14 hospitals in the UK and 3 in Sweden (UK)

When is the study starting and how long is it expected to run for?
January 2015 to February 2020

Who is funding the study?
National Institute for Health Research (UK)

Who is the main contact?
Mr Simon Strange

Contact information

Type(s)
Scientific

Contact name
Mr Simon Strange

Contact details
INFORM Programme Manager
Musculoskeletal Research Unit
Level 1, Learning and Research
Southmead Hospital
Bristol
United Kingdom
BS10 5NB
+44 (0)7789777053
simon.strange1@nhs.net

Additional identifiers

Protocol serial number
18159

Study information

Scientific Title
A multicentre randomised trial to compare 1-stage with 2-stage revision surgery for prosthetic hip joint infection

Study objectives
The aim of this study is to determine if there is any difference in patient-reported outcomes (pain, stiffness or physical function) after 1-stage or 2-stage revision hip surgery for prosthetic joint infection (PJI).

Ethics approval required
Old ethics approval format

Ethics approval(s)
NRES Committee South-West - Frenchay, 31/12/2014, ref: 14/SW/1166

Study design

Randomised; Interventional and Observational; Design type: Not specified, Treatment, Qualitative

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Topic: Surgery; Subtopic: Surgery; Disease: All Surgery

Interventions

Revision hip surgery

1-stage or 2-stage revision for infected prosthetic hip joint; Follow Up Length: 18 month(s)

Intervention Type

Other

Primary outcome(s)

Patient-reported hip pain, stiffness, function; Timepoint(s): WOMAC at 18 months post-randomisation

Key secondary outcome(s))

1. Quality of life, measured using Euroqol EQ-5D-5L and HOOS
2. All hip-related complications (including continuing and reinfection) recorded throughout study period
3. Post-op pain, measured using the Brief Pain Inventory
4. Patient-reported hip function, measured using the Oxford Hip Score
5. Objective measure of hip function, measured using the 20 metre timed walk test
6. Depression and anxiety, measured using the Hospital Anxiety and Depression Score
7. Cost-effectiveness of interventions, measured using health resource use

Completion date

28/02/2020

Eligibility

Key inclusion criteria

1. Age 18 or above
 2. A clinical diagnosis of prosthetic hip joint infection
 3. Require revision surgery (either 1-stage or 2-stage) to treat prosthetic hip joint infection, in the opinion of the treating consultant orthopaedic surgeon
- Target Gender: Male & Female ; Lower Age Limit 18 years

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

177

Key exclusion criteria

1. Unable or unwilling to undergo either 1-stage or 2-stage revision surgery
2. Lack capacity to give written informed consent

Date of first enrolment

04/03/2015

Date of final enrolment

31/08/2018

Locations

Countries of recruitment

United Kingdom

England

Sweden

Study participating centre

North Bristol NHS Trust

Trust Headquarters

Beckspool Road

Frenchay

Bristol

United Kingdom

B16 1JE

Sponsor information

Organisation

North Bristol NHS Trust

ROR

<https://ror.org/036x6gt55>

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

Participant level data will be made available at the data.bris (University of Bristol Research Data repository) beginning 1 year following publication of the trial results. Access to data will 'controlled' and request forms can be found at <http://www.bristol.ac.uk/staff/researchers/data/accessing-research-data/>

IPD sharing plan summary

Stored in repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		31/10/2022	01/11/2022	Yes	No
Results article		01/11/2022	13/02/2024	Yes	No
Protocol article	protocol	17/02/2016		Yes	No

[HRA research summary](#)

28/06/2023 No No

[Other publications](#)

What are patients' preferences for revision surgery after periprosthetic joint infection? A discrete choice experiment

21/01/2020 13/02/2024 Yes No

[Other publications](#)

qualitative study

03/08/2020 13/02/2024 Yes No

[Participant information sheet](#)

Participant information sheet

11/11/2025 11/11/2025 No Yes