# Prediction of adverse pregnancy outcomes following induction of labour

Submission date	Recruitment status  No longer recruiting	[X] Prospectively registered		
26/01/2016		Protocol		
Registration date	Overall study status Completed Condition category	Statistical analysis plan		
02/02/2016		Results		
Last Edited		Individual participant data		
16/05/2016	Pregnancy and Childbirth	<ul><li>Record updated in last year</li></ul>		

## Plain English summary of protocol

Background and study aims

Although most labours begin naturally, however around one third of women need help to get started. In these cases, a healthcare professional will often "induce" labour. Although the aim of induction is to achieve a vaginal delivery, it is not always successful and some women end up having a caesarean section (an operation where the child is delivered through a cut in the abdomen). Currently, there are no effective methods to predict which women will not respond favourably to induction of labor or how long this process is likely to take. The aim of this study is to develop techniques to help predict whether induction will result in vaginal delivery or caesarean section.

#### Who can participate?

Women aged 18 or over with a single baby pregnancy, who have been scheduled for an induction of labour at Medway Maritime Hospital.

#### What does the study involve?

All women at Medway Maritime Hospital receive a routine clinical assessment in the preinduction clinic, which involves an ultrasound scan of the baby, digital vaginal examination and a
blood test. Participants are also offered an ultrasound scan to measure the length and dilation
of the neck of the womb (cervix) and to find out how low the baby's head is in the pelvis. For
those agreeing to take part in the study, a swab of the vagina and another blood sample is taken
to be tested in the laboratory for natural indicators (biochemical markers) as well as receiving an
ultrasound scan which can help to predict the result of the induction. The amount of women
who go into labour following induction are then recorded and compared to the results of
prediction from the biochemical markers and ultrasound scan to find out how well the tests are
able to predict the results of the induction.

What are the possible benefits and risks of participating? There are no direct benefits or risks for participants taking part in this study.

Where is the study run from? Medway Maritime Hospital (UK) When is the study starting and how long is it expected to run for? November 2015 to June 2017

Who is funding the study? Fetal Medicine Foundation (UK)

Who is the main contact?

- 1. Dr Alexander Frick (public)
- 2. Mr Ranjit Akolekar (scientific)

# Contact information

## Type(s)

Public

#### Contact name

Dr Alexander Frick

#### **ORCID ID**

https://orcid.org/0000-0001-8013-3679

#### Contact details

Fetal Medicine Research Institute 16-20 Windsor walk London United Kingdom SE5 8BB

# Type(s)

Scientific

#### Contact name

Prof Ranjit Akolekar

#### Contact details

Medway Maritime Hospital Fetal Medicine Unit Windmill Road Gillingham United Kingdom ME7 5NY

# Additional identifiers

Protocol serial number

2.0

# Study information

Scientific Title

Prediction Of adverse Pregnancy outcomes following INduction of labour in singleton pregnancies: A single centre observational study

#### Acronym

**POPIN** 

#### **Study objectives**

The aim of this study is to assess if a combination of ultrasound findings and biochemical markers from vaginal secretions and maternal blood can better predict vaginal delivery than traditional methods such as the Bishop Score.

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

London – Dulwich Research Ethics Committee, 31/03/2016, ref: 16/LO/0367

#### Study design

Single-centre prospective observational cohort study

#### Primary study design

Observational

#### Study type(s)

Screening

## Health condition(s) or problem(s) studied

Pregnancy and childbirth

#### **Interventions**

Women attending a pre-induction clinic who are participating in the study will have a vaginal swab taken to test levels of quantitative fetal fibronectin and placental alpha-macroglobulin-1 prior to induction of labour. They will also have a transperineal ultrasound (TPUS) to measure the angle of progression, head-perineum distance and sonographic assessment of cervical dilatation. The results from these assessments will be combined with maternal characteristics to predict mode of delivery.

Patients will then be followed throughout the induction and labour process, with the transperineal ultrasound assessments repeated on a 4 hourly basis to compare the effectiveness of ultrasound with clinical vaginal examination.

Women will be followed until delivery to determine the accuracy of the pre-induction test and intrapartum ultrasound assessments. The total duration of observation will vary from depending on the length of induction and labour, but will typically be 1-3 days. There will be no need for follow up after delivery.

## Intervention Type

Other

# Primary outcome(s)

Detection rate and false positive rate for predicting vaginal delivery following induction of labour is measured using a combination of maternal characteristics, ultrasound findings, and biochemical markers from a pre-induction clinic at the time of delivery.

#### Key secondary outcome(s))

- 1. Detection rate for caesarean section for failure to progress is measured using a combination of maternal characteristics, ultrasound findings, and biochemical markers from a pre-induction clinic at the time of delivery
- 2. Detection rate for caesarean section for fetal distress is measured using a combination of maternal characteristics, ultrasound findings, and biochemical markers from a pre-induction clinic at the time of delivery
- 3. Detection rate for maternal haemorrhage greater than 1500 ml is measured using a combination of maternal characteristics, ultrasound findings, and biochemical markers from a pre-induction clinic at the time of delivery
- 4. Detection rate for neonatal intensive care admission is measured using a combination of maternal characteristics, ultrasound findings, and biochemical markers from a pre-induction clinic at the time of delivery
- 5. Effectiveness of intrapartum ultrasound in predicting vaginal delivery as compared to traditional vaginal examination is determined at the time of delivery

#### Completion date

01/10/2017

# Eligibility

#### Key inclusion criteria

- 1. Female
- 2. Viable singleton pregnancy
- 3. Aged 18 years or over
- 4. Cephalic presentation
- 5. Attending pre-induction of labour clinic
- 6. Informed written consent

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Adult

## Lower age limit

18 years

#### Sex

Female

#### Key exclusion criteria

- 1. Multiple pregnancies
- 2. Women with fetal demise
- 3. Women less than 18 years
- 4. Women who are unconscious or severely ill, those with learning difficulties, and serious mental illness
- 5. Malpresentation

#### Date of first enrolment

01/03/2016

#### Date of final enrolment

01/03/2017

# Locations

#### Countries of recruitment

United Kingdom

**England** 

## Study participating centre Medway Maritime Hospital

Medway NHS Foundation Trust Windmill Road Gillingham United Kingdom ME7 5NY

# Sponsor information

## Organisation

Medway NHS Foundation Trust

#### **ROR**

https://ror.org/01apxt611

# Funder(s)

# Funder type

Charity

#### **Funder Name**

#### Fetal Medicine Foundation

## Alternative Name(s)

**FMF** 

# **Funding Body Type**

Private sector organisation

# Funding Body Subtype

Trusts, charities, foundations (both public and private)

#### Location

**United Kingdom** 

# **Results and Publications**

Individual participant data (IPD) sharing plan

# IPD sharing plan summary

Available on request

# **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes